

International Journal of Statistics and Applied Mathematics

ISSN: 2456-1452
Maths 2018; 3(3): 226-230
© 2018 Stats & Maths
www.mathsjournal.com
Received: 18-03-2018
Accepted: 22-04-2018

B Amudhambigai
Assistant Professor of
Mathematics, Department of
Mathematics Sri Sarada College
for Women Salem, Tamil Nadu,
India

G Vasuki
Associate Professor of
Mathematics, Department of
Mathematics Sri Sarada College
for Women Salem, Tamil Nadu,
India

N Santhiya
Research Scholar, Department of
Mathematics Sri Sarada College
for Women Salem, Tamil Nadu,
India

Correspondence
B Amudhambigai
Assistant Professor of
Mathematics, Department of
Mathematics Sri Sarada College
for Women Salem, Tamil Nadu,
India

A view on rb-convergence in multiset topological spaces

B Amudhambigai, G Vasuki and N Santhiya

Abstract

This paper deals with the concepts of regular b-closed M-sets and regular b-continuous M-set functions are studied with necessary examples. Also the concepts of rb-convergence to a point, rb-accumulates to a point, rb-S-closed M-spaces are studied and some interesting properties are discussed. Finally, sequence of rb-convergence and rb-Hausdorff M-spaces are discussed.

Keywords: regular b-closed M-sets, regular b-continuous M-set functions, rb-convergence, rb-accumulates, rb-S-closed M-spaces

2010 AMS Subject Classification: 54A05, 54A10, 54A20

1. Introduction

The notion of M-topological space and the concept of open M-sets are introduced by Girish and sunil Jacob John ^[4]. Moreover, topologies on multisets can be associated to IC-bags or nk-bags introduced by Chakrabarthy ^[3] (Chakrabarty, 2000; Chakrabarty and Despi, 2007) with the help of rough set theory. The association of rough set theory and topologies on multisets through bags with interval counts (Chakrabarty and Despi, 2007) can be used to develop theoretical study of covering based rough sets with respect to universe as multisets. Nagaveni and Narmadha ^[9, 10] introduced the notions of regular b-closed (rb-closed) sets in topological space. Regular open sets and b-open sets have been introduced and investigated by Stone ^[12] and Andrijević ^[1]. Generalized α -closed sets have been introduced and investigated Maki, Devi and Balachandran ^[8].

2. Preliminaries

Definition 2.1 ^[4]: Let $M \in [X]_w$ and $\tau \subseteq P^*(M)$. Then τ is called a Multiset topology of M if τ satisfies the following properties.

1. The M-set M and the empty M-set ϕ are in τ .
2. The M-set union of the elements of any sub collection of τ is in τ .
3. The M-set intersection of the elements of any finite sub collection of τ is in τ .

Definition 2.2 ^[4] Given a subM-set A of an M-topological space M in $[X]_w$, the interior of A is denoted by $\text{Int}(A)$ and it is defined as follows: $\text{Int}(A) = \bigcup \{G \subseteq M: G \text{ is an open M-set and } G \subseteq A\}$ and $C_{\text{Int}(A)}(x) = \max \{C_G(x) : G \subseteq A, G \in \tau\}$.

Definition 2.3 ^[4]: Given a subM-set A of an M-topological space M in $[X]_w$, the closure of A is denoted by $\text{Cl}(A)$ and is defined as follows: $\text{Cl}(A) = \bigcap \{K \subseteq M: K \text{ is a closed M-set and } A \subseteq K\}$ and $C_{\text{Cl}(A)}(x) = \min \{C_K(x) : A \subseteq K, K \in \tau^c\}$.

3. On rb-Closed M-Sets And rb-Continuous M-Set Functions

Definition 3.1: Let (M, τ) be an M-topological space in $[X]^w$. Any subM-set A of M is called a regular open M-set if $A = \text{int}(\text{cl}(A))$ with $C_A(x) = C_{\text{int}(\text{cl}(A))}(x)$, for all $x \in X$.

Definition 3.2: Let (M, τ) be an M -topological space in $[X]^W$. Any sub M -set A of M is called a regular closed M -set if $A = \text{cl}(\text{int}(A))$ with $C_A(x) = C_{\text{cl}(\text{int}(A))}(x)$, for all $x \in X$.

Definition 3.3: Let (M, τ) be an M -topological space in $[X]^W$ and A be any sub M -set of M . The M -set union of all regular open M -sets of (M, τ) contained in A is called the regular interior of A and is denoted by $\text{rint}(A)$ i.e. $\text{rint}(A) = \bigcup \{G \subseteq M: \text{each } G \text{ is a regular open } M\text{-set and } G \subseteq A\}$ with $C_{\text{rint}(A)}(x) = \text{Max} \{C_G(x): G \subseteq A, \text{ each } G \text{ is a regular open } M\text{-set}\}$, for all $x \in X$.

Definition 3.4: Let (M, τ) be an M -topological space in $[X]^W$ and A be any sub M -set of M . The M -set intersection of all regular closed M -sets of (M, τ) containing A is called the regular closure of A and it is denoted by $\text{rcl}(A)$ i.e. $\text{rcl}(A) = \bigcap \{K \subseteq M: \text{each } K \text{ is a regular closed } M\text{-set and } A \subseteq K\}$ with $C_{\text{rcl}(A)}(x) = \text{Min} \{C_K(x): A \subseteq K, \text{ each } K \text{ is a regular closed } M\text{-set}\}$, for all $x \in X$.

Definition 3.5: Let (M, τ) be an M -topological space in $[X]^W$. Any sub M -set A of M is called a b -open M -set if $A \subseteq \text{int}(\text{cl}(A)) \cup \text{cl}(\text{int}(A))$ with $C_A(x) \leq \text{Max} \{C_{\text{int}(\text{cl}(A))}(x), C_{\text{cl}(\text{int}(A))}(x)\}$, for all $x \in X$.

Definition 3.6: Let (M, τ) be an M -topological space in $[X]^W$. Any sub M -set A of M is called a regular b -closed (rb-closed) M -set if $\text{rcl}(A) \subseteq U$ with $C_{\text{rcl}(A)}(x) \leq C_U(x)$, whenever $A \subseteq U$ with $C_A(x) \leq C_U(x)$ and U is a b -open M -set in (M, τ) , for all $x \in X$. The complement of a regular b -closed M -set is called a regular b -open (rb-open) M -set.

Definition 3.7: Let (M, τ) be an M -topological space in $[X]^W$. For any sub M -set A of M , the rb -interior of A is defined as the M -set union of all rb -open M -sets of (M, τ) contained in A and is denoted by $\text{rb-int}(A)$ i.e. $\text{rb-int}(A) = \bigcup \{G \subseteq M: \text{each } G \text{ is a } rb\text{-open } M\text{-set and } G \subseteq A\}$ with $C_{\text{rb-int}(A)}(x) = \text{Max} \{C_G(x): G \subseteq A, \text{ each } G \text{ is a } rb\text{-open } M\text{-set}\}$, for all $x \in X$.

Definition 2.8: Let (M, τ) be an M -topological space in $[X]^W$. For any sub M -set A of M , the rb -closure of A is defined as the M -set intersection of all rb -closed M -sets of (M, τ) containing A and is denoted by $\text{rb-cl}(A)$ i.e. $\text{rb-cl}(A) = \bigcap \{K \subseteq M: \text{each } K \text{ is a } rb\text{-closed } M\text{-set and } A \subseteq K\}$ with $C_{\text{rb-cl}(A)}(x) = \text{Min} \{C_K(x): A \subseteq K, \text{ each } K \text{ is a } rb\text{-closed } M\text{-set}\}$, for all $x \in X$.

Definition 3.9: Let (M, τ) be an M -topological space in $[X]^W$. Any sub M -set A of M is called an α -open M -set if $A \subseteq \text{int}(\text{cl}(\text{int}(A)))$ with $C_A(x) \leq C_{\text{int}(\text{cl}(\text{int}(A)))}(x)$, for all $x \in X$. The complement of an α -open M -set is called an α -closed M -set.

Definition 3.10: Let (M, τ) be an M -topological space in $[X]^W$ and A be any sub M -set of M . The M -set intersection of all α -closed M -sets containing A is called the α -closure of A and is denoted by $\alpha\text{cl}(A)$. Similarly, the M -set union of all α -open M -sets of (M, τ) contained in A is called the α -interior of A and is denoted by $\alpha\text{int}(A)$ i.e. $\alpha\text{cl}(A) = \bigcap \{K \subseteq M: \text{each } K \text{ is an } \alpha\text{-closed } M\text{-set and } A \subseteq K\}$ with $C_{\alpha\text{cl}(A)}(x) = \text{Min} \{C_K(x): A \subseteq K, \text{ each } K \text{ is an } \alpha\text{-closed } M\text{-set}\}$, for all $x \in X$. Similarly, $\alpha\text{int}(A) = \bigcup \{K \subseteq M: \text{each } K \text{ is an } \alpha\text{-open } M\text{-set and } A \supseteq K\}$ with $C_{\alpha\text{int}(A)}(x) = \text{Max} \{C_K(x): A \supseteq K, \text{ each } K \text{ is an } \alpha\text{-open } M\text{-set}\}$, for all $x \in X$.

Definition 3.11: Let (M, τ) be an M -topological space in $[X]^W$. Any sub M -set A of M is called a generalized α -closed (ga -closed) M -set if $\alpha\text{cl}(A) \subseteq U$ with $C_{\alpha\text{cl}(A)}(x) \leq C_U(x)$, whenever $A \subseteq U$ with $C_A(x) \leq C_U(x)$, and U is an α -open M -set in (M, τ) , for all $x \in X$.

Proposition 3.1.1: Every regular closed M -set is a rb -closed M -set.

Remark 3.1.2: the converse of the above Proposition 2.1.1 need not be true as shown in the Example 2.1.12.

Example 3.1.3 Let $X = \{a, b, c\}$, $W = 2$. Let $M = \{1/a, 2/b, 1/c\}$. Let $\tau = \{M, \phi, \{1/a\}, \{2/b\}, \{1/a, 2/b\}\}$. Then $\tau^c = \{\phi, M, \{2/b, 1/c\}, \{1/a, 1/c\}, \{1/c\}\}$. Clearly, τ is an M -topology and the ordered pair (M, τ) is an M -topological space. Also, the collection of the regular closed M -sets of (M, τ) is $\{M, \phi, \{2/b, 1/c\}, \{1/a, 1/c\}\}$ and the collection of the b -open M -sets of (M, τ) is $\{\phi, M, \{1/a\}, \{2/b\}, \{1/b\}, \{1/a, 2/b\}, \{2/b, 1/c\}, \{1/a, 1/c\}, \{1/b, 1/a\}, \{1/a, 1/b, 1/c\}\}$.

Let $A = \{1/c\}$ be a sub M -set of M and $A \subset U = \{1/a, 1/c\}$ is the b -open M -set in (M, τ) . Then $\text{rcl}(A) = \{1/c\} \subset U$. Therefore, $\text{rcl}(A) \subset U$ with $C_{\text{rcl}(A)}(x) < C_U(x)$, for all $x \in X$. whenever $A \subseteq U$ and U is b -open M -set with $C_A(x) \leq C_U(x)$, for all $x \in X$. Hence, A is a rb -closed M -set. But $\text{int}(A) = \phi$ and $\text{cl}(\text{int}(A)) = \phi$. Thus, $A \neq \text{cl}(\text{int}(A))$ with $C_A(c) = 1 \neq 0 = C_{\text{cl}(\text{int}(A))}(c)$, $C_A(a) = C_{\text{cl}(\text{int}(A))}(a) = 0$ and $C_A(c) = C_{\text{cl}(\text{int}(A))}(c) = 0$, for all $x \in X$. Hence, A is not a regular closed M -set.

Proposition 3.1.4: Every regular closed M -set is a ga -closed M -set.

Remark 3.1.5: The converse of the Proposition 2.1.2 need not be true as shown in the Example 2.1.13

Example 3.1.6: Let $X = \{a, b, c\}$, $W = 2$. Let $M = \{1/a, 2/b, 1/c\}$. Let $\tau = \{\phi, M, \{1/a\}, \{1/a, 2/b\}, \{2/b\}\}$. Then $\tau^c = \{M, \phi, \{2/b, 1/c\}, \{1/c\}, \{1/a, 1/c\}\}$. Clearly, τ is an M -topology and the ordered pair (M, τ) is an M -topological space. Now the collection of the α -open M -sets of (M, τ) is $\{M, \phi, \{1/a\}, \{1/a, 2/b\}, \{2/b\}\}$ and α -closed M -sets of (M, τ) is $\{M, \phi, \{2/b, 1/c\}, \{1/c\}, \{1/a, 1/c\}\}$.

Let $A = \{1/b, 1/c\}$ be a sub M -set of M . Then $A \subset U = M$ is the α -open M -set in (M, τ) . Then $\alpha\text{cl}(A) = \{2/b, 1/c\} \subset U$. Therefore, $\alpha\text{cl}(A) \subset U$ with $C_{\alpha\text{cl}(A)}(x) < C_U(x)$, for all $x \in X$ whenever $A \subset U$ and U is an α -open M -set. Hence, A is a ga -closed M -set. But $\text{int}(A) = \phi$ and $\text{cl}(\text{int}(A)) = \phi$. Thus $A \neq \text{cl}(\text{int}(A))$ with $C_A(c) = 1 \neq 0 = C_{\text{cl}(\text{int}(A))}(c)$, $C_A(a) = C_{\text{cl}(\text{int}(A))}(a) = 0$ and $C_A(b) = C_{\text{cl}(\text{int}(A))}(b) = 0$. Hence, A is not a regular closed M -set.

Proposition 3.1.7: Every rb -closed M -set is a ga -closed M -set.


Remark 3.1.8: The converse of the Proposition 2.1.3 need not be true as shown in the Example 2.1.14

Example 3.1.9: Let $X = \{a, b, c\}$, $W = 2$. Let $M = \{1/a, 2/b, 1/c\}$. Let $\tau = \{\phi, M, \{1/a\}, \{1/a, 2/b\}, \{2/b\}\}$. Then $\tau^c = \{M, \phi, \{2/b, 1/c\}, \{1/c\}, \{1/a, 1/c\}\}$. Now τ is an M -topology and the ordered pair (M, τ) is an M -topological space. Now the collection of the regular-closed M -sets of (M, τ) is $\{\phi, M, \{1/c, 2/b\}, \{1/c, 1/a\}\}$, the collection of the α -closed M -sets of (M, τ) is $\{M, \phi, \{2/b, 1/c\}, \{1/c\}, \{1/a, 1/c\}\}$.

$1/c\}$ }, the collection of the b-open M-sets of (M, τ) is $\{\phi, M, \{1/a\}, \{2/b\}, \{1/b\}, \{1/a, 2/b\}, \{2/b, 1/c\}, \{1/a, 1/c\}, \{1/b, 1/a\}, \{1/a, 1/b, 1/c\}\}$ and the collection of the α -open M-sets of (M, τ) is $\{M, \phi, \{2/a\}, \{1/b\}, \{2/a, 1/b\}\}$. Let $A = \{1/a, 1/b, 1/c\}$ be a sub M-set of M and $A \subset U = M$ is the α -open M-set in (M, τ) . Then $\alpha cl(A) = M = U$. Therefore, $\alpha cl(A) = U$ with $C_{\alpha cl(A)}(x) = C_U(x)$ whenever $A \subset U$ and U is α -open M-set with $C_A(x) < C_U(x)$, for all $x \in X$. Hence, A is a ga -closed M-set. But $A = U = \{1/a, 1/b, 1/c\}$ is the b-open M-set. Then $rcl(A) = M \not\subset U$ with $C_{rcl(A)}(x) \not\leq C_U(x)$, for all $x \in X$, whenever $A = U$ and U is b-open M-set. Hence, A is not a rb-closed M-set.

Remark 3.1.10: From the above discussions, the following implications hold: In the following diagram by $A \longrightarrow B$ means A implies B

$A \longleftarrow B$ does not imply A .



3.2 Regular b-continuous M-set functions

Definition 3.2.1 Let (M, τ) and (N, σ) be any two M-topological spaces. Any M-set function $f: (M, \tau) \rightarrow (N, \sigma)$ is said to be a continuous M-set function if $f^{-1}(A)$ is closed (resp. open) M-set in (M, τ) , for every closed (resp. open) M-set A in (N, σ) .

Definition 3.2.2 Let (M, τ) and (N, σ) be any two M-topological spaces. Any M-set function $f: (M, \tau) \rightarrow (N, \sigma)$ is said to be a regular continuous M-set function if $f^{-1}(A)$ is regular closed (resp. regular open) M-set in (M, τ) , for every closed (resp. open) M-set A in (N, σ) .

Definition 3.2.3 Let (M, τ) and (N, σ) be any two M-topological spaces. Any M-set function $f: (M, \tau) \rightarrow (N, \sigma)$ is said to be a regular b-continuous (rb-continuous) M-set function if $f^{-1}(A)$ is rb-closed (res. rb-open) M-set in (M, τ) , for every closed (res. open) M-set A in (N, σ) .

Definition 3.2.4 Let (M, τ) and (N, σ) be any two M-topological spaces. Any M-set function $f: (M, \tau) \rightarrow (N, \sigma)$ is said to be a generalized α -continuous (ga -continuous) M-set function if $f^{-1}(A)$ is ga -closed (res. ga -open) M-set in (M, τ) , for every closed (res. open) M-set A in (N, σ) .

Proposition 3.2.5 Let (M, τ) and (N, σ) be any two M-topological spaces in $[X]^w$. For any M-set function $f: (M, \tau) \rightarrow (N, \sigma)$, the following statements are equivalent:

- f is a rb-continuous M-set function.
- $f(rb-cl(A)) \subseteq cl(f(A))$, for each $A \subseteq M$ with $C_{f(rb-cl(A))}(x) \leq C_{cl(f(A))}(x)$, for all $x \in X$
- $rb-cl(f(B)) \subseteq f^{-1}(cl(B))$, for each $B \subseteq N$ with $C_{rb-cl(f^{-1}(B))}(x) \leq C_{f^{-1}(cl(B))}(x)$, for all $x \in X$.
- $f^{-1}(int(B)) \subseteq rb-int(f^{-1}(B))$, for each $B \subseteq N$ with
- $C_{(f^{-1}(int(B)))}(x) \leq C_{rb-int(f^{-1}(B))}(x)$, for all $x \in X$.

Proposition 3.2.6: Every regular continuous M-set function is a continuous M-set function.

Remark 3.2.7: The converse of the above Proposition need not be true as shown in the Example 2.2.5

Example 3.2.8: Let $X = \{a, b, c\}$, $W_1 = 2$, $Y = \{x, y, z\}$, and $W_2 = 1$. Let $M = \{1/a, 2/b, 1/c\}$ and $N = \{1/x, 1/y, 1/z\}$ be two M-sets. Let $\tau = \{M, \phi, \{1/a\}, \{2/b\}, \{1/a, 2/b\}\}$ and $\sigma = \{N, \phi, \{1/x, 1/y\}\}$ be the M-topologies on M and N respectively. Then (M, τ) and (N, σ) are M-topological spaces. Now the collection of the closed M-sets of (N, σ) is $\{\phi, N, \{1/z\}\}$ and the collection of the regular closed M-sets of (M, τ) is $\{M, \phi, \{2/b, 1/c\}, \{1/a, 1/c\}\}$ and the collection of the closed M-sets of (M, τ) is $\{\phi, M, \{2/b, 1/c\}, \{1/a, 1/c\}, \{1/c\}\}$.

Let M-set function $f: (M, \tau) \rightarrow (N, \sigma)$ be defined by $f = \{(1/a, 1/x)/1, (2/b, 1/y)/2, (1/c, 1/z)/1\}$. For the closed M-set $A = \{1/z\}$ in (N, σ) , $f^{-1}(A) = \{1/c\}$ is an closed M-set in (M, τ) . Trivially $f^{-1}(\phi) = \phi$ and $f^{-1}(N) = M$ are closed M-sets in (M, τ) . But $\{1/c\}$ is not a regular closed M-set in (M, τ) . Hence, every continuous M-set function need not be a regular continuous M-set function.

Proposition 3.2.9: Every regular continuous M-set function is a rb-continuous M-set function.

Remark 3.2.10: The converse of the Proposition 2.2.3 need not be true as shown in following Example.

Example 3.2.11: Let $X = \{a, b, c\}$, $W_1 = 2$, $Y = \{x, y, z\}$, and $W_2 = 1$. Let $M = \{1/a, 2/b, 1/c\}$ and $N = \{1/x, 1/y, 2/z\}$ be two M-sets. Let $\tau = \{M, \phi, \{1/a\}, \{2/b\}, \{1/a, 2/b\}\}$ and $\sigma = \{N, \phi, \{1/x, 1/y\}\}$ be two M-topologies on M and N respectively. Then (M, τ) and (N, σ) are M-topological spaces. Now the collection of the regular closed M-sets of (M, τ) is $\{M, \phi, \{2/b, 1/c\}, \{1/a, 1/c\}\}$ and the collection of the rb-closed M-sets of (M, τ) is $\{M, \phi, \{1/c, 2/b\}, \{1/a, 1/c\}, \{1/c\}\}$ and the collection of the closed M-sets of (N, σ) is $\{\phi, N, \{2/z\}\}$.

Let M-set function $f: (M, \tau) \rightarrow (N, \sigma)$ be defined by $f = \{(1/a, 1/x)/1, (2/b, 1/y)/2, (1/c, 2/z)/2\}$. For the closed M-set $A = \{2/z\}$ in (N, σ) , $f^{-1}(A) = \{1/c\}$ is an rb-closed M-set in (M, τ) . Trivially $f^{-1}(\phi) = \phi$ and $f^{-1}(N) = M$ are rb-closed M-sets in (M, τ) . But $\{1/c\}$ is not a regular closed M-set in (M, τ) . Hence, every rb-continuous M-set function need not be a regular continuous M-set function.

Proposition 3.2.12: Every rb-continuous M-set function is a ga -continuous M-set function.

Remark 3.2.13: The converse of the Proposition 2.2.4 need not be true as shown in the Example 2.2.7

Example 3.2.14: Let $X = \{a, b, c\}$, $W_1 = 2$, $Y = \{x, y, z\}$ and $W_2 = 2$. Let $M = \{1/a, 2/b, 1/c\}$ and $N = \{2/x, 1/y, 1/z\}$ be two M-sets. Let $\tau = \{\phi, M, \{1/a\}, \{2/b\}, \{1/a, 2/b\}\}$ and $\sigma = \{N, \phi, \{2/x\}\}$ be the M-topologies on M and N respectively. Then (M, τ) and (N, σ) are M-topological spaces. Now the collection of the closed M-sets of (N, σ) is $\{\phi, N, \{1/y, 1/z\}\}$ and the collection of the rb-closed M-sets of (M, τ) is $\{M, \phi, \{1/c\}, \{2/b, 1/c\}, \{1/c, 1/a\}\}$ and the collection of the ga -closed M-sets of (M, τ) is $\{M, \phi, \{1/c\}, \{1/b, 1/c\}, \{1/c, 2/b\}, \{1/c, 1/a\}, \{1/a, 1/b, 1/c\}\}$.

Let M-set function $f: (M, \tau) \rightarrow (N, \sigma)$ be defined by $f = \{(1/a, 2/x)/2, (1/b, 1/y)/2, (1/c, 1/z)/1\}$. For the closed M-set $A = \{1/y, 1/z\}$ in (N, σ) , $f^{-1}(A) = \{1/b, 1/c\}$ is a ga -closed M-set in (M, τ) . Trivially $f^{-1}(\phi) = \phi$ and $f^{-1}(N) = M$ are ga -

closed M-sets in (M, τ) , but $\{1/b, 1/c\}$ is not a rb-closed in (M, τ) . Hence every $g\alpha$ -continuous M-set function need not be a rb-continuous M-set function.

Proposition 3.2.15: Every regular continuous M-set function is $g\alpha$ -continuous M-set function.


Remark 3.2.16: The converse of the Proposition 2.2.5 need not be true as shown in the Example 2.2.6

Example 3.2.17: Let $X = \{a, b, c\}$, $W_1 = 2$, $Y = \{x, y, z\}$ and $W_2 = 2$. Let $M = \{1/a, 2/b, 1/c\}$ and $N = \{2/x, 1/y, 1/z\}$ be two M-sets. Let $\tau = \{\phi, M, \{1/a\}, \{2/b\}, \{1/a, 2/b\}\}$ and $\sigma = \{N, \phi, \{2/x, 1/y\}\}$ be the M-topologies on M and N respectively. Then (M, τ) and (N, σ) are M-topological spaces. Now the collection of the closed M-sets of (N, σ) is $\{\phi, N, \{1/z\}\}$ and the collection of the regular closed M-sets of (M, τ) is $\{M, \phi, \{2/b, 1/c\}, \{1/c, 1/a\}\}$ and the collection of the $g\alpha$ -closed M-sets of (M, τ) is $\{M, \phi, \{2/b, 1/c\}, \{1/c\}, \{1/b, 1/c\}, \{1/c, 1/a\}, \{1/a, 1/b, 1/c\}\}$.

Let M-set function $f: (M, \tau) \rightarrow (N, \sigma)$ be defined by $f = \{(1/a, 2/x)/2, (2/b, 1/y)/2, (1/c, 1/z)/1\}$. For the closed M-set $A = \{1/z\}$ in (N, σ) , $f^{-1}(A) = \{1/c\}$ is an $g\alpha$ -closed M-set in (M, τ) . Trivially $f^{-1}(\phi) = \phi$ and $f^{-1}(N) = M$ are $g\alpha$ -closed M-sets in (M, τ) , but $\{1/c\}$ is not a regular closed in (M, τ) . Hence, every $g\alpha$ -continuous M-set function need not be a regular continuous M-set function.

Remark 3.2.18: From the above discussions, the following implications hold: In the following diagram by A B means A implies B

$A \not\Rightarrow B$ does not imply A.



4. On rb-Convergence in Multiset Topological Spaces

In this chapter, the concepts of rb-convergence to a point, rb-accumulates to a point, rb-S-closed M-spaces are studied and some interesting properties are discussed. Finally, sequence of rb-convergence and rb-Hausdorff M-spaces are discussed.

Definition 4.1: A filter is a non-empty collection F of subM-sets of an M-topological space (M, τ) such that

1. $\phi \notin F$;
2. If $A \in F$ and $B \supseteq A$ with $C_B(x) \geq C_A(x)$, for all $x \in X$, then $B \in F$ and
3. If $A \in F$ and $B \in F$, then $A \cap B \in F$ with $C_{A \cap B}(x) \leq C_F(x)$, for all $x \in X$.

Definition 4.2: Let (M, τ) be an M-topological space. A filter base of a filter F is a sub collection B of M such that for every $F \in F$ there exists $B \in B$ with $B \subseteq F$ with $C_B(x) \leq C_F(x)$, for

all $x \in X$. Such a subcollection B satisfies the following conditions:

1. $\phi \notin B$
2. For every $B_1, B_2 \in B$ there exists $B_3 \in B$ with $B_3 \subseteq B_1 \cap B_2$ with $C_{B_3}(x) \leq C_{B_1}(x) \cap C_{B_2}(x)$, for all $x \in X$.

Definition 4.3: Let F be a filter on B. Then F is a maximal filter if every filter including F coincides with F

Definition 4.4: Let (M, τ) be an M-topological space. A filter base $B = \{ \{ A_\alpha \}_{\alpha \in J} \}$, where J is an indexed set and $A_\alpha \in M$ } rb-converges to a point $x_0 \in {}^m M$ if for each rb-open M-set V containing m/x_0 there exists an $A_\alpha \in B$ such that $A_\alpha \subseteq \text{rb-cl}(V)$ with $C_{A_\alpha}(x) \leq C_{\text{rb-cl}(V)}(x)$, for all $x \in X$.

Definition 4.5: A filter base $B = \{ \{ A_\alpha \}_{\alpha \in J} \}$, where J is indexed set and $A_\alpha \in M$ } rb-accumulates to a point $x_0 \in {}^m M$ if for each rb-open M-set V containing m/x_0 and $A_\alpha \in B$, $A_\alpha \cap \text{rbcl}(V) \neq \phi$ with $\text{Min} \{ C_{A_\alpha}(x), C_{\text{rb-cl}(V)}(x) \} \neq 0$, for all $x \in X$.

Definition 4.6: Let (M, τ) be an M-topological space. A collection $\mathcal{A}$ of sub M-sets of a space M is said to cover M, or to be a covering of M, if the union of the elements of $\mathcal{A}$ is equal to M. It is called an open covering of M if its elements are open sub M-sets of M.

Definition 4.7: Let (M, τ) be an M-topological space. An M-topological space $(M, \square)$ is said to be rb-S-closed if for every cover $\{ V_\alpha \}_{\alpha \in I}$, I is an indexed set } of (M, τ) by rb-open M-sets of M, there exists a finite subM-set I_0 of I such that $\bigcup \{ \text{rb-cl}(V_\alpha) \}_{\alpha \in I_0} = M$ with $\text{Max} \{ C_{\text{rb-cl}(V_\alpha)}(x) \}_{\alpha \in I_0} = C_M(x)$, for all $x \in X$.

Definition 4.8: Let (M, τ) be an M-topological space. A subM-set K of an M-topological space $(M, \square)$ is said to be rb-closed relative to M if for any cover $\{ U_i \}_{i \in I}$, I is an indexed set } of K by rb-open M-sets, there exists a finite subM-set I_0 of I such that $K \subseteq \bigcup \{ \text{rb-cl}(U_i) \}_{i \in I_0}$ with $C_K(x) \leq \text{Max} \{ C_{\text{rb-cl}(U_i)}(x) \}_{i \in I_0}$, for all $x \in X$.

Remark 4.9: The family of all rb-open (res, rb-closed) M-sets of (M, τ) is denoted by $\text{RBO}(X)$ (res, $\text{RBC}(X)$). The family of all rb-open M-sets of (M, τ) containing a point $x \in {}^m M$ is denoted by $\text{RBO}(M, m/x)$.

Remark 4.10: A filter base B is rb-convergent to a point $x \in {}^m M$ if and only if B contains the collection $\{ \text{rbcl}(U) : U \in \text{RBO}(M, m/x) \}$.

Proposition 4.11: For an M-topological space (M, τ) , the following statements are equivalent:

1. M is an rb-S-closed M-space.
2. For each family of rb-closed M-sets $\{ F_\alpha \}_{\alpha \in I}$, where I is an indexed set, such that $\bigcap (F_\alpha) = \phi$ with $\text{Min} \{ C_{F_\alpha}(x), \alpha \in I \} = 0$, for all $x \in X$, there exists a finite subfamily $\{ F_{\alpha_i} \}_{i=1}^n$ such that $\bigcap_{i=1}^n \text{rb-int}(F_{\alpha_i}) = \phi$ with $\text{Min} \{ C_{\text{rb-int}(F_{\alpha_i})}(x), i = 1, 2, \dots, n \} = 0$, for all $x \in X$.
3. Each filter base $B = \{ A_\alpha \}_{\alpha \in I}$ with $C_B(x) = C_{\{ A_\alpha \}_{\alpha \in I}}(x)$, for all $x \in X$, rb-accumulates to some point $x_0 \in {}^m M$.
4. Each maximum filter base B Rb-converges.

Proposition 4.12: A subM-set A of a M -topological space (M, τ) is rb -closed relative to M if and only if every filter base B on M with $A \in B$ rb -converges to a point in A .

Definition 4.13: Let (M, τ) be an M -topological space. A sequence in a M -set M is a function $f: \mathbb{N} \rightarrow M$. Then $m/x_n = f(n)$ for all $n \in \mathbb{N}$, where $\mathbb{N}$ is positive integer and denoted the sequence as $(m/x_n)_{n \in \mathbb{N}}$.

Definition 4.14: Let (M, τ) be an M -topological space, let $x \in {}^m M$ and $N \subseteq M$. Then N is said to be a rb -neighborhood of m/x if there is an rb -open M -set V in τ such that $x \in {}^m V$ and $C_V(y) \subseteq C_N(y)$, for all $y \neq x$.

I.e. a rb -neighborhood of m/x in M means any rb -open M -set containing m/x . Here m/x is said to be an interior point of N .

Definition 4.15: Let (M, τ) be an M -topological space. A sequence $m/x_1, m/x_2, m/x_3, \dots$ of points in M is said to be rb -converges to a point $x \in {}^m M$ if for each rb -neighborhood U of m/x , there is a positive integer $\mathbb{N}$ such that $x_n \in {}^m U$, for all $n \geq \mathbb{N}$. It is denoted by $(m/x_n) \rightarrow m/x$.

Definition 4.16: An M -topological space (M, τ) is said to be a rb -Hausdorff M -space, if each pair of distinct points m/x and m/y in M , there exist rb -open M -sets U and V containing m/x and m/y respectively such that $U \cap V = \emptyset$ with $C_U \cap C_V(x) = \emptyset$, for all $x \in X$.

Definition 4.17: Let (M, τ) be an M -topological space. A rb -countable basis at a point m/x is a countable M -set $\{U_n \mid n \in \mathbb{N}\}$ of rb -neighbourhood of m/x , such that for any rb -neighbourhood V of m/x there is an $n \in \mathbb{N}$ such that $U_n \subseteq V$ with $C_{U_n}(x) \subseteq C_V(x)$, for all $x \in X$.

An M -topological space (M, τ) is said to be first rb -countable if every point has a rb -countable basis.

Proposition 4.3 Let (M, τ) be an M -topological space. Let M be a rb -Hausdorff space and $x_n \in {}^m M$ a rb -convergent sequence. Then the limit $\lim_{n \rightarrow \infty} m/x_n$ is unique.

5. Acknowledgement

The authors would like to express their sincere gratitude to the referees for their valuable suggestions and comments for the betterment of this paper.

6. References

1. Andrijevic D. On b -open sets, mat. Vesnik. 1996; 48:59-64.
2. Beceren Y. Almost α -irresolute functions, bull. Calcutta Math. Soc. 2000; 92:213-218.
3. Chakrabarty, Kankana. Ioan Despi. n^k -bags. Int. J Intell. Syst. 2007s; 22(2):223-236.
4. Girish KP. Sunil Jacob John. On Multiset Topologies, 2007s.
5. Computer and Mathematics with Applications, 2012, 37-52
6. James Munkres R. Topology, Prentice Hall of India Private Limited, New Delhi.
7. Jeyanthi V, Janaki C. On π gr-continuous functions, International Journal of Engineering Research and Applications, Jan-Feb. 2013; 3(1):1861-1870.
8. John Kelly L. General Topology, Spring India Private Limited.

9. Maki H, Devi R, Balachandran K. Associated topologies of generalized α -closed sets and α -generalized closed sets, Mem, Fac. Sci, Kochi Univ. Ser. A Math. 1994; 15:51-63.
10. Nagaveni N, Narmadha A. A note on regular b -open sets in topological spaces, Proc, UGC and DST-Curie sponsored International Conference on Mathematics and its Applications-A New Wave, ICMANW, Dec. 2011, 93-100, 21-22.
11. Nagaveni N, Narmadha A. On regular b -closed sets in topological spaces, Heber International Conference on Applications of Mathematics and Statistics, HICAMS, 2012, 5-7, 81-87.
12. Narmadha A, Nagaveni N, Noiri T. On Regular b -open sets in Topological spaces, Int. Journal of Math. Analysis. 2013; 7(19):937-948.
13. Stone MH. Applications of the theory of Boolean rings to general topology", Trans. Math. Soc. 1937; 41:375-381.
14. Thomson T. S-closed spaces, Proc. Amer. Math. Soc. 1976; 60:335-338.
15. Thomas Wright. Convergence of sequences and Nets in Metric and Topological space, July 1, 2013, 1-6.