

# International Journal of Statistics and Applied Mathematics

ISSN: 2456-1452  
Maths 2018; 3(6): 19-21  
© 2018 Stats & Maths  
www.mathsjournal.com  
Received: 05-09-2018  
Accepted: 11-10-2018

**Showkat Ahmad Rather**  
Research Scholar Ashmuji  
Kulgam, Jammu and Kashmir,  
India

**Riyaz Ahmad Shah**  
HOD Department of  
Mathematics GDC Kulgam,  
Jammu and Kashmir, India

## Concircular curvature tensor in Kenmotsu manifold

**Showkat Ahmad Rather and Riyaz Ahmad Shah**

### Abstract

In this paper we study Concircular curvature tensor in Kenmotsu manifolds. We consider curvature tensor, scalar curvature in a Kenmotsu manifold admitting concircular curvature tensor and prove the condition for concircular curvature tensor to be shrinking, steady and expanding.

**Keywords:** Curvature tensor, scalar curvature, Ricci tensor, Einstein, Kenmotsu manifold

### Introduction

The Concircular curvature  $\tilde{C}$  on Kenmostu manifolds of dimensional  $n$  is defined by

$$\tilde{C}(X, Y)Z = R(X, Y)Z - \frac{r}{n(n-1)} [g(Y, Z)X - g(X, Z)Y] \quad (1.1)$$

for any vector fields  $X, Y, Z$  where  $R$  is the curvature tensor and  $r$  is the scalar curvature.

In an  $n$ - dimensional Kenmotsu manifold, we have

$$\eta(R(X, Y)Z) = g(X, Z) \eta(Y) - g(Y, Z) \eta(X), \quad (1.2)$$

$$R(X, Y)\xi = \eta(X)Y - \eta(Y)X, \quad (1.3)$$

$$R(\xi, X)Y = \eta(Y)X - g(X, Y)\xi, \quad (1.4)$$

$$R(\xi, X)\xi = X - \eta(X)\xi, \quad (1.5)$$

where  $R$  is the Riemannian curvature tensor.

Taking  $Z = \xi$  in equation (1.1) and using equation (1.3), we get

$$\begin{aligned} \tilde{C}(X, Y)\xi &= R(X, Y)\xi - \frac{r}{n(n-1)} [g(Y, \xi)X - g(X, \xi)Y] \\ \Rightarrow \tilde{C}(X, Y)\xi &= [1 + \frac{r}{n(n-1)}] [\eta(X)Y - \eta(Y)X]. \end{aligned} \quad (1.6)$$

Taking inner product of equation (1.1) with  $\xi$ , we get

$$\eta(\tilde{C}(X, Y)Z) = \eta(R(X, Y)Z) - \frac{r}{n(n-1)} [g(Y, Z)\eta(X) - g(X, Z)\eta(Y)]. \quad (1.7)$$

Using equation (1.2) in above equation, we get

$$\eta(\tilde{C}(X, Y)Z) = [1 + \frac{r}{n(n-1)}] [g(X, Z)\eta(Y) - g(Y, Z)\eta(X)] \quad (1.8)$$

### 2.2 Preliminaries

An  $n$ - dimensional differential manifold  $M$  is said to be an almost contact metric manifold [3]. If it admits an almost contact metric structure  $(\phi, \xi, \eta, g)$  consisting of a tensor field  $\phi$  of type

### Correspondence

**Showkat Ahmad Rather**  
Research Scholar Ashmuji  
Kulgam, Jammu and Kashmir,  
India

(1, 1) a vector field  $\xi$ , a 1-form  $\eta$ , and a Riemannian metric  $g$  compatible with  $(\phi, \xi, \eta, g)$  satisfying.

$$\phi^2 = -I + \otimes \xi, \eta(\xi) = 1, \quad (2.1)$$

$$\eta \circ \phi = 0, \phi \xi = 0 \quad (2.2)$$

$$g(\phi X, \phi Y) = g(X, Y) - \eta(X)\eta(Y), g(X, \xi) = \eta(X), \quad (2.3)$$

for all vector fields  $X, Y$  on  $M$ .

An almost contact metric manifold  $M(\phi, \xi, \eta, g)$  is said to be Kenmotsu manifold if

$$(\nabla_X \phi)Y = g(\phi X, Y)\xi - \eta(Y)\phi X. \quad (2.4)$$

From equation (2.4), we have

$$\nabla_X \xi = X - \eta(X)\xi, \quad (2.5)$$

where  $\nabla$  denotes the Riemannian connection of  $g$ .

Let  $(g, V, \lambda)$  be a Ricci soliton is an  $n$ -dimensional Kenmotsu manifold  $M$ .

From equation (2.5), we have

$$(\mathcal{L}_\xi g)(X, Y) = 2[g(X, Y) - \eta(X)\eta(Y)], \quad (2.6)$$

From equations (2.1) and (2.6), we have

$$S(X, Y) = -(\lambda + 1)g(X, Y) + \eta(X)\eta(Y), \quad (2.7)$$

$$S(\phi X, \phi Y) = S(X, Y) + \lambda \eta(X)\eta(Y) \quad (2.8)$$

From equation (2.8), we have

$$QX = -(\lambda + 1)X + \eta(X)\xi, \quad (2.9)$$

$$S(X, \xi) = -\lambda \eta(X), \quad (2.10)$$

$$r = -\lambda n - (n - 1), \quad (2.11)$$

where  $S$  is the Ricci tensor,  $Q$  is the Ricci operator and  $r$  is the scalar curvature on  $M$ .

**Theorem 3.1:** A Concircular curvature tensor  $\tilde{C}$  in Kenmostu manifold satisfying  $R(\xi, X) \cdot \tilde{C} = 0$  is expanding.

**Proof:** Let us suppose  $(R(\xi, X) \cdot \tilde{C})(U, V)W = 0$

$$\Rightarrow R(\xi, X) \tilde{C}(U, V)W - \tilde{C}(R(\xi, X)U, V)W - \tilde{C}(U, R(\xi, X)V)W - \tilde{C}(U, V)R(\xi, X)W = 0. \quad (3.1)$$

Using equation (2.2.8) in above equation, we get

$$\begin{aligned} & \eta(\tilde{C}(U, V)W)X - g(X, \tilde{C}(U, V)W)\xi - \tilde{C}(\eta(U)X - g(X, U)\xi, V)W - \tilde{C}(U, \eta(V)X - g(X, V)\xi)W \\ & - \tilde{C}(U, V)\{\eta(W)X - (X, W)\xi\} = 0, \\ & \Rightarrow \eta(\tilde{C}(U, V)W)X - g(X, \tilde{C}(U, V)W)\xi - \eta(U)\tilde{C}(X, V)W + g(X, U)\tilde{C}(\xi, V)W - \eta(V)\tilde{C}(U, X)W \\ & + g(X, V)\tilde{C}(U, \xi)W - \eta(W)\tilde{C}(U, V)X + g(X, W)\tilde{C}(U, V)\xi = 0. \end{aligned} \quad (3.2)$$

Now taking an inner product of above equation with  $\xi$ , we get

$$\begin{aligned} & \Rightarrow \eta(\tilde{C}(U, V)W)\eta(X) - g(X, \tilde{C}(U, V)W) - \eta(U)\eta(\tilde{C}(X, V)W) + g(X, U)\eta(\tilde{C}(\xi, V)W) - \eta(V)\tilde{C}(U, V)W + \\ & g(X, V)\eta(\tilde{C}(U, \xi)W) - \eta(W)\eta(\tilde{C}(U, V)X) + g(X, W)\eta(\tilde{C}(U, V)\xi) = 0. \end{aligned} \quad (3.3)$$

Using equation (1.6) and (1.7) in above equation, we get

$$\begin{aligned} & \left[1 + \frac{r}{n(n-1)}\right][g(U, W)\eta(V) - g(V, W)\eta(U)]\eta(X) - g(X, \tilde{C}(U, V)W) - \eta(U)\left[1 + \frac{r}{n(n-1)}\right][g(X, W) \\ & \eta(V) - g(V, W)\eta(X)] + g(X, U)\left[1 + \frac{r}{n(n-1)}\right][g(\xi, W)\eta(V) - g(V, W)\eta(\xi)] - \eta(V)\left[1 + \frac{r}{n(n-1)}\right] \end{aligned}$$

$$\begin{aligned}
& [g(U, W) \eta(X) - g(X, W)\eta(U)] + g(X, V) \left[1 + \frac{r}{n(n-1)}\right] [g(U, W) \eta(\xi) - g(\xi, W)\eta(U)] - \eta(W) \left[1 + \frac{r}{n(n-1)}\right] [g(U, X) \eta(V) - \\
& g(V, X)\eta(U)] + g(X, W) \left[1 + \frac{r}{n(n-1)}\right] [g(U, \xi) \eta(V) - g(V, \xi)\eta(U)] = 0, \\
& \Rightarrow \left[1 + \frac{r}{n(n-1)}\right] [g(U, W) \eta(X)\eta(V) - g(V, W)\eta(U)\eta(X) - g(X, W) \eta(V)\eta(U) + g(V, W)\eta(X)\eta(U) + g(X, \\
& U) \eta(W)\eta(V) - g(X, U) g(V, W) - g(U, W) \eta(V)\eta(X) + g(X, W) \eta(V)\eta(U) + g(X, V) g(U, W) - g(X, V) \\
& \eta(W)\eta(U) + g(U, X) \eta(W)\eta(V) + g(V, X) \eta(W)\eta(U) + g(X, W) \eta(U)\eta(V) - g(X, W) \eta(U)\eta(V)] - \\
& g(X, \tilde{C}(U, V)W) = 0, \\
& \Rightarrow \left[1 + \frac{r}{n(n-1)}\right] [g(X, V) g(U, W) - g(X, U) g(V, W)] - g(X, \tilde{C}(U, V)W) = 0.
\end{aligned}$$

The above equation can be written as

$$-g(X, \tilde{C}(U, V)W) + \left[1 + \frac{r}{n(n-1)}\right] [g(X, V) g(U, W) - g(X, U) g(V, W)] = 0. \quad (3.4)$$

In view of equation (1.1), above equation takes the form

$$\begin{aligned}
& -g(X, R(U, V)W) - \frac{r}{n(n-1)} [g(V, W)U - g(U, W)V] + \left[1 + \frac{r}{n(n-1)}\right] [g(X, V) g(U, W) \\
& - g(X, U) g(V, W)] = 0 \\
& \Rightarrow -g(X, R(U, V)W) - \frac{r}{n(n-1)} [g(X, V) g(U, W) - g(X, U) g(V, W)] + \left[1 + \frac{r}{n(n-1)}\right] [g(X, V) g(U, W) \\
& - g(X, U) g(V, W)] = 0 \\
& \Rightarrow -g(X, R(U, V)W) + [g(X, V) g(U, W) - g(X, U) g(V, W)] = 0
\end{aligned} \quad (3.5)$$

Taking  $X = U = e_i$  in above equation and summing over  $i$ ,  $1 \leq i \leq n$ , we get

$$\begin{aligned}
& -R(e_i, V, W, e_i) + [g(e_i, v) g(e_i, W) - g(e_i, e_i) g(V, W)] = 0 \\
& \Rightarrow -S(V, W) + [g(V, W) - ng(V, W)] = 0.
\end{aligned} \quad (3.6)$$

Taking  $V = W = \xi$  in above equation, we get

$$\begin{aligned}
& -S(\xi, \xi) + [g(\xi, \xi) - ng(\xi, \xi)] = 0 \\
& \Rightarrow \lambda + (1 - n) = 0 \\
& \Rightarrow \lambda = n - 1.
\end{aligned}$$

This shows that  $\lambda$  is positive, it means that the concircular curvative tensor  $\tilde{C}$  in Kenmostu manifold satisfying  $R(\xi, X).\tilde{C} = 0$  is expanding. This completes the proof.

## References

1. Blair DE. On the geometric meaning of the Bochner tensor. *Geometriae Dedicata*. 1975; 4:1.
2. Kenmotsu K. A class of almost contact Riemannian manifolds. *The Tohoku Mathematical Journal*. 1972, 24.
3. Blair DE. Contact Manifolds in Riemannian Geometry, vol. 509 of Lecture Notes in Mathematics, Springer, Berlin, Germany, 1976.
4. Sharma R. On the curvature of contact metric manifolds, *J Geom*. 1995, 53.
5. Bernardini A. Riemannian manifolds with concircular curvature tensor, *Atti, sem. Mat. Fis. Univ. Modena*, 1981, 30.
6. Sharma SR. On almost contact manifold, *Indian J. Pure and Applied Math*, 1974, 5.
7. Perelman G. The entropy formula for the Ricci flow and its geometric applications, *Math*, 1972, 29.
8. Ively T. Ricci solitons on compact 3- manifolds, *Different. Geom. Appl.* 1993, 3.
9. Hamilton RS. Almost contact structures and curvature tensors, *Kodai Math. J.* 1992; 4:1-27.
10. Sasaki S. On differentiable manifolds with certain structures which are closely related to almost contact structure, *Tohoku Math. J.* 1960; 2:459-476.
11. Tanno S. Ricci curvatures of contact Riemannian manifolds, *Tohoku Math. J.* 1988; 40:441-448.