



International Journal of Research in Agronomy

E-ISSN: 2618-0618

P-ISSN: 2618-060X

© Agronomy

www.agronomyjournals.com

2024; 7(8): 572-574

Received: 25-05-2024

Accepted: 29-06-2024

PM Chaudhary

Senior Research Fellow,
Agricultural Research Station,
Anand Agricultural University,
Jabugam, Gujarat, India

HN Luea

Assistant Professor & Head,
Department of Fruit Science,
College of Horticulture,
Sardarkrushinagar Dantiwada
Agricultural University, Jagudan,
Gujarat, India

PD Patel

Assistant Professor & Head,
Department of Horticulture,
College of Agriculture, Anand
Agricultural University, Jabugam,
Gujarat, India

SN Makwana

Research Associate, Agricultural
Research Station, Anand
Agricultural University, Jabugam,
Gujarat, India

VM Gohil

Ph.D. Scholar, Department of
Fruit Science, ASPEE College of
Horticulture, Navsari Agricultural
University, Navsari, Gujarat, India

Corresponding Author:

PM Chaudhary

Senior Research Fellow,
Agricultural Research Station,
Anand Agricultural University,
Jabugam, Gujarat, India

Impact of different seed treatments on germination of chironji (*Buchanania lanzan* Spreng.)

PM Chaudhary, HN Luea, PD Patel, SN Makwana and VM Gohil

DOI: <https://doi.org/10.33545/2618060X.2024.v7.i8h.1319>

Abstract

An experiment was conducted to study the effect of different seed treatments on germination and seedling growth of chironji (*Buchanania lanzan* Spreng.) carried out under propagation chamber at College Farm, College of Horticulture, Jagudan during October, 2022 to March, 2023 and laid out by Completely Randomized Design with three repetition and fifteen treatments viz., T₁: Absolute control; T₂: Water soaking for 24 hrs.; T₃: Hot water treatment (99±1 °C for 10 second); T₄: Hot water treatment (99±1 °C for 30 second); T₅: Mechanical scarification (break seed coat by hammering); T₆: H₂SO₄ @ 2.5% for 10 minute; T₇: H₂SO₄ @ 5% for 10 minute; T₈: GA₃ @ 200 ppm for 24 hrs.; T₉: GA₃ @ 400 ppm for 24 hrs.; T₁₀: Cow urine @ 10% for 24 hrs.; T₁₁: Cow urine @ 20% for 24 hrs.; T₁₂: Cow dung slurry @ 10% for 24 hrs.; T₁₃: Cow dung slurry @ 20% for 24 hrs.; T₁₄: Beejamrut @ 5% for 24 hrs.; T₁₅: Beejamrut @ 10% for 24 hours. Among seed soaking treatments, T₅ [Mechanical scarification (break seed coat by hammering)] significantly recorded minimum number of days taken for germination (24.53) and maximum germination percentage (63.67).

Keywords: Germination, GA₃, H₂SO₄, cow urine, cow dung slurry, Beejamrut and seed

Introduction

Chironji (*Buchanania lanzan* Spreng.) a member of the family Anacardiaceae, originated in the Indian sub-continent. The tree is found growing as wild in the tropical deciduous forests of north, west and central India mostly in the states of Madhya Pradesh, Bihar, Orissa, Andhra Pradesh, Chhattisgarh, Jharkhand, Gujarat, Rajasthan and Maharashtra. In Gujarat, it is found growing especially in Dang, Valsad, Vadodara and Dahod district. Chironji is a small to medium size semi evergreen minor fruit tree with straight cylindrical trunk up to 15 meter height. Trees have the alternate bearing nature like mango. Tree shows deciduous nature for short time in summer and new leaves come in the late May. Leaves are 6-10 inches in length, oblong, obtuse; flowers whitish green, sessile; fruit drupe, green when immature and dark black at ripened stage. Flowering occurs in the month of January-February and fruits ripen in April-May.

The flesh of ripe fruit is very palatable and is largely eaten raw or roasted. The oily kernels are the most important part which are used in preparation of puddings. Kernel contains 34-47 percent oil. Chironji oil is non-repellent and non-toxic which is suitable for human consumption (Banerjee and Jain, 1988) ^[1]. Hemawathy and Prabhakar (1988) ^[3] reported that the total kernel lipid extracted from chironji was 65.60 percent of dry kernel and consisted of 90.40 percent neutral lipids, 3.40 percent glycolipids and 6.20 percent phospholipids. The kernels are reported to be used by tribal of Gujarat as brain tonic. An ointment made out of the kernels is used to cure itches of the skin and to remove blemishes from the face. The bark furnishes a natural varnish and is used in tanning in Kerala. The leaves are reported to be valued for their tonic and cardiotonic properties and their powder is a common medicine for wounds.

In this regard, different seed treatments are being used for accelerating seed germination and seedling growth of chironji. GA₃ is used for weakening of the seed coat so that the radical of the seedling can come out through the seed coat. GA₃ induces the synthesis of amylase and other hydrolytic enzymes during the early stages of seed germination. GA₃ controls mobilization of starch which acts as a respiratory substrate leading to immediate enhancement in cell elongation. Gibberellins also help in enhancing the availability of reserved mineral elements which promote

the germination process.

Cow dung slurry voided by solid excreta of animals represents the digested and non-digested portion of the animal feed. It contains 70-80% water, admixed with digestive juices, tissues of alimentary canal, microbial cells. Cow dung slurry also contains some growth promoting substances like auxin, N, P, K, micronutrients and biodigestible enzymes which have been cause for softening of seed coat and thereby enhances seed germination and growth of seedlings (Raj *et al.*, 2014) [9].

Cow urine is also known as “*Gomutra*”. Cow urine contains about 1.21% N, 0.01% P₂O₅ and 1.35% K₂O (Subramanian, 2005) [13] and also contains physiologically active substances viz., growth regulators, nutrients (Kamalam and Rajappan, 1989) [4] and trace elements (Munoz, 1988). Cow urine contains iron, urea, uric acid, estrogens and progesterone which affects the inhibitory responses to seed germination, shoot growth and seedling vigour.

Beejamrut is a technique of seed treatment from locally available ingredients. It protects the crop from harmful soil borne and seed borne pathogens during the initial stages of germination and establishment.

Materials and Methods

The present experiment was carried out during 2022-23 at College Farm, College of Horticulture, Sardarkrushinagar Dantiwada Agricultural University, Jagudan in propagation chamber. A completely randomized design was adopted with three replications. The experiment included fifteen treatments viz., T₁: Absolute control; T₂: Water soaking for 24 hrs.; T₃: Hot water treatment (99±1 °C for 10 second); T₄: Hot water treatment (99±1 °C for 30 second); T₅: Mechanical scarification (break seed coat by hammering); T₆: H₂SO₄ @ 2.5% for 10 minute; T₇: H₂SO₄ @ 5% for 10 minute; T₈: GA₃ @ 200 ppm for 24 hrs.; T₉: GA₃ @ 400 ppm for 24 hrs.; T₁₀: Cow urine @ 10% for 24 hrs.; T₁₁: Cow urine @ 20% for 24 hrs.; T₁₂: Cow dung slurry @ 10% for 24 hrs.; T₁₃: Cow dung slurry @ 20% for 24 hrs.; T₁₄: *Beejamrut* @ 5% for 24 hrs.; T₁₅: *Beejamrut* @ 10% for 24 hours. Sowing were done just after the compilation

of seed treatments. Sowing of seed in 20 cm × 15 cm size polythene bags, for improve the drainage, holes were made in polythene bags and filled with potting mixture which was prepared by mixing one part of soil, one part of sand and one part of well- rotted FYM at 1 cm depth and were kept in propagation chamber. The polythene bags were watered daily till final data were recorded. Observations were recorded daily on germination parameters. The date of first plumule emergence was recorded and computing the difference between date of sowing and plumule emergence was recorded as a number of days taken for germination. The germination percentage was calculated by using below formula:

$$\text{Germination percentage} = \frac{\text{Number of seeds germinated}}{\text{Total number of seeds sown}} \times 100$$

The data collected from the five labeled seedlings in each treatment in repetition were average and subjected statistical analysis (Panse and Sukhatme, 1985) [8].

Results and Discussion

Number of days taken for germination

The statistical analysed data presented in table 1, revealed that number of days taken for germination of chironji were significantly influenced by different pre sowing treatments. The seed subjected to significantly minimum number of days taken for germination (24.53 days) were recorded in treatment T₅ [Mechanical scarification (break seed coat by hammering)] which was statistically at par with T₈ (25.47 days) and T₉ (26.80 days). The maximum number of days taken for germination (39.53 days) was noted in absolute control (T₁).

It might be due to the fact that practice of mechanical scarification injured the seed coat thereby, it increased the permeability of air and water through seed, which ended up with the elongation of the embryonic axis and favours the early germination (Mahadev *et al.*, 2021) [6]. These results are conformation with Shukla and Solanki (2000) [12] and Sharma (2016) [11] in chironji.

Table 1: Effect of seed soaking treatment on number of days taken for germination of chironji

Treatment	Number of days taken for germination
T ₁ : Absolute control	39.53
T ₂ : Water soaking for 24 hrs.	34.00
T ₃ : Hot water treatment (99±1 °C for 10 second)	29.20
T ₄ : Hot water treatment (99±1 °C for 30 second)	29.47
T ₅ : Mechanical scarification (break seed coat by hammering)	24.53
T ₆ : H ₂ SO ₄ @ 2.5% for 10 minute	29.87
T ₇ : H ₂ SO ₄ @ 5% for 10 minute	28.73
T ₈ : GA ₃ @ 200 ppm for 24 hrs.	25.47
T ₉ : GA ₃ @ 400 ppm for 24 hrs.	26.80
T ₁₀ : Cow urine @ 10% for 24 hrs.	32.80
T ₁₁ : Cow urine @ 20% for 24 hrs.	33.67
T ₁₂ : Cow dung slurry @ 10% for 24 hrs.	31.98
T ₁₃ : Cow dung slurry @ 20% for 24 hrs.	33.93
T ₁₄ : <i>Beejamrut</i> @ 5% for 24 hrs.	31.00
T ₁₅ : <i>Beejamrut</i> @ 10% for 24 hrs.	32.07
S.Em. ±	1.01
C.D. at 5%	2.88
C.V.%	5.69

Germination percentage

A perusal of data presented in the table 2 reveals that the maximum germination percentage maximum germination percentage (63.67) were recorded in treatment T₅ [Mechanical

scarification (break seed coat by hammering)] which was statistically at par with T₈ (62.50%) and T₉ (60.00%). The minimum germination percentage (36.67) was observed in absolute control (T₁).It might be due to higher germination

percentage is as result of mechanical scarification which facilitated entry of water into the seed kernels. Similar results

were reported by Hassen *et al.* (2005)^[2] in ber; Saied *et al.* (2008)^[10] in ber and Khobragade and Rai (2022)^[5] in chironji.

Table 2: Effect of seed soaking treatments on germination percentage of chironji

Treatment	Germination percentage
T ₁ : Absolute control	36.67
T ₂ : Water soaking for 24 hrs.	40.00
T ₃ : Hot water treatment (99±1 °C for 10 second)	43.33
T ₄ : Hot water treatment (99±1 °C for 30 second)	38.33
T ₅ : Mechanical scarification (break seed coat by hammering)	63.67
T ₆ : H ₂ SO ₄ @ 2.5% for 10 minute	41.67
T ₇ : H ₂ SO ₄ @ 5% for 10 minute	45.00
T ₈ : GA ₃ @ 200 ppm for 24 hrs.	62.50
T ₉ : GA ₃ @ 400 ppm for 24 hrs.	60.00
T ₁₀ : Cow urine @ 10% for 24 hrs.	46.67
T ₁₁ : Cow urine @ 20% for 24 hrs.	48.33
T ₁₂ : Cow dung slurry @ 10% for 24 hrs.	50.00
T ₁₃ : Cow dung slurry @ 20% for 24 hrs.	43.33
T ₁₄ : Beejamrut @ 5% for 24 hrs.	46.67
T ₁₅ : Beejamrut @ 10% for 24 hrs.	48.33
S.E.m. ±	1.39
C.D. at 5%	3.95
C.V.%	5.04

Conclusion

The present investigation can be concluded that among the different seed soaking treatments, mechanical scarification (break seed coat by hammering) significantly exhibited maximum germination percentage along with minimum number of days taken for germination.

References

- Banerjee A, Jain M. Investigations on *Buchanania lanzan* seed oil. *Fitoterapia*. 1988;59:406.
- Hassen A, Rethman NFG, Niekerk VWA. Effect of different seed treatment options on dormancy breaking, germination and emergence of *Ziziphus mucronata* seed. *Trop Grassl*. 2005;39:124-128.
- Hemawathy J, Prabhakar JV. Lipid composition of chironji (*Buchanania lanzan* Spreng.) kernel. *J Food Compos Anal*. 1988;1:366-370.
- Kamalam J, Rajappan N. Effect of seed hardening on germination and seedling vigour in paddy. *Seed Res*. 1989;17(2):188-190.
- Khobragade ND, Rai N. Effect of pretreatments on seed germination and seedling growth of *Buchanania lanzan*. *Pharma Innov J*. 2022;11(1):104-106.
- Mahadev LP, Ahire DB, Patel HD. Influence of different chemicals on vegetative growth of chironji (*Buchanania lanzan* Spreng.). *J Agric Res Life Sci*. 2021;5(2):27-30.
- Munoz AM. Increasing the vigour of rice seeds with trace element application. *Arrazy*. 1988;37:20-27.
- Panse VG, Sukhatme PV. Statistical methods for agricultural workers. 4th ed. New Delhi: ICAR; c1985. p. 87-89.
- Raj A, Jhariya M, Toppo P. Cow dung for eco-friendly and sustainable productive farming. *Int. J Sci. Res*. 2014;3(8):201-202.
- Saied A, Gebauer J, Buerkert A. Effects of different scarification methods on germination of *Ziziphus spinachristi* seeds. *Seed Sci. Technol*. 2008;36:201-205.
- Sharma DK. Effect of plant growth regulators and scarification on germination and seedling growth of chironji (*Buchanania lanzan* Spreng.). *Adv. Life Sci*. 2016;5(8):3237-3241.
- Shukla SK, Solanki KR. Studies on seed germination, plant survival and growth of chironji (*Buchanania lanzan* Spreng.). *J Trop For*. 2000;16(1):44-49.
- Subramanian A. Effect of panchagavya on *Escherichia coli* in procured milk. *Indian Vet J*. 2005;82:799-800.