





International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
IJABR 2024; 8(8): 948-950
www.biochemjournal.com
Received: 05-07-2024
Accepted: 11-08-2024

Dr. Tasmiya Imtiyaz
Department of Agricultural
Microbiology, University of
Agricultural Science,
Bangalore, Karnataka, India

Acid stress and microbial resilience: Insights into adaptive responses

Dr. Tasmiya Imtiyaz

DOI: <https://doi.org/10.33545/26174693.2024.v8.i8l.1908>

Abstract

Microbes encounter fluctuations in environmental factors, including pH, which directly affect microbial growth. Acid stress, defined as the combined biological effect of H⁺ ions and weak acid concentrations, triggers various microbial responses. This article examines the adaptive mechanisms employed by bacteria to survive and thrive in acidic conditions. Acidophilic, alkalophilic, and neutralophilic bacteria each demonstrate unique strategies for maintaining pH homeostasis. By producing enzymes, modulating proton pumps, and employing acid tolerance responses (ATRs), these microbes can survive and adapt to harsh acidic environments. Understanding these mechanisms provides insights into microbial survival and has implications for industrial applications and pathogenic control.

Keywords: Microbes, environmental fluctuations, pH, microbial growth

Introduction

Microorganisms exist in diverse environments where they are frequently subjected to fluctuating conditions that can significantly impact their growth and survival. Among these environmental factors, pH variation plays a critical role. Microbes exhibit a remarkable ability to grow over a wide range of hydrogen ion concentrations, from acidic to alkaline conditions. For example, acidophilic bacteria thrive in highly acidic environments such as sulfur springs with pH levels as low as 1, while alkalophilic bacteria prefer highly alkaline conditions, such as those found in soda lakes with pH levels up to 11.

Most bacteria that inhabit human-associated environments, including common species like *Escherichia coli*, *Salmonella enterica*, *Streptococcus mutans*, and *Bacillus subtilis*, prefer neutral pH values and are classified as neutralophilic. These bacteria generally grow within a pH range of 5 to 9. Although neutralophiles do not typically grow under extreme acidic or basic conditions, they can survive such exposures through adaptive transitions where pH gradually changes, allowing them to adjust their internal mechanisms accordingly.

The shift in environmental pH towards acidity imposes acid stress on microbes. Acid stress encompasses the combined biological effects of increased H⁺ ions and weak acid concentrations. To cope with acidifying conditions, microbes employ various strategies to maintain pH homeostasis and ensure survival. These strategies include producing enzymes that convert acidic metabolites to neutral or alkaline products, such as glutamate decarboxylase, lysine decarboxylase, and arginine decarboxylase, which are upregulated in response to external acidic pH (Peter *et al.*, 2019) ^[2].

Homeostasis Mechanisms in Bacteria

Both Gram-negative and Gram-positive bacteria have developed sophisticated homeostasis mechanisms to regulate internal pH during growth. These mechanisms involve the modulation of primary proton pumps and the activity of K⁺/H⁺, Na⁺/H⁺ antiporters, and Mg²⁺-dependent proton-translocating ATPase. These systems are critical for maintaining internal pH within a viable range, especially when external pH conditions fluctuate.

1. Proton Pumps: These include primary proton pumps such as the F₁F₀-ATPase and other proton translocating ATPases that expel excess protons from the cytoplasm to maintain a stable internal pH.

Corresponding Author:
Dr. Tasmiya Imtiyaz
Department of Agricultural
Microbiology, University of
Agricultural Science,
Bangalore, Karnataka, India

- 2. Antiporters:** K^+/H^+ and Na^+/H^+ antiporters play a significant role in exchanging intracellular H^+ ions with extracellular K^+ or Na^+ ions. This exchange helps to alleviate the acid load inside the cell.
- 3. Buffers and Compatible Solutes:** Bacteria also produce intracellular buffers and compatible solutes, such as glutamate and proline, which help to neutralize the internal pH and protect cellular functions.

Inducible Systems and Acid Tolerance Response (ATR)

When pH conditions fall outside the optimal growth range, bacteria activate inducible systems that provide additional protection. These adaptive mechanisms include the acid tolerance response (ATR), which involves genetic and physiological changes that enhance the cell's ability to withstand acidic conditions. The complete activation of ATR results in the increased expression of proteins known as acid shock proteins, some of which are also induced by heat shock (Krulwich *et al.*, 2011) [6].

Acid Shock Proteins: These proteins are involved in protecting cellular components from damage caused by acid stress. They include chaperones that refold damaged proteins, proteases that degrade irreparably damaged proteins, and other proteins that help to stabilize DNA and RNA under acidic conditions.

Pathogenic Bacteria and Acid Stress

Pathogenic bacteria such as *Listeria monocytogenes*, *E. coli*, and *Bacillus cereus* frequently encounter acidic environments, particularly in the human gastrointestinal tract. These bacteria have evolved various mechanisms to survive acid stress, including the glutamate decarboxylase system, arginine decarboxylase system, and DNA repair systems (Skandamis *et al.*, 2008) [5]. For instance, *Streptococcus mutans* forms dense biofilms to protect itself against extracellular acid shock (Robert and Dennis, 2010) [3]. Additionally, urease-producing pathogens like *Helicobacter pylori* can survive in extremely acidic conditions by generating cytoplasmic ammonia through urea hydrolysis, thereby neutralizing the surrounding acidity.

Specific Adaptations in Pathogens

- 1. Glutamate Decarboxylase System:** Converts glutamate to gamma-aminobutyric acid (GABA), consuming protons in the process and thereby increasing the internal pH.
- 2. Arginine Decarboxylase System:** Catalyzes the conversion of arginine to agmatine, which also consumes protons and helps to mitigate acid stress.
- 3. DNA Repair Systems:** Enhanced DNA repair

mechanisms ensure the integrity of genetic material under acidic conditions, preventing mutations and maintaining cellular functions.

- 4. Biofilm Formation:** In *Streptococcus mutans*, biofilm formation provides a physical barrier that protects bacterial communities from acidic environments, allowing them to persist and cause dental caries.

Industrially Relevant Bacteria and Acid Stress

Certain bacteria have harnessed their acid tolerance mechanisms for industrial applications. Acetic acid bacteria, obligate aerobes used in vinegar production, exhibit remarkable tolerance to acetic acid. This ability is attributed to their efficient ethanol oxidation pathway, involving alcohol dehydrogenase, aldehyde dehydrogenase, and terminal oxidase enzymes on the cell membrane (Shigeru *et al.*, 2006) [4]. Lactic acid bacteria, widely used in the production of fermented foods and probiotics, can also induce acid tolerance in response to mild acid conditions, enhancing their survival and functional properties (Chongde *et al.*, 2014) [1].

Applications in Industry

- 1. Vinegar Production:** Acetic acid bacteria are used for their ability to oxidize ethanol to acetic acid, even in high-acid environments. This trait is essential for producing high-quality vinegar.
- 2. Fermented Foods:** Lactic acid bacteria are critical in the production of yogurt, cheese, sauerkraut, and other fermented foods. Their ability to tolerate and produce acidic conditions is central to the fermentation process.
- 3. Probiotics:** The acid tolerance of lactic acid bacteria makes them suitable for surviving the acidic conditions of the stomach, allowing them to confer health benefits to the host.

Ecological and Evolutionary Implications

The conservation of acid tolerance mechanisms across diverse bacterial species suggests that these strategies provide significant adaptive advantages. By modulating the activity of RNAP and thus the transcriptional profile of the cell, 6S RNA allows bacteria to quickly adjust to fluctuating environmental conditions. This ability to rapidly switch between different transcriptional states is likely crucial for survival in habitats where nutrient availability and other conditions can change rapidly.

Furthermore, the ability of 6S RNA to act as a template for pRNA synthesis represents an elegant example of molecular mimicry and regulatory versatility. This feature may provide insights into the evolution of regulatory RNAs and their integration into complex cellular networks.




Fig 1: Acid stress responses in microbial cells

Conclusion

Bacteria employ a variety of acid stress mechanisms to survive and thrive in acidic environments. These mechanisms, including enzyme production, proton pump modulation, and inducible acid tolerance responses, work synergistically to ensure microbial growth and metabolism under hostile conditions. Understanding these adaptive strategies not only sheds light on microbial survival but also has significant implications for industrial applications and the control of pathogenic bacteria. Further research into these mechanisms will undoubtedly reveal new insights into the resilience and versatility of microbial life.

References

1. Chongde W, Jun H, Rongqing Z. Progress in engineering acid stress resistance of lactic acid bacteria. *Appl Microbiol Biotechnol*. 2014;98:1055-1063.
2. Peter L, Angela T, Daniela D. Coping with low pH: molecular strategies in neutrophilic bacteria. *FEMS Microbiol Rev*. 2019;8:1-35.
3. Robert M, Dennis C. Acid tolerance mechanisms utilized by *Streptococcus mutans*. *Front Microbiol*. 2010;5:403-417.
4. Shigeru N, Masahiro F, Sueharu H. Putative ABC transporter responsible for acetic acid resistance in *Acetobacter aceti*. *Appl. Environ Microbiol*. 2006;72:497-505.
5. Skandamis PN, Yoon Y, Stopforth JD, Kendall PA, Sofos JN. Acid tolerance of *Listeria monocytogenes* after exposure to single and multiple sublethal stresses. *Food Microbiol*. 2008;25:294-303.
6. Krulwich A, George S, Etana P. Molecular aspects of bacterial pH sensing and homeostasis. *FEMS Microbiol Rev*. 2011;9:330-343.