



International Journal of Agriculture and Food Science

ISSN Print: 2664-844X
ISSN Online: 2664-8458
IJAFS 2024; 6(2): 87-90
www.agriculturaljournals.com
Received: 06-08-2024
Accepted: 11-09-2024

José Antonio Costa
Senior Lecturer, Department
of Agronomy, Ceará
Agricultural College, Fortaleza,
Brazil

Assessing different weed control methods in soybean fields

José Antonio Costa

DOI: <https://doi.org/10.33545/2664844X.2024.v6.i2b.208>

Abstract

Weed management is a critical aspect of soybean production, as weeds compete for resources, reduce yield potential, and increase production costs. This paper evaluates and compares various weed control methods cultural, mechanical, chemical, and biological based on their effectiveness, environmental impact, economic feasibility, and relevance in sustainable agriculture. While each method has distinct advantages, limitations such as herbicide resistance, soil degradation, and labor requirements underscore the importance of an integrated approach. Integrated Weed Management (IWM) emerges as the most effective strategy, combining diverse methods to enhance weed suppression, improve crop yields, and promote environmental sustainability. By reviewing relevant studies and field practices, this paper highlights the need for adaptive and innovative weed control systems to address modern agricultural challenges.

Keywords: Weed management, Soybean production, Integrated Weed Management (IWM), Sustainable agriculture, Herbicide resistance

Introduction


Soybean (*Glycine max*) is one of the most widely cultivated and economically significant crops globally, valued for its high protein content and versatile applications in food, feed, and industrial products. However, achieving optimal yields in soybean production is often hampered by weed infestations, which compete for essential resources such as sunlight, water, and nutrients. Weeds not only reduce crop productivity but also serve as hosts for pests and diseases, further compounding the challenges faced by farmers. Effective weed management is therefore crucial for maintaining crop health and maximizing economic returns. The selection of appropriate weed control methods is influenced by several factors, including the type of weeds, environmental conditions, farming system, and economic constraints. Traditionally, weed management has relied heavily on chemical herbicides due to their rapid and reliable action. However, the excessive and unregulated use of herbicides has led to significant problems, including the development of herbicide-resistant weed species, soil and water contamination, and risks to non-target organisms. These issues underscore the need for sustainable and integrated approaches to weed management. Mechanical methods, such as tillage and inter-row cultivation, provide immediate weed suppression but can result in soil erosion and compaction when overused. Biological control, which utilizes natural enemies or allelopathic plants, offers environmentally friendly alternatives but often requires longer periods to demonstrate effectiveness. Cultural methods, including crop rotation and cover cropping, enhance soil health and disrupt weed life cycles but may not be sufficient to manage high weed densities. Each of these methods has unique strengths and limitations, necessitating a comprehensive evaluation to identify the most effective strategies for soybean fields. Integrated Weed Management (IWM) has emerged as a promising approach to address these challenges. By combining multiple weed control methods, IWM leverages the strengths of each while mitigating their individual drawbacks. This paper aims to evaluate the effectiveness, environmental impact, and economic feasibility of various weed control strategies, with a particular focus on their integration into IWM systems. Drawing on relevant studies and practical field applications, this research provides insights into sustainable weed management practices for soybean production, highlighting the importance of innovation and adaptability in modern agriculture.

Corresponding Author:
José Antonio Costa
Senior Lecturer, Department
of Agronomy, Ceará
Agricultural College, Fortaleza,
Brazil

Objective

The objective of this paper is to evaluate and compare various weed control methods in soybean fields, focusing on their effectiveness, environmental impact, and economic feasibility, while highlighting the importance of integrated approaches for sustainable agriculture.

Classification of weed control methods



Cultural weed control methods

Cultural weed control methods are fundamental to sustainable agricultural practices, relying on agronomic techniques to minimize weed competition and promote soybean growth. These methods, including crop rotation, cover cropping, mulching, and optimized planting practices, offer an environmentally friendly alternative to chemical herbicides. Crop rotation disrupts weed life cycles by alternating soybean with crops like maize or wheat, reducing the prevalence of weed species that thrive in continuous soybean systems. Similarly, cover crops, such as rye or clover, suppress weeds by outcompeting them for sunlight, nutrients, and water, while improving soil structure and fertility. Mulching provides an additional layer of protection by physically blocking sunlight, preventing weed germination, and conserving soil moisture. Optimized planting practices, such as adjusting row spacing and planting density, increase crop competitiveness against weeds, reducing their growth and establishment. These methods are particularly advantageous in organic farming systems, where chemical inputs are restricted, and in regions facing herbicide resistance, providing a sustainable solution to persistent weed challenges. Compared to mechanical or chemical methods, cultural weed control is less immediate but offers long-term benefits. Unlike herbicides, which may lead to resistance and environmental contamination, cultural methods enhance soil health, promote biodiversity, and reduce reliance on external inputs. Studies such as those by Liebman and Dyck (1993) have demonstrated the efficacy of crop rotation in reducing weed biomass and seedbank densities. Similarly, research by Teasdale *et al.*, (2007) highlights that cover crops can reduce weed density by up to 80%, aligning with findings that mulching can significantly suppress weed growth and conserve soil moisture. Economic feasibility is another critical factor in comparing cultural weed control with other methods. While the initial investment for practices like cover cropping or mulching may be higher, the reduced dependence on herbicides and improved yield potential often offset these costs over time. This contrasts with chemical methods, which, despite their immediate effectiveness, can lead to increased costs due to resistance management and the need for multiple applications. In summary, cultural weed control methods provide a sustainable and economically viable approach to managing weeds in soybean fields. They complement mechanical and chemical strategies, addressing the limitations of each. Previous studies consistently support their effectiveness and highlight their role in promoting environmental and economic sustainability. By integrating

cultural practices into a broader weed management plan, farmers can achieve long-term productivity and ecological balance

Mechanical weed control methods

Mechanical weed control methods involve the physical removal or destruction of weeds using tools or machinery. These methods have been used for centuries and remain integral to modern weed management, particularly in systems aiming to reduce chemical inputs. Common techniques include tillage, inter-row cultivation, and hand weeding, each varying in effectiveness, feasibility, and environmental impact.

Tillage is one of the most widely used mechanical methods, breaking the soil to uproot weeds and disrupt their growth cycle. This method is particularly effective against annual weeds, as it buries weed seeds deeper into the soil, reducing their chances of germination. However, over-reliance on tillage can lead to soil erosion, compaction, and disruption of beneficial soil organisms, which has been noted in studies like those by Triplett and Dick (2008), highlighting the environmental trade-offs of this approach.

Inter-row cultivation involves using specialized machinery to remove weeds between crop rows without disturbing the soybean plants. This method is precise and efficient, offering excellent weed control in early to mid-growth stages of the crop. It minimizes competition for resources, improving soybean yield. However, its success depends on accurate row spacing and machinery calibration to prevent crop damage. Studies such as Knezevic *et al.*, (2002) emphasize the importance of timing and precision for optimal results with inter-row cultivation.

Hand weeding, while labor-intensive, is highly effective in small-scale or organic farming systems. This method allows for the selective removal of weeds without harming the crop. It is especially useful for managing weeds in areas where machinery cannot be used or when dealing with small weed infestations. Despite its effectiveness, the high labor cost and time requirements limit its feasibility for large-scale soybean production. Compared to cultural and chemical methods, mechanical weed control provides immediate results, making it an appealing choice for addressing severe weed infestations. However, the environmental impact and economic feasibility vary significantly among techniques. For example, while hand weeding is environmentally neutral, tillage can degrade soil structure over time. In contrast, inter-row cultivation offers a middle ground, balancing effectiveness with lower environmental costs when used judiciously. Studies show that integrating mechanical methods with other strategies, such as cultural or chemical weed control, enhances overall effectiveness and sustainability. For example, combining tillage with cover cropping can reduce weed pressure while mitigating soil erosion. This synergy highlights the role of mechanical weed control as a critical component of Integrated Weed Management (IWM).

In conclusion, mechanical weed control methods are effective and versatile tools for managing weeds in soybean fields. While they offer immediate weed suppression, their environmental impact and economic feasibility must be carefully evaluated. Incorporating these methods into a comprehensive weed management plan can maximize their benefits while minimizing their limitations, as supported by various studies and practical field applications.

Chemical weed control methods

Chemical weed control methods involve the use of herbicides to manage weed populations effectively. These methods are widely adopted due to their ability to provide quick and reliable results, particularly in large-scale agricultural systems like soybean production. Herbicides can be classified into pre-emergence and post-emergence types, each with specific applications, advantages, and limitations. Pre-emergence herbicides are applied to the soil before weed and crop emergence, forming a protective barrier that prevents weed germination. These herbicides, such as metolachlor and pendimethalin, are particularly effective in controlling annual grasses and broadleaf weeds during their early growth stages. Studies, including those by Culpepper *et al.*, (2004), show that pre-emergence herbicides can reduce early weed competition significantly, leading to higher crop yields. However, their effectiveness relies on proper timing and soil conditions, such as adequate moisture to activate the herbicide. Post-emergence herbicides are applied after weeds and crops have emerged, targeting weeds actively competing with the soybean crop. These include selective herbicides like glyphosate or dicamba, which are designed to kill specific weed species without harming the crop, especially in herbicide-tolerant soybean varieties. Post-emergence herbicides offer flexibility, as they can be applied when weed problems become visible. However, over-reliance on these herbicides has led to the widespread issue of herbicide-resistant weeds, as documented in studies like those by Powles and Yu (2010). The environmental impact of chemical weed control is a significant concern. While herbicides are efficient in managing weeds, their misuse or overuse can result in soil and water contamination, harm to non-target organisms, and biodiversity loss. Herbicide resistance further complicates this issue, requiring higher doses or alternative chemicals, increasing the ecological footprint. Integrated strategies that rotate herbicides with different modes of action, as recommended by the Weed Science Society of America, can mitigate resistance development and reduce environmental risks.

Economically, chemical weed control is initially cost-effective due to reduced labor and machinery costs. However, the emergence of resistant weed populations can lead to increased expenses for alternative herbicides or supplementary control measures. Studies, such as those by Shaner (2014), highlight the need for sustainable chemical usage to balance immediate cost savings with long-term economic feasibility. Compared to cultural and mechanical methods, chemical weed control provides unparalleled speed and effectiveness but comes with ecological and resistance-related challenges. Incorporating herbicides into an Integrated Weed Management (IWM) system, combining them with cultural practices like crop rotation or mechanical methods such as inter-row cultivation, can enhance their benefits while addressing their limitations. In conclusion, chemical weed control methods remain an essential component of soybean weed management, offering rapid and effective results. However, their long-term sustainability depends on responsible use, resistance management, and integration with other weed control strategies. Studies underscore the importance of adopting a balanced approach to maximize productivity while minimizing environmental and economic risks.

Biological weed control methods

Biological weed control methods involve the use of living organisms, natural processes, or their derivatives to suppress weed growth. These methods are rooted in ecological principles and are increasingly recognized for their potential to provide sustainable weed management while minimizing environmental harm. Biological methods can include the introduction of natural enemies, the use of allelopathic plants, or fostering microbial activity in the soil. Natural enemies, such as insects, pathogens, or grazing animals, are used to target specific weed species. For example, insects like the leafy spurge flea beetle have been deployed to control invasive weeds in certain regions, demonstrating effectiveness in reducing weed biomass. Similarly, fungal pathogens, such as rust fungi, have been introduced to suppress weeds like Canada thistle. Studies, such as those by Charudattan (2001), highlight the success of biocontrol agents in managing weed populations without harming crops or non-target plants. However, the effectiveness of biocontrol agents can be limited by environmental conditions, requiring careful selection and monitoring. Allelopathy is another key approach, where certain plants release biochemicals that inhibit the germination and growth of surrounding weeds. Crops like sorghum and mustard have demonstrated allelopathic properties, making them effective as cover crops or rotational choices. Studies, such as Weston and Duke (2003), emphasize the potential of allelopathic plants in integrated weed management systems. While promising, their effectiveness may vary with soil type, climate, and the specific weed-crop interactions. Soil microbial activity also plays a role in biological weed control. By promoting beneficial microorganisms, such as mycorrhizal fungi or rhizobacteria, farmers can enhance soil health and suppress weed proliferation indirectly. Microbial herbicides, derived from naturally occurring organisms, are being developed as eco-friendly alternatives to synthetic herbicides. These bioherbicides, such as those based on *Phytophthora palmivora*, show potential for managing persistent weed species while reducing chemical inputs. The environmental impact of biological weed control is highly favorable, as these methods align with natural ecosystem processes and reduce dependency on synthetic herbicides. They enhance biodiversity, improve soil health, and pose minimal risks to non-target species when implemented carefully. However, the slower action of biological methods compared to chemical or mechanical approaches can be a limitation, particularly in high-weed-pressure scenarios. Economically, biological methods can be cost-effective in the long term. Initial investments in biocontrol agent research, propagation, or allelopathic crop establishment may be higher, but these costs are offset by reduced reliance on herbicides and improved crop and soil health. However, the site-specific nature of many biological methods may require tailored solutions, which can increase complexity and costs for large-scale implementation. Compared to cultural, mechanical, and chemical methods, biological weed control provides a sustainable and environmentally sound option but often requires integration with other strategies for maximum effectiveness. For example, using allelopathic plants as part of a cover cropping system alongside mechanical weed removal can provide comprehensive control with reduced ecological impact.

Integrated Weed Management (IWM)

Integrated Weed Management (IWM) is a comprehensive approach to weed control that combines cultural, mechanical, chemical, and biological methods to achieve effective and sustainable management. Unlike relying on a single method, IWM integrates these strategies to address weeds at different growth stages and tackle diverse weed species. This approach reduces reliance on chemical herbicides, mitigating environmental risks such as soil and water contamination while also addressing the growing issue of herbicide resistance. Practices like crop rotation, cover cropping, and inter-row cultivation work alongside selective herbicide applications to optimize weed control. IWM has demonstrated its effectiveness in various studies, showing significant reductions in weed biomass and seedbanks while maintaining crop yields. It also enhances soil health and biodiversity, making it a more sustainable option compared to chemical-intensive methods. Economically, while IWM may involve higher initial costs due to investments in equipment or biocontrol agents, its long-term benefits include lower herbicide usage, improved crop health, and reduced resistance-related expenses. Research has consistently highlighted the advantages of IWM, with studies emphasizing its role in maintaining ecological balance, ensuring economic feasibility, and enhancing crop productivity. Despite challenges like increased complexity and labor requirements, IWM offers a balanced, adaptive, and resilient framework for modern weed management in soybean fields, aligning with the goals of sustainable agriculture.

Conclusion

In conclusion, effective weed management in soybean fields is critical for ensuring optimal crop yield, economic viability, and environmental sustainability. Each weed control method cultural, mechanical, chemical, and biological has distinct strengths and limitations. While chemical methods offer quick and reliable results, they come with risks of herbicide resistance and environmental degradation. Cultural and biological methods provide sustainable alternatives, enhancing soil health and biodiversity but requiring longer implementation periods. Mechanical methods offer immediate weed suppression but may impact soil structure and demand significant labor. Integrated Weed Management (IWM) emerges as the most effective strategy, combining the strengths of diverse approaches to achieve holistic weed control. By reducing reliance on any single method and adapting practices to specific field conditions, IWM ensures long-term productivity and sustainability. Research highlights the importance of adopting such multifaceted approaches to address the complex challenges posed by weeds. Future advancements in technology, farmer education, and policy support will further enhance the efficiency and adoption of integrated weed management systems, paving the way for resilient agricultural practices.

References

1. Liebman M, Dyck E. Crop rotation and intercropping strategies for weed management. *Ecol Appl.* 1993;3(1):92-122. Available from: <https://doi.org/10.2307/1941795>
2. Teasdale JR, Devine TE, Mosjidis JA, Bellinder RR, Beste CE. Weed suppression by hairy vetch in no-

tillage corn. *Weed Sci.* 2007;55(3):219-25. Available from: <https://doi.org/10.1614/ws-06-114.1>

3. Knezevic SZ, Evans SP, Mainz M. Row spacing influences the critical timing for weed removal in soybean (*Glycine max*). *Weed Technol.* 2002;17(4):666-673. Available from: [https://doi.org/10.1614/0890-037X\(2003\)017\[0666:RSITCT\]2.0.CO;2](https://doi.org/10.1614/0890-037X(2003)017[0666:RSITCT]2.0.CO;2)
4. Triplett GB, Dick WA. No-tillage crop production: A revolution in agriculture! *Agron J.* 2008;100(3):S-153-S-165. Available from: <https://doi.org/10.2134/agronj2007.0005c>
5. Culpepper AS, Grey TL, Vencill WK, Kichler JM, Webster TM, Brown SM, *et al.* Glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) confirmed in Georgia. *Weed Sci.* 2004;54(6):620-626. Available from: <https://doi.org/10.1614/ws-04-015r2.1>
6. Powles SB, Yu Q. Evolution in action: Plants resistant to herbicides. *Annu Rev Plant Biol.* 2010;61:317-347. Available from: <https://doi.org/10.1146/annurev-arplant-042809-112119>
7. Weston LA, Duke SO. Weed and crop allelopathy. *Crit Rev Plant Sci.* 2003;22(3-4):367-389. Available from: <https://doi.org/10.1080/713610858>
8. Nuru Seid Tehulie, Tenalem Misgan, Tigist Awoke. Review on weeds and weed controlling methods in soybean (*Glycine max L.*). *J Curr Res Food Sci.* 2021;2(1):1-6.
9. Shaner DL. Lessons learned from the history of herbicide resistance. *Weed Sci.* 2014;62(2):427-31. Available from: <https://doi.org/10.1614/ws-d-13-00062.1>
10. Charudattan R. Biological control of weeds by means of plant pathogens: Significance for integrated weed management in modern agro-ecology. *Biocontrol.* 2001;46(2):229-260. Available from: <https://doi.org/10.1023/A:1011474521182>
11. Norsworthy JK, Ward SM, Shaw DR, Llewellyn RS, Nichols RL, Webster TM, *et al.* Reducing the risks of herbicide resistance: Best management practices and recommendations. *Weed Sci.* 2012;60(1):31-62. Available from: <https://doi.org/10.1614/ws-d-11-00155.1>
12. Pannell DJ, Llewellyn RS, Corbeels M. Herbicide resistance: Economic and environmental challenges. *Agric Syst.* 2004;79(2):159-161. Available from: <https://doi.org/10.1016/j.agsy.2004.06.001>.