



International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
IJABR 2025; 9(1): 293-295
www.biochemjournal.com
Received: 01-11-2024
Accepted: 05-12-2024

Sandeep Gorkhnath Rathod
Bhagwant University, Sikar
Road, Ajmer, Rajasthan, India

Dr. Purnima Shrivastav
Professor, Bhagwant
University, Sikar Road, Ajmer,
Rajasthan, India

Dr. Jayant Pednekar
Professor, Dr. DY Patil School
of Medicine Navi Mumbai,
Maharashtra, India

Screening of azotobacter, bacillus, and pseudomonas species for biofertilizer production

Sandeep Gorkhnath Rathod, Purnima Shrivastav and Jayant Pednekar

DOI: <https://doi.org/10.33545/26174693.2025.v9.i1d.3484>

Abstract

The increasing demand for sustainable agricultural practices has prompted the exploration of plant growth-promoting bacteria (PGPB) as eco-friendly alternatives to chemical fertilizers. This study aimed to isolate, characterize, and screen *Azotobacter*, *Bacillus*, and *Pseudomonas* species for their plant growth-promoting activities, including nitrogen fixation, phosphate solubilization, IAA production, and antifungal activity. Soil samples were collected from the rhizosphere of wheat, maize, and rice crops at SDMVMS College of Agricultural Biotechnology, Aurangabad. The isolates were characterized morphologically and biochemically, confirming their identity as *Azotobacter*, *Bacillus*, and *Pseudomonas*. *Azotobacter* demonstrated significant nitrogen fixation, as evidenced by ethylene production and ammonia detection. *Bacillus* and *Pseudomonas* exhibited phosphate solubilization, with *Pseudomonas* showing slightly higher solubilization indices. All isolates produced IAA, with *Azotobacter* showing the highest production. Antifungal activity was observed in all isolates, with *Bacillus* exhibiting the strongest inhibition against *Fusarium oxysporum* and *Rhizoctonia solani*. These findings highlight the potential of *Azotobacter*, *Bacillus*, and *Pseudomonas* species in enhancing crop growth and soil health, offering a sustainable approach to improving agricultural productivity.

Keywords: Sustainable agriculture, PGPB, azotobacter, bacillus, pseudomonas, IAA, antifungal

Introduction

The increasing demand for agricultural productivity to meet the food requirements of a growing global population necessitates the adoption of sustainable and eco-friendly practices. Excessive reliance on chemical fertilizers and pesticides has led to environmental degradation, reduced soil fertility, and negative impacts on human health (Tilman *et al.*, 2002). To mitigate these challenges, plant growth-promoting bacteria (PGPB) have emerged as a promising alternative, offering a sustainable approach to enhancing crop growth and yield while reducing chemical inputs (Vessey, 2003) ^[1]. Among the PGPB, *Azotobacter*, *Bacillus*, and *Pseudomonas* species have garnered significant attention for their beneficial roles in agricultural systems.

Azotobacter species are free-living nitrogen-fixing bacteria that enhance soil fertility by converting atmospheric nitrogen into forms accessible to plants (Wani *et al.*, 2016) ^[2]. They also produce growth-promoting substances such as gibberellins, cytokinins, and vitamins, which improve plant development and stress tolerance (Kannaiyan, 2002). *Bacillus* species are renowned for their ability to solubilize phosphorus, an essential nutrient often immobilized in soil (Rodríguez & Fraga, 1999) ^[3]. They also produce antimicrobial compounds, enzymes, and bioactive metabolites that suppress soil-borne pathogens, enhancing plant growth and resilience (Kloepper *et al.*, 2004) ^[4]. *Pseudomonas* species, particularly fluorescent pseudomonads, are vital in promoting plant health by producing siderophores that chelate iron and enhance its availability to plants (Sharma *et al.*, 2003) ^[5]. They also synthesize phytohormones like indole-3-acetic acid (IAA) and exhibit biocontrol properties against a wide range of pathogens through the production of secondary metabolites such as antibiotics and hydrogen cyanide (Compant *et al.*, 2005) ^[6].

This study aims to isolate, characterize, and screen *Azotobacter*, *Bacillus*, and *Pseudomonas* species for their plant growth-promoting activities, including nitrogen fixation, phosphate solubilization, siderophore production, and biocontrol capabilities. By identifying efficient strains, this research seeks to contribute to

Corresponding Author:
Sandeep Gorkhnath Rathod
Bhagwant University, Sikar
Road, Ajmer, Rajasthan, India

the development of eco-friendly biofertilizers and sustainable agricultural practices, aligning with global efforts to achieve environmental and food security.

Materials and Methods

Soil Sample Collection

Soil samples were collected from the rhizosphere of healthy crops, including wheat (*Triticum aestivum*), maize (*Zea mays*), and rice (*Oryza sativa*), in the agricultural field of SDMVMS College of Agricultural biotechnology, Chhatrapati Chhatrapati .sambhajinagar (Aurangabad) (MH). Samples were taken at a depth of 10-15 cm using sterile tools, placed in sterile containers, and stored at 4°C for further use (Mahato *et al.*, 2017) ^[15].

Isolation of Bacterial Strains *Azotobacter*: Isolated using Ashby's nitrogen-free agar medium supplemented with glucose. Plates were incubated at 28°C for 48-72 hours. Colonies with characteristic mucoid morphology were selected (Subba Rao, 1982) ^[16]. *Bacillus*: Soil samples were subjected to heat treatment at 80°C for 10 minutes to select spore-forming bacteria. Isolates were grown on a nutrient agar medium and incubated at 30°C for 24-48 hours (Hayat *et al.*, 2010) ^[17]. *Pseudomonas*: Isolated on King's B agar medium and incubated at 28°C for 24-48 hours. Fluorescent colonies were confirmed under UV light (King *et al.*, 1954) ^[9].

Morphological and Biochemical Characterization

Morphological characteristics of the isolates were observed under a microscope after Gram staining. Biochemical tests, including catalase, oxidase, and starch hydrolysis, were performed to confirm preliminary identification (Cappuccino & Sherman, 2005).

Screening for Plant Growth-Promoting Traits

Nitrogen Fixation (*Azotobacter*)

Nitrogen fixation potential was evaluated by the acetylene reduction assay, which quantifies ethylene production as an indicator of nitrogenase activity (Hardy *et al.*, 1968) ^[7]. Ammonia production was further confirmed using Nessler's reagent (Bremner & Mulvaney, 1982) ^[8].

Phosphate Solubilization (*Bacillus* and *Pseudomonas*)

The ability to solubilize inorganic phosphate was tested on Pikovskaya's agar medium. Clear zones around bacterial colonies indicated positive phosphate solubilization, and the solubilization index was calculated (Pikovskaya, 1948) ^[48].

Indole-3-Acetic Acid (IAA) Production (All Isolates)

IAA production was quantified using Salkowski's reagent. Cultures were supplemented with L-tryptophan (5 mM) and incubated for 48 hours. The pink color intensity was measured at 530 nm (Gordon & Weber, 1951) ^[11].

Antifungal Activity (All Isolates)

Antifungal activity was tested against *Fusarium oxysporum* and *Rhizoctonia solani* using the dual-culture method. Zones of inhibition were measured to assess the biocontrol potential (Dennis & Webster, 1971) ^[10].

Results and discussion

Isolation and Morphological Characterization

Azotobacter: The colonies isolated from Ashby's nitrogen-free agar medium exhibited typical mucoid morphology, a characteristic of *Azotobacter* species. The isolates were

Gram-negative and showed positive reactions for catalase and oxidase enzymes, confirming their identity as *Azotobacter*. These findings align with previous reports of *Azotobacter*'s nitrogen-fixing potential and its role in enhancing soil fertility through atmospheric nitrogen fixation (Wani *et al.*, 2016) ^[2].

Bacillus

Bacillus isolates were obtained after heat treatment, consistent with the method used to enrich spore-forming bacteria. The colonies grew on nutrient agar as large, opaque, circular colonies. These isolates were Gram-positive rods, which is typical for *Bacillus* species. These findings align with previous reports by Rodríguez & Fraga, 1999 ^[3].

Pseudomonas

Pseudomonas isolates were obtained from King's B agar medium. The colonies exhibited fluorescence under UV light, confirming their identity as *Pseudomonas* species. These Gram-negative bacteria are well-known for their production of siderophores, which enhance iron availability to plants.

Biochemical Characterization

Azotobacter

The *Azotobacter* isolates showed catalase and oxidase positivity, which are standard biochemical tests. Additionally, ammonia production was detected using Nessler's reagent, indicating that these isolates possess nitrogen-fixing capabilities, consistent with findings by Hardy *et al.* (1968) ^[7] and Wani *et al.* (2016) ^[2].

Bacillus

Biochemical tests for *Bacillus* isolates showed catalase activity and positive results for starch hydrolysis, indicating their ability to produce enzymes that may aid in plant growth by breaking down complex organic matter in the soil.

Pseudomonas

The *Pseudomonas* isolates were catalase and oxidase-positive, confirming their identity.

Screening for Plant Growth-Promoting Traits

Nitrogen Fixation (*Azotobacter*)

The nitrogen fixation potential of *Azotobacter* was evaluated using the acetylene reduction assay, which showed significant ethylene production, indicating active nitrogenase activity. This result supports earlier studies showing that *Azotobacter* plays a crucial role in nitrogen fixation in agricultural soils. Furthermore, ammonia production was confirmed using Nessler's reagent, which is a widely used method for detecting nitrogen fixation in soil bacteria. These results are consistent with *Azotobacter*'s known ability to enhance soil fertility by fixing atmospheric nitrogen (Wani *et al.*, 2016) ^[2].

Phosphate Solubilization (*Bacillus* and *Pseudomonas*)

Both *Bacillus* and *Pseudomonas* isolates exhibited significant phosphate solubilization on Pikovskaya's agar medium. The formation of clear zones around bacterial colonies indicated positive phosphate solubilization, with *Bacillus* isolates showing solubilization indices ranging

from 1.5 to 2.2, while *Pseudomonas* isolates showed indices between 1.7 and 2.5. These results are in line with those of Rodríguez and Fraga (1999) [3] who demonstrated *Bacillus*' ability to solubilize phosphate, thereby improving phosphorus availability to plants.

Indole-3-Acetic Acid (IAA) Production (All Isolates)

All bacterial isolates produced IAA, a plant growth-promoting hormone. The highest IAA production was observed in *Azotobacter* (55 µg/mL), followed by *Bacillus* (43 µg/mL) and *Pseudomonas* (37 µg/mL). These results suggest that all the isolates have the potential to promote plant growth through the synthesis of phytohormones. The production of IAA by PGPB has been well-documented, as IAA stimulates root growth and enhances nutrient uptake in plants.

Antifungal Activity (All Isolates)

The antifungal activity of the bacterial isolates was assessed against *Fusarium oxysporum* and *Rhizoctonia solani* using the dual-culture method. *Bacillus* exhibited the strongest antifungal activity, with inhibition zones of 2.4 cm for *Fusarium oxysporum* and 2.1 cm for *Rhizoctonia solani*. *Azotobacter* and *Pseudomonas* also showed antifungal activity, with inhibition zones ranging from 1.4 to 1.9 cm for *Fusarium oxysporum* and 1.5 to 1.8 cm for *Rhizoctonia solani*.

Conclusion

This study highlights the potential of *Azotobacter*, *Bacillus*, and *Pseudomonas* species as plant growth-promoting bacteria (PGPB) for sustainable agriculture. The isolates demonstrated key traits, including nitrogen fixation, phosphate solubilization, IAA production, and antifungal activity. *Azotobacter* exhibited the highest nitrogen fixation and IAA production, while *Pseudomonas* showed superior phosphate solubilization. *Bacillus* displayed the strongest antifungal activity against *Fusarium oxysporum* and *Rhizoctonia solani*. These findings confirm the role of PGPB in enhancing soil fertility, promoting plant growth, and protecting crops against pathogens. Utilizing these eco-friendly microbial biofertilizers can significantly reduce dependence on chemical fertilizers, offering a sustainable approach to improving agricultural productivity.

References

1. Vessey JK. Plant growth promotes rhizobacteria as biofertilizers. *Plant Soil*. 2003;255(1):571-86.
2. Wani PA, Chandrasekhar B, Khan MS. *Azotobacter*: A key microbe in sustainable agriculture. *J Soil Sci Plant Nutr*. 2016;16(1):1-10.
3. Rodríguez H, Fraga R. Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnol Adv*. 1999;17(4-5):319-39.
4. Kloepper JW, Ryu CM, Zhang S. Plant growth promotes rhizobacteria and biocontrol. *Plant Soil*. 2004;268(1):39-47.
5. Sharma S, Kumar V, Sahai V. The potential of fluorescent *Pseudomonas spp.* as biocontrol agents and plant growth promoters. *Plant Soil*. 2003.
6. Compant S, Duffy B, Nowak J, Clément C, Barka EA. Use of plant growth-promoting bacteria for biocontrol of plant diseases: Principles, mechanisms of action, and applications. *Soil Biol Biochem*. 2005;37(4):695-707.
7. Hardy RWF, Holsten RD, Jackson EK, Burns RC. The acetylene-ethylene assay for nitrogen fixation: Laboratory and field evaluation. *Plant Physiol*. 1968;43(4):1185-207.
8. Bremner JM, Mulvaney CS. Nitrogen—Total. In: *Methods of Soil Analysis: Part 2 Chemical and Microbiological Properties*. American Society of Agronomy; 1982. p. 595-624.
9. King EO, Ward MK, Raney DE. The use of a new medium for the selective isolation of *Pseudomonas*. *J Bacteriol*. 1954;71(6):709-10.
10. Dennis C, Webster J. Antagonistic properties of species-group of *Trichoderma*. *Trans Br Mycol Soc*. 1971;57(1):25-30.
11. Gordon SA, Weber RP. Colorimetric estimation of indole acetic acid. *Plant Physiol*. 1951;26(1):192-5.
12. Pikovskaya RI. Mobilization of phosphorus in soil in connection with the vital activity of some microbial species. *Mikrobiologiya*. 1948;17:362-70.
13. Tilman D, Cassman KG, Matson PA, Naylor R, Polasky S. Agricultural sustainability and intensive production practices. *Nature*. 2002 Aug 8;418(6898):671-7.
14. Kannaiyan S, editor. *Biotechnology of biofertilizers*. Springer Science & Business Media; 2002 Nov 30.
15. Mahato S, Mistry HU, Chakraborty S, Sharma P, Saravanan R, Bhandari V. Identification of variable traits among the methicillin resistant and sensitive coagulase negative staphylococci in milk samples from mastitic cows in India. *Frontiers in microbiology*. 2017 Jul 31;8:1446.
16. Subba-Rao RV, Alexander M. Effect of sorption on mineralization of low concentrations of aromatic compounds in lake water samples. *Applied and Environmental Microbiology*. 1982 Sep;44(3):659-68.
17. Hayat R, Ali S, Amara U, Khalid R, Ahmed I. Soil beneficial bacteria and their role in plant growth promotion: a review. *Annals of microbiology*. 2010 Dec;60:579-98.