



International Journal of Research in Agronomy

E-ISSN: 2618-0618

P-ISSN: 2618-060X

© Agronomy

www.agronomyjournals.com

2024; 7(12): 960-964

Received: 26-10-2024

Accepted: 29-11-2024

Amol R Pawar

Ph.D. Scholar, Department of
Agronomy, Lovely Professional
University, Phagwara, Punjab, India

Sangeeta Kumari

Assistant Professor, Department of
Botany RPS Degree College, Balana,
Mahendergarh, Haryana, India

Kartikey Sootrakar

Teaching Assistant,
Department of Genetics and Plant
Breeding Maharaja Chhatrasal
Bundelkhand University,
Chhatarpur, Madhya Pradesh, India

Shivanand Hiremath

Scientist Plant Protection, ICAR
KLE KVK, Mattikoppa, Karnataka,
India

Neha Singh

Assistant Professor, Department of
Zoology HVM (P.G.) College, Raisi,
Haridwar affiliated to Sri Dev Suman
Uttarakhand University,
Badshahithaul, Uttarakhand, India

Akshita Awasthi

Ph.D. Research Scholar, Department
of Plant Breeding and Genetics,
Punjab Agricultural University,
Ludhiana, Punjab, India

Kalisetty Hanuma Sai Dinesh

M.S Agriculture for Sustainable
Development, Research scholar
Department of Agriculture
University of Greenwich, Kent,
United Kingdom (ME4 4TB)

Thiruvengadam K

Assistant Professor in Entomology,
Department of Agricultural
Entomology, RVS Agricultural
College, Thanjavur, Tamil Nadu,
India

Corresponding Author:

Amol R Pawar

Ph.D. Scholar, Department of
Agronomy, Lovely Professional
University, Phagwara, Punjab, India

Advancing integrated pest management: Strategies for minimizing pesticide use

Amol R Pawar, Sangeeta Kumari, Kartikey Sootrakar, Shivanand Hiremath, Neha Singh, Akshita Awasthi, Kalisetty Hanuma Sai Dinesh and Thiruvengadam K

DOI: <https://doi.org/10.33545/2618060X.2024.v7.i12l.2359>

Abstract

Integrated Pest Management (IPM) is a holistic approach to pest control that emphasizes ecological principles and minimizes reliance on synthetic pesticides. This paper provides a concise overview of the key strategies for advancing IPM and minimizing pesticide use. It emphasizes the need for a multifaceted approach that combines various control methods and considers the ecological and economic implications of pest management decisions. The key strategies for advancing IPM and reducing pesticide use, including: Habitat Manipulation, Cultural Controls, Biological Control, Monitoring and Early Detection, Pesticide Use Optimization. This paper highlights the importance of interdisciplinary research, farmer education, and policy support to effectively implement IPM strategies and minimize pesticide use, leading to more sustainable and environmentally friendly agricultural practices.

Keywords: Integrated Pest Management (IPM), pesticide reduction, habitat manipulation, biological control, cultural control, monitoring, sustainable agriculture

Introduction

Integrated Pest Management (IPM)

Integrated Pest Management (IPM) is a holistic approach to pest control that emphasizes ecological principles and sustainable practices. It is a comprehensive approach to pest control that aims to minimize pesticide use while effectively managing pest populations (Barzman *et al.*, 2015) [4]. It involves a combination of strategies, including cultural, physical, biological, and chemical methods. Since the initiation of "integrated control," which is defined as "applied pest control which combines and integrates biological and chemical control" (Smith *et al.* 1959) [13], IPM has advanced significantly. Entomologists first came up with the idea in response to widespread use of broad-spectrum insecticides, insect outbreaks brought on by the eradication of natural enemies, and the rise of pesticide resistance (Barzman *et al.*, 2015) [4]. Today, IPM covers every facet of plant protection. The definition of IPM that is primarily based on the one provided by the Food and Agriculture Organization of the United Nations, adding the concept of "ecological justification" to that of "economic justification" and replacing the term "pest control techniques" with "plant protection methods".

However, IPM is a multifaceted method that involves multiple economic sectors and draws from a variety of disciplines. Producing specific recommendations that are applicable to the diversity of global agriculture is a challenging task (Barzman *et al.* 2014) [3]. In order to keep pest populations below economically damaging levels while reducing risks to the environment and public health, IPM is a comprehensive, ecological approach to managing pests in agricultural systems. It involves the strategic integration of multiple control methods, including cultural, biological, and chemical tactics (Angon *et al.*, 2023) [1]. Rather than depending exclusively on reactive pesticide treatments, IPM stresses the use of preventive measures, monitoring, and decision-making based on defined criteria (Rossi *et al.*, 2023; Deguine *et al.*, 2021) [10, 6]. IPM includes: preventing pest issues through cultural practices like crop production and sanitation; tracking pest populations and their natural enemies; utilizing economic thresholds to inform

management choices; utilizing a combination of chemical, physical, and biological control methods; and assessing the efficacy of interventions (Scheff and Phillips, 2022; Han *et al.*, 2024) ^[12, 7]. IPM aims to maximize agricultural productivity while maintaining ecosystem services and reducing the negative effects of conventional pesticide dependence by combining multidisciplinary knowledge and a systems-based approach (Saroop and Tamchos, 2024; Baker *et al.*, 2020) ^[11, 2].

General principles of IPM (Barzman *et al.*, 2015) ^[4].

The general principles of IPM are shown in Fig. 1 and are as follows:




Fig 1: General principles of IPM

1. Prevention and suppression

The prevention and/or suppression of harmful organisms can be achieved or supported by:

- Crop rotation
- Use of adequate cultivation techniques
- Use of balanced fertilization, liming and irrigation/drainage practices,
- Use, where appropriate, of resistant/tolerant cultivars and standard/ certified seed and planting material,
- Preventing the spreading of harmful organisms by hygiene measures (e.g., by regular cleansing of machinery and equipment),

2. Protection and enhancement of important beneficial organisms

- By adequate plant protection measures
- Utilization of ecological infrastructures inside and outside production sites.

3. Monitoring Harmful organisms must be monitored

- observations in the field,
- scientifically sound warning,
- forecasting and early diagnosis systems, where feasible,
- Use of advice from professionally qualified advisors.

4. Decision-making

- Based on the results of the monitoring the professional user has to decide whether and when to apply plant protection measures.
- Robust and scientifically sound threshold values

- For harmful organisms, threshold levels defined for the region, specific areas, crops and particular climatic conditions must be taken into account before treatment.

5. Non-chemical methods

- Sustainable biological, physical and other non-chemical methods must be chosen than chemical methods if they provide reasonable pest control.

6. Pesticide selection

- The pesticides applied shall be as specific as possible for the target
- shall have the least side effects on human health, non-target organisms and the environment.

7. Reduced pesticide use

- by reduced doses,
- reduced application frequency or partial applications,
- considering that the level of risk in vegetation is acceptable and
- they do not increase the risk for development of resistance in populations of harmful organisms.

8. Anti-resistance strategies

- Use of multiple pesticides with different modes of action.

9. Evaluation

- should check the success of the applied plant protection measures

Key Strategies for Minimizing Pesticide Use in IPM (Zhou *et al.*, 2024; Barzman *et al.*, 2015) ^[14, 4] (Fig. 2):




Fig 2: Key strategies for minimizing pesticide use in IPM

1. **Cultural Controls:** Employing agricultural practices that minimize pest infestations. This includes techniques like proper sanitation, timely planting and harvesting, and the use of pest-resistant varieties (Ahuja *et al.*, 2015; Leskey *et al.*, 2020) ^[16, 17]
- **Crop Rotation:** Planting different crops in sequence can disrupt the life cycles of pests and reduce their populations.
- **Sanitation:** Removing weeds, debris, and other potential breeding grounds for pests can help prevent infestations.
- **Healthy Plants:** Providing adequate water, nutrients, and sunlight can help plants resist pests and diseases.
- **Planting pest-resistant varieties:** Reduces the need for pesticides.
- **Proper irrigation:** Avoids creating conditions that favor pests, such as excessive moisture.

- **Row covers:** Protect plants from insect pests and can improve pollination.
- **Timing of planting and harvesting:** Can help avoid periods of peak pest activity.

Pros

- Relatively easy to implement
- Can be effective for a wide range of pests
- Can improve overall crop health
- Often low-cost or no-cost

Cons

- May not be effective for all pests
- May require ongoing management
- Effectiveness can vary depending on environmental conditions

2. Physical Controls (Ragunathan and Divakar, 2020)^[15]

Regular monitoring of pest populations and early detection of infestations are crucial for effective IPM. This allows for timely intervention with the least disruptive control methods.

- **Visual inspections:** Regularly checking plants for signs of pests and damage.
- **Scouting:** Conducting regular field surveys to identify problem areas.
- **Sampling techniques:** Collecting and analyzing samples of plants or soil to detect pests.
- **Handpicking:** Manually removing pests from plants can be effective for small infestations.
- **Trapping:** Using traps to capture and kill pests, such as sticky traps for insects or snap traps for rodents.
- **Exclusion:** Preventing pests from entering areas by using barriers, such as screens or netting.

Pros

- Allows for timely intervention
- Can prevent infestations from reaching damaging levels
- Minimizes the need for broad-spectrum pesticides
- Can reduce the risk of pesticide resistance

Cons

- Requires ongoing effort and resources
- May require specialized training for personnel
- Effectiveness depends on accurate and timely monitoring

3. Biological Controls (Alam *et al.*, 2016; Baker *et al.*, 2020; Leskey *et al.*, 2020)^[19, 2, 17]

Utilizing natural enemies of pests, such as predators, parasites, and pathogens, to suppress pest populations. This involves the conservation and augmentation of existing beneficial organisms and the introduction of new biocontrol agents.

- **Conservation biological control:** Protecting and enhancing existing populations of natural enemies.
- **Augmentation biological control:** Introducing additional natural enemies to an area.
- **Inoculative releases:** Introducing small numbers of natural enemies to establish a permanent population.
- **Inundative releases:** Releasing large numbers of natural enemies for immediate pest control.
- **Microbial pesticides:** Using bacteria, fungi, or viruses to kill pests.

Pros

- Highly specific to target pests
- Can be very effective in reducing pest populations
- Environmentally friendly
- Can reduce reliance on chemical pesticides

Cons

- Can be expensive to establish and maintain
- May require specialized knowledge and expertise
- Effectiveness can be influenced by environmental factors

4. Chemical Controls

- **Pesticide Use as a Last Resort:** Only using pesticides when other methods are ineffective and when the potential benefits outweigh the risks.
- **Selective Pesticides:** Choosing pesticides that are specific to the target pest and have minimal impact on beneficial organisms.
- **Integrated Pest Management (IPM) programs:** Using a combination of cultural, physical, and biological controls to reduce the need for pesticides.

5. Habitat Manipulation (Ahuja *et al.*, 2015; Paez Chavez *et al.*, 2018)^[16, 18]

Altering the environment to make it less favorable for pests. This includes techniques like crop rotation, intercropping, and the use of cover crops to disrupt pest life cycles and promote beneficial organisms. Examples:

- **Planting flowering plants:** Attracts beneficial insects like bees, butterflies, and ladybugs, which prey on pests.
- **Creating hedgerows:** Provides shelter and nesting sites for birds, bats, and other predators.
- **Crop rotation:** Disrupts the life cycles of pests that are specific to certain crops.
- **Cover crops:** Can suppress weeds, improve soil health, and provide habitat for beneficial organisms.

Pros

- Environmentally friendly
- Cost-effective in the long term
- Can enhance biodiversity
- Improves soil health and water quality

Cons

- May require significant upfront investment
- Effectiveness can vary depending on the specific pest and ecosystem
- Requires careful planning and management

6. Pesticide Use Reduction

When pesticides are necessary, IPM emphasizes the use of the least toxic and most selective options. This includes the use of reduced-risk pesticides, targeted applications, and the development of pest management decision support systems (Ahuja *et al.*, 2015; Alam *et al.*, 2016; Ragunathan and Divakar, 2020)^[16, 19, 15]. Examples:

- **Using selective pesticides:** Targeting specific pests while minimizing harm to beneficial organisms.
- **Applying pesticides accurately and efficiently:** Reducing the amount of pesticide used and preventing drift.
- **Adhering to integrated pest management thresholds:** Only applying pesticides when the pest population exceeds

a certain level.

- **Using reduced-risk pesticides:** Choosing pesticides with lower environmental impact.

Pros

- Can provide effective control when necessary
- Minimizes the risk of environmental contamination and human exposure
- Reduces the development of pesticide resistance

Cons

- Requires careful planning and implementation
- May still have some negative environmental impacts
- Requires knowledge of pesticide use and safety

Comparing different Strategies of IPM

IPM encompasses a variety of strategies to minimize pesticide use. Habitat manipulation focuses on modifying the environment to favor beneficial organisms and deter pests, offering long-term benefits like biodiversity enhancement but requiring upfront investment (Ragunathan and Divakar, 2020) ^[15]. Cultural controls involve adjusting agricultural practices, such as planting resistant varieties or improving sanitation, for relatively easy and often cost-effective pest prevention (Ahuja *et al.*, 2015) ^[16]. Biological control leverages natural enemies, like predators and parasites, for highly specific and environmentally friendly control, though it may require specialized knowledge and can be costly (Alam *et al.*, 2016) ^[19]. Monitoring and early detection enable timely interventions, minimizing the need for broad-spectrum pesticides, but require consistent effort and potentially specialized training (Paez Chavez *et al.*, 2018) ^[18]. Finally, pesticide use optimization emphasizes judicious use of pesticides when necessary, prioritizing selective options and adhering to thresholds to minimize environmental impact and reduce resistance risks (Leskey *et al.*, 2020) ^[17]. Choosing the right IPM strategy depends on a variety of factors, including the specific pest, the crop being grown, the environmental conditions, and the available resources. Often, the most effective approach is to combine multiple strategies to create a comprehensive IPM strategy.

Benefits of Advancing IPM

- **Reduced Environmental Impact:** Minimizing pesticide use protects beneficial organisms, reduces water pollution, and conserves natural resources.
- **Improved Human Health:** Reducing exposure to pesticides can improve human health and safety (Boudh and Singh, 2019 and Rejesus and Jones, 2020) ^[5, 9].
- **Sustainable Agriculture:** IPM practices promote long-term sustainability of agricultural systems.
- **Cost-Effectiveness:** IPM can be more cost-effective in the long run compared to relying solely on pesticides.
- **Enhanced agricultural sustainability:** Promotes long-term productivity and profitability by reducing pest-related losses and improving soil health (Narayanan *et al.*, 2022 and Rejesus and Jones, 2020) ^[8, 9].
- **Improved food quality:** Reduces pesticide residues on crops, ensuring safer and healthier food for consumers (Boudh and Singh, 2019; Narayanan *et al.*, 2022 and Rejesus and Jones, 2020) ^[5, 8, 9].
- **Increased biodiversity:** Supports the conservation of beneficial organisms and promotes a healthier agroecosystem (Boudh and Singh, 2019 and Rejesus and

Jones, 2020) ^[5, 9].

Challenges and Future Directions

- **Developing and Implementing IPM Programs:** Requires knowledge, training, and resources for farmers and other stakeholders.
- **Addressing Pest Resistance:** Pests can develop resistance to pesticides, necessitating the development of new control methods.
- **Integrating New Technologies:** Incorporating new technologies, such as precision agriculture and remote sensing, to improve IPM decision-making.
- **Promoting Public Awareness:** Raising awareness among the public about the benefits of IPM and encouraging its adoption.

Conclusion

Advancing IPM through the implementation of these strategies is essential for achieving sustainable agriculture and mitigating the negative impacts of pesticide use. Continued research, development, and implementation of innovative IPM approaches are crucial for addressing the challenges of pest management in a changing world. By advancing IPM strategies, we can create a more sustainable and resilient agricultural system that minimizes pesticide use and protects both human and environmental health.

References

1. Angon PB, Mondal S, Jahan I, Datto M, Antu UB, Ayshi FJ, *et al.* Integrated pest management (IPM) in agriculture and its role in maintaining ecological balance and biodiversity. *Adv Agric.* 2023;2023(1):5546373.
2. Baker BP, Green TA, Loker AJ. Biological control and integrated pest management in organic and conventional systems. *Biol Control.* 2020;140:104095.
3. Barzman MS, Bertschinger L, Dachbrodt-Saaydeh S, Graf B, Jensen JE, Joergensen LN, *et al.* Integrated Pest Management policy, research and implementation: European initiatives. In: *Integrated Pest Management: Experiences with Implementation, Global Overview*, Vol. 4. 2014. p. 415-428.
4. Barzman M, Bàrberi P, Birch ANE, Boonekamp P, Dachbrodt-Saaydeh S, Graf B, *et al.* Eight principles of integrated pest management. *Agron Sustain Dev.* 2015;35(4):1199-1215.
5. Boudh S, Singh JS. Pesticide contamination: environmental problems and remediation strategies. In: *Emerging and eco-friendly approaches for waste management*. 2019. p. 245-269.
6. Deguine JP, Aubertot JN, Flor RJ, Lescourret F, Wyckhuys KA, Ratnadass A. Integrated pest management: good intentions, hard realities. A review. *Agron Sustain Dev.* 2021;41(3):38.
7. Han P, Rodriguez-Saona C, Zalucki MP, Liu SS, Desneux N. A theoretical framework to improve the adoption of green Integrated Pest Management tactics. *Commun Biol.* 2024;7(1):337.
8. Narayanan M, Kandasamy S, He Z, Kumarasamy S. Ecological impacts of pesticides on soil and water ecosystems and its natural degradation process. In: *Pesticides in the natural environment*. Elsevier; 2022. p. 23-49.
9. Rejesus RM, Jones MS. Perspective: enhancing economic evaluations and impacts of integrated pest management farmer field schools (IPM-FFS) in low-income countries.

- Pest Manag Sci. 2020;76(11):3527-3536.
10. Rossi V, Caffi T, Salotti I, Fedele G. Sharing decision-making tools for pest management may foster implementation of Integrated Pest Management. Food Sec. 2023;15(6):1459-1474.
 11. Saroop S, Tamchos S. Impact of pesticide application: Positive and negative side. In: Pesticides in a Changing Environment. 2024. p. 155-178.
 12. Scheff DS, Phillips TW. Integrated pest management. In: Storage of Cereal Grains and Their Products. Woodhead Publishing; c2022. p. 661-675.
 13. Smith R, Van Den Bosch R, Hagen K, Stern V. The integration of chemical and biological control of the spotted alfalfa aphid: the integrated control concept. Hilgardia. 1959;29(2):81-101.
 14. Zhou W, Arcot Y, Medina RF, Bernal J, Cisneros-Zevallos L, Akbulut ME. Integrated pest management: an update on the sustainability approach to crop protection. ACS Omega. 2024;9(40):41130-41147.
 15. Ragunathan V, Divakar BJ. Integrated pest management strategies. In: Molecular Biology of the Biological Control of Pests and Diseases of Plants. CRC Press; 2020. p. 173-196.
 16. Ahuja DB, Ahuja UR, Singh SK, Singh N. Comparison of integrated pest management approaches and conventional (non-IPM) practices in late-winter-season cauliflower in Northern India. Crop Prot. 2015;78:232-238.
 17. Leskey TC, Short BD, Ludwick D. Comparison and refinement of integrated pest management tactics for *Halyomorpha halys* (Hemiptera: Pentatomidae) management in apple orchards. J Econ Entomol. 2020;113(4):1725-1734.
 18. Paez Chavez J, Jungmann D, Siegmund S. A comparative study of integrated pest management strategies based on impulsive control. J Biol Dyn. 2018;12(1):318-341.
 19. Alam MZ, Haque MM, Islam MS, Hossain E, Hasan SB, Hasan SB, Hossain MS. Comparative study of integrated pest management and farmers practices on sustainable environment in the rice ecosystem. Int J Zool. 2016;2016(1):7286040.