

International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
IJABR 2025; SP-9(1): 489-493
www.biochemjournal.com
Received: 08-10-2024
Accepted: 13-11-2024

Preeti Verma
Physiology and climatology
Division, ICAR- Indian Veterinary
Research Institute, Izatnagar
Bareilly, Uttar Pradesh, India

Susmita Majumder
Physiology and climatology
Division, ICAR- Indian Veterinary
Research Institute, Izatnagar
Bareilly, Uttar Pradesh, India

Lata Kant
Physiology and climatology
Division, ICAR- Indian Veterinary
Research Institute, Izatnagar
Bareilly, Uttar Pradesh, India

Suboni Nunisa
Physiology and climatology
Division, ICAR- Indian Veterinary
Research Institute, Izatnagar
Bareilly, Uttar Pradesh, India

HA Samad
Physiology and climatology
Division, ICAR- Indian Veterinary
Research Institute, Izatnagar
Bareilly, Uttar Pradesh, India

VS Chouhan
Physiology and climatology
Division, ICAR- Indian Veterinary
Research Institute, Izatnagar
Bareilly, Uttar Pradesh, India

Gynendra Singh
Physiology and climatology
Division, ICAR- Indian Veterinary
Research Institute, Izatnagar
Bareilly, Uttar Pradesh, India

VP Maurya
Physiology and climatology
Division, ICAR- Indian Veterinary
Research Institute, Izatnagar
Bareilly, Uttar Pradesh, India

Corresponding Author:
Preeti Verma
Physiology and climatology
Division, ICAR- Indian Veterinary
Research Institute, Izatnagar
Bareilly, Uttar Pradesh, India

A review on: Effect of dietary chromium and selenium supplementation on phsio-biochemical, antioxidant activity and immune function of Vrindavani cattle exposed to thermal stress

Preeti Verma, Susmita Majumder, Lata Kant, Suboni Nunisa, HA Samad, VS Chouhan, Gynendra Singh and VP Maurya

DOI: <https://doi.org/10.33545/26174693.2025.v9.i1Sg.3538>

Abstract

Heat stress has harmful effect on the animal's physio-biochemical, hormonal, antioxidant activity and immunity of animals which hampers the overall productive potential of animals. There is an urgent need to address the problem of heat stress so that the productivity of animals may be maintained up to some extent. Antioxidants are added in animal's diets as part of nutritional management to minimize oxidative damage during heat stress (Chauhan *et al.*, 2020). Now a days a number of feed additives, such as niacin, vitamin A,C, E, selenium (Se) and chromium (Cr) are added in animal's diet to reduce the adverse consequences of heat exposure by preventing oxidative damage in the animals experiencing heat stress (Chauhan *et al.*, 2020). Selenium (Se) is dietary trace element which acts as an essential co-factor of glutathione peroxidase. Dietary selenium supplementation has been demonstrated to improve gastrointestinal health, reproductive physiology, and production performance during heat stress (Zheng *et al.*, 2022). Chromium (Cr) is essential micro mineral which is vital for the metabolism in animals (Lashkari *et al.*, 2018). The dietary supplementations of Cr in heat stressed calves helps in maintaining their health and immune system (Shan *et al.*, 2020). Further research is urgently required to study the combined effect of Se and Cr on productive and reproductive potential of cattle exposed to thermal stress.

Keywords: Heat stress, antioxidants, selenium, chromium, oxidative damage, nutritional management

Introduction

India is the country with the largest dairy herd in the world compared to industrialized countries that produce milk (Rhoads *et al.*, 2009) [34]. The vast climatic condition in India influences the adaptive capabilities and immunity of different milch breeds of cow which intern reduces the milk production. During the heat stress animal adopt various biological mechanism to maintain their internal milieu by reducing their feed intake, altered endocrine status and reduction in rumination (Collier *et al.*, 2006) [11]. Antioxidants are added in animal's diets as part of nutritional management to minimize oxidative damage during heat stress (Chauhan *et al.*, 2020) [7]. Now a days a number of feed additives, such as niacin, vitamin A,C, E, selenium (Se) and chromium (Cr) are added in animal's diet to reduce the adverse consequences of heat exposure by preventing oxidative damage in the animals experiencing heat stress (Chauhan *et al.*, 2020) [7].

Selenium (Se) is a dietary trace element which acts as an essential co-factor of glutathione peroxidase and acting as an antioxidant enzyme which is very essential for the animal's productivity. Dietary selenium supplementation has been demonstrated to improve gastrointestinal health, reproductive physiology, and production performance during heat stress (Zheng *et al.*, 2022) [45]. Chromium (Cr) is essential micro mineral which is vital for the metabolism in animals (Lashkari *et al.*, 2018) [21]. Cr plays a crucial role in the cellular breakdown of fat, protein, and carbohydrates and is a very important component of the glucose metabolism and it also act as an antioxidant role. The dietary supplementations of Cr in heat stressed calves helps in maintaining their health and immune system up to some extent.

It is an affective antioxidant that inhibits the lipid peroxidation caused by heat stress and also increases the activity of insulin in tissues which helps in enhancing animal production (Chouraddi *et al.*, 2022) ^[9]. In this review, we summarized the effect of chromium and selenium feeding on physio-biochemical response, anti oxidant activity and immune function in Vrindavani cattle exposed to thermal stress.

Physiological response status

The various physiological responses like body temperature, heart rate and respiration rate are drastically affected by the fluctuation in the environmental condition. In addition to this these parameters also act as an indicator of severity of environmental stress (Maurya *et al.*, 2020) ^[25].

The change in rectal temperature by 1 °C have its impact on feed intake of animals. (Ranjitkar *et al.*, 2020) ^[33]. A heat stress which generally affects the rectal temperature of the animal and also it has negative impact on milk yield and fertility of animals (Mousavi *et al.*, 2019) ^[28]. The response of RT elevated when ambient temperature was higher. Research indicates that heat stress elevates the body temperature cattle's (Kargar *et al.*, 2018) ^[19]. Studies has indicated that while under heat stress, growing calves and lams have an increased heart rate and rectal temperature (Maurya *et al.*, 2010; Kargar *et al.*, 2018) ^[23, 19]. Mudgal *et al* (2008) ^[29], reported an ameliorative effect of Se and chromium on rectal temperature rise.

The Respiration Rate (RR) is very crucial parameters which used to increase during heat exposure (Maurya *et al.*, 2020) ^[25]. Respiratory rate is an marker of heat stress that is measurable noninvasively (Chrast *et al.*, 2023) ^[10]. Aleena *et al* (2020) ^[1], reported that in a hot environment, breathing rate of animals increases to dissipate heat form the body and to maintain a normal body temperature. (Djelailia *et al.*, 2021) ^[15] reported that the reparation rate of cow increases when they are being exposed to elevated temperature. The supplementation of selenium and chromium reduces the respiration rate during summer as compared to nontreated animals (Srikandakumar *et al.*, 2004) ^[41].

The pulse rate serves as an important indicator of circulatory balance and the overall metabolic state of animals in various environmental conditions (Maurya *et al.*, 2015; Hooda and Upadhyay., 2015) ^[22, 18]. The exposure to heat stress may affects the distribution of blood to peripheral tissues and resulting in increased heat loss through sensible and insensible means respectively (Mohr *et al.*, 2002) ^[27]. In all exposure environments, PR was much greater in crossbred Vrindavani animals than in Tharparkar cattle, possibly due to an acclimation process of a breed that may able to generate excess heat. Beneficial effect of 1.0 ppm of dietary chromium supplementation in 42 °C ambient temperature exposed a calves was reported by (Yari *et al.*, 2010) ^[44]. According to reports a selenium dose of 0.3 mg / kg DMI in the male buffalo calve was sufficient to maintain pulse rate under thermal stress (Shinde *et al.*, 2009) ^[39].

Stress Hormones status

Heat stress drastically affects the kinetics of various endocrine glands, causing the production of various hormones such as cortisol, thyroxine, adrenaline, and nor-adrenaline respectively (Djelailia *et al.*, 2021) ^[15]. The cortisol is a stress hormone produced by adrenal gland and plays a vital role and significant role in immune system

(Degen and young 2002) ^[14]. It has been reported by Shaji *et al* (2017) ^[37], that heat stress stimulates the hypothalamo-pituitary axis which results release glucocorticoids from the adrenal glands and secretion of cortisol as well. Cortisol is a good biomarker for determining the level of stress in different animals (Chen *et al.*, 2023) ^[8]. Ghorbani *et al* (2012) ^[17], observed that Se supplementation to stressed transportation stress or birth stress calves decreased blood cortisol. (Raghunandan *et al.*, 2022) ^[32] reported a decrease in cortisol levels when animal were supplemented with chromium and selenium as compared to non-supplemented animals under thermal stress. The thyroid is responsible for accurately maintaining both body temperature and heat production. Thyroid gland activity varies seasonally and is influenced by surrounding temperature and humidity levels (Maurya *et al.*, 2015) ^[15]. T3 and T4 hormones are involved in thermogenesis in animals and also act as significant indicator for adaptability of animals during stress (Maurya *et al.*, 2018) ^[24]. However (Mousavi *et al.*, 2019) ^[28] reported higher concentration of T3 in heat stressed animal supplemented with Cr and selenium as compared to the non-stressed animals.

Biochemical parameters status

Sejian *et al* (2012) ^[36], reported the considerable decreased the value of albumin and total protein under heat exposure conditions. Garcia *et al.* (2015) ^[16] observed that in heat stressed cow experiencing minor dehydration due to reduced feed intake, there was a notable decrease in albumin levels. Al-Saiady *et al* (2004) ^[3], conducted a study on cow supplemented with chromium and found that there was reduction in the total blood proteins and globulins under heat stress conditions. Contrastingly Yadav *et at.*, (2016) ^[43], reported an rise in the total protein concentration in dairy cattle under to heat stress condition. Mousavi *et al* (2019) ^[28], supplemented Cr and Se in summer exposed dairy calf and found that the concentration of total protein and albumin was higher in supplemented animal as compared non supplemented calves.

The heat stress affects the urea and serum creatinine levels in animals significantly, which also depend upon other factors such as intensity and duration of heat exposure, age, breed and manage mental practices (Saeed *et al.*, 2021) ^[35]. In one study Bernabucci *et al* (2010) ^[4], reported that cross-bred dairy calves exposed to heat stress exhibited higher blood creatinine and blood urea nitrogen (BUN) levels. Cr and Se in feeding animal during heat stress condition reduce the blood creatinine and blood urea nitrogen (BUN) levels (Kargar *et al* 2018; Alhidary *et al.*, 2015) ^[19, 2]. In addition to this Raghunandan *et al.* (2022) ^[32] also reported a decrease level of AST and ALT heat exposed calf as compared to control animals.

Heat Shock Proteins status

Heat Shock Proteins are very important and potent molecular chaperons which help in reducing stress and enhance stress tolerance in farm animals. (Pearl *et al.* 2008). HSP70 is the most sensitive of all the HSPs and is regarded as a crucial regulator of thermal adaptation in animals under heat stress (Dangi *et al.*, 2014) ^[13]. HSP70 helps animals adjust to heat stress by promoting correct folding of recently synthesized proteins and restoring denatured proteins to their original condition (Kishore *et al.*, 2014) ^[20]. Increased HSP70 synthesis cells under heat stress may indicate that

HSP70 is involved in reducing the harmful effect of elevated temperature to preserve goat cell structure and equilibrium (Mishra and Palai 2014) ^[26]. The dietary supplementation Cr and Se reduces the HSP 70 expression in heat exposed cows (Son *et al.*, 2022) ^[40]. In continuation of previous finding (Parashuramulu *et al.*, 2015) ^[30] also reported that Cr and Se feeding reduces the HSP 70 expression. HSP-90 is an important molecular chaperon and highly abundant in the endoplasmic reticulum, cytoplasm and nucleus and is responsible for protein translocation under normal conditions. (Bharti *et al.*, 2017) ^[5]. It is regarded as a flexible chaperone protein that, in the face of heat stress, preserves cellular integrity and homeostasis (Kishore *et al.*, 2014) ^[20]. Dangi *et al.* (2012) ^[12] reported a higher level of HSP-90 in goat exposed to elevated temperature. In heat-exposed cows, the dietary supplements Cr and Se decrease expression of HSP 90 (Son *et al.*, 2022) ^[40]. In continuation of previous finding (Parashuramulu *et al.*, 2015) ^[30] also reported that Cr and Se feeding reduces the HSP 90 expression.

Antioxidants status

Catalase (CAT), total antioxidant capacity (TAC), glutathione peroxidase (GPx), and SOD are the primary components of the intracellular antioxidant defense system (Shaji *et al.*, 2017) ^[37]. The enhanced SOD activity during the summer results in increased production of H₂O₂ and protection from this ROS would only be conferred by the coordinated elevate in the activity of CAT and GPx (Chandra and Aggarwal, 2009) ^[6]. According to Sun *et al.* 2018, GPx activity in calves and dairy cows increase with Dietary Se supplementation. According to Kumbhar *et al.*, 2018, increased SOD activity with Se supplementation have been observed in broilers. During the summer, Tharparkar cattle had the highest total antioxidant activity and catalase (CAT), followed by Sahiwal cattle and then crossbred cattle (Tejaswi *et al.*, 2020) ^[42]. Found that CAT and T-AOC activities increased in the serum, indicating that Cr and Se supplementation improved the antioxidant capacity of heat-stressed dairy cows. (Kargar *et al.*, 2018) ^[19] reported that Cr and Se feeding to heat stressed Holstein calve increases in the activity of catalase and and T- AOC .

Conclusion

The heat stress drastically affects animal production and reproduction. There is an urgent need to have some strategies so that live stock owner can manage their animal's productive potential during thermal stress. The Chromium and Se supplement could serve as a nutritional strategy to help animals to reduce negative effects on their production performances and health during heat stress period. The Cr and Se restores productive performance, antioxidant profile and immune status up to some extent. Therefore, dietary supplementation of Cr and Se may be used as nutritional strategy to minimize the negative effects of heat stress in Vrindavani cattle.

Reference

1. Aleena J, Sejian V, Krishnan G, Bagath M, Pragna P, Bhatta R. Heat stress impact on blood biochemical response and plasma aldosterone level in three different indigenous goat breeds. *Journal of Animal Behaviour and Biometeorology*. 2020;8(4):266-275.
2. Alhidary IA, Shini S, Al Jassim RAM, Abudabos AM, Gaughan JB. Effects of selenium and vitamin E on performance, physiological response, and selenium balance in heat-stressed sheep. *Journal of Animal Science*. 2015;93(2):576-588.
3. Al-Saiady MY, Al-Shaikh MA, Al-Mufarrej SI, Al-Showeimi TA, Mogawer HH, Dirrar A. Effect of chelated chromium supplementation on lactation performance and blood parameters of Holstein cows under heat stress. *Animal Feed Science and Technology*. 2004;117(3-4):223-233.
4. Bernabucci U, Lacetera N, Baumgard LH, Rhoads RP, Ronchi B, Nardone A. Metabolic and hormonal acclimation to heat stress in domesticated ruminants. *Journal of Animal*. 2010;4(7):1167-1183.
5. Bharati J, Dangi SS, Bag S, Maurya VP, Singh G, Kumar P, *et al.* Expression dynamics of HSP90 and nitric oxide synthase (NOS) isoforms during heat stress acclimation in Tharparkar cattle. *International Journal of Biometeorology*. 2017;61:1461-1469.
6. Chandra G, Aggarwal A. Effect of DL-Tocopherol Acetate on calving induced oxidative stress in periparturient crossbred cows during summer and winter seasons. *Indian Journal of Animal Research*. 2009;26(3):204-210.
7. Chauhan SS, Dunshea FR, Plozza TE, Hopkins DL, Ponnampalam EN. The impact of antioxidant supplementation and heat stress on carcass characteristics, muscle nutritional profile and functionality of lamb meat. *Animals*. 2020;10(8):1286.
8. Chen X, Shu H, Sun F, Yao J, Gu X. Impact of heat stress on blood, production, and physiological indicators in heat-tolerant and heat-sensitive dairy cows. *Animals*. 2023;13(16):2562.
9. Chouraddi R, Sreeja SJ, Johnson A, Naveen P, Gujjalkar P, Singh AK, *et al.* A review on the effects of micronutrients in heat stress alleviation in dairy animals. *Current Applied Science and Technology*. 2022;41(36):59-68.
10. Chrast V, Langova L, Novotna I, Zemanova M, Vrtkova I, Urban T, *et al.* Effect of temperature-humidity index on physiological and haematological indicators in dairy cows. *Journal of Central European Agriculture*. 2023;24(4):802-808.
11. Collier RJ, Stiening CM, Pollard BC, VanBaale MJ, Baumgard LH, Gentry PC, *et al.* Use of gene expression microarrays for evaluating environmental stress tolerance at the cellular level in cattle. *Journal of Animal Science*. 2006;84(13):E1-E13.
12. Dangi SS, Gupta M, Maurya D, Yadav VP, Panda RP, Singh G, *et al.* Expression profile of HSP genes during different seasons in goats (*Capra hircus*). *Tropical Animal Health and Production*. 2012;44:1905-1912.
13. Dangi SS, Gupta M, Nagar V, Yadav VP, Dangi SK, Shankar OM, *et al.* Impact of short-term heat stress on physiological responses and expression profile of HSPs in Barbari goats. *International Journal of Biometeorology*. 2014;58:2085-2093.
14. Degen AA, Young BA. Effect of air temperature and energy intake on body mass, body composition and energy requirements in sheep. *Journal of Agricultural Science*. 2002;138(2):221-226.
15. Djelailia H, M'hamdi N, Bouraoui R, Najjar T. Effects of thermal stress on physiological state and hormone

- concentrations in Holstein cows under arid climatic conditions. *South African Journal of Animal Science*. 2021;51(4):452-459.
16. Garcia AB, Angeli N, Machado L, de Cardoso FC, Gonzalez F. Relationships between heat stress and metabolic and milk parameters in dairy cows in southern Brazil. *Tropical Animal Health and Production*. 2015;47:889-894.
 17. Ghorbani A, Sadri H, Alizadeh AR, Bruckmaier RM. Performance and metabolic responses of Holstein calves to supplemental chromium in colostrum and milk. *Journal of Dairy Science*. 2012;95(10):5760-5769.
 18. Hooda OK, Upadhyay RC. Growth rate, hormonal and physiological responses of kids subjected to thermal and exercise stress. *Journal of Environmental Research*. 2015;9(4):1095-1101.
 19. Kargar S, Mousavi F, Karimi-Dehkordi S. Effects of chromium supplementation on weight gain, feeding behaviour, health and metabolic criteria of environmentally heat-loaded Holstein dairy calves from birth to weaning. *Archives of Animal Nutrition*. 2018;72(6):443-457.
 20. Kishore A, Sodhi M, Kumari P, Mohanty AK, Sadana DK, Kapila N, *et al.* Peripheral blood mononuclear cells: a potential cellular system to understand differential heat shock response across native cattle (*Bos indicus*), exotic cattle (*Bos taurus*), and riverine buffaloes (*Bubalus bubalis*) of India. *Cell Stress and Chaperones*. 2014;19(5):613-621.
 21. Lashkari S, Habibi M, Jensen SK. A review on the role of chromium supplementation in ruminant nutrition effects on productive performance, blood metabolites, antioxidant status, and immunocompetence. *Biological Trace Element Research*. 2018;186:305-321.
 22. Maurya VP, Sejian V, Gupta M, Dangi SS, Kushwaha A, Singh G, *et al.* Adaptive mechanisms of livestock to changing climate. *Climate Change Impact on Livestock: Adaptation and Mitigation*. Springer, New Delhi; 2015. p. 123-138.
 23. Maurya VP, Sejian V, Kumar D, Naqvi SMK. Effect of induced body condition score differences on sexual behavior, scrotal measurements, semen attributes and endocrine responses in Malpura rams under hot semi-arid environment. *Journal of Animal Physiology and Animal Nutrition*. 2010;94(6):308-317.
 24. Maurya VP, Sejian V, Kumar D, Naqvi SMK. Biological ability of Malpura rams to counter heat stress challenges and its consequences on production performance in a semi-arid tropical environment. *Biological Rhythm Research*. 2018;49(3):479-493.
 25. Maurya VP, Sejian V, Kumar D, Naqvi SMK. Impact of heat stress, nutritional stress and their combinations on the adaptive capability of Malpura sheep under hot semi-arid tropical environment. *J Anim Behav Biometeorol*. 2020;7(1):17-23.
 26. Mishra SR, Palai TK. Importance of heat shock protein 70 in livestock at cellular level. *J Mol Pathophysiol*. 2014;3(2):1-5.
 27. Mohr E, Langbein J, Nürnberg G. Heart rate variability: a noninvasive approach to measure stress in calves and cows. *Physiol Behav*. 2002;75(1-2):251-259.
 28. Mousavi F, Karimi-Dehkordi S, Kargar S, Ghaffari MH. Effect of chromium supplementation on growth performance, meal pattern, metabolic and antioxidant status and insulin sensitivity of summer-exposed weaned dairy calves. *Anim*. 2019;13(5):968-974.
 29. Mudgal V, Garg AK, Dass RS, Varshney VP. Effect of selenium and copper supplementation on blood metabolic profile in male buffalo (*Bubalus bubalis*) calves. *Biol Trace Elem Res*. 2008;121(1):31-38.
 30. Parashuramulu S, Nagalakshmi D, Rao DS, Kumar MK, Swain PS. Effect of zinc supplementation on antioxidant status and immune response in buffalo calves. *Anim Nutr Feed Technol*. 2015;15(2):179-188.
 31. Pearl LH, Prodromou C, Workman P. The Hsp90 molecular chaperone: an open and shut case for treatment. *J Biochem*. 2008;410(3):439-453.
 32. Raghunandan T, Sultana JR, Chandra AS, Prakash MG, Venkateswarlu M, Ramana DBV. Effect of dietary chromium, vitamin E and selenium supplementation on biochemical and physiological parameters of Holstein Friesian cows under heat stress. *Indian J Anim Sci*. 2022;92(7):858-864.
 33. Ranjitkar S, Bu D, Van Wijk M, Ma Y, Ma L, Zhao L, *et al.* Will heat stress take its toll on milk production in China? *Clim Change*. 2020;161(4):637-652.
 34. Rhoads ML, Rhoads RP, VanBaale MJ, Collier RJ, Sanders SR, Weber WJ, *et al.* Effects of heat stress and plane of nutrition on lactating Holstein cows: I. Production, metabolism, and aspects of circulating somatotropin. *J Dairy Sci*. 2009;92(5):1986-1997.
 35. Saeed OA, Jaber BT, Mohammed MTA, Sani UM, Ziara KS, Saad HM. Impacts of heat stress on blood metabolic in different periods of lactation and pregnancy in Holstein cows. In: *IOP Conference Series: Environ Earth Sci*. 2021;779(1):012013.
 36. Sejian V, Maurya VP, Naqvi SM. Effect of walking stress on growth, physiological adaptability and endocrine responses in Malpura ewes in a semi-arid tropical environment. *Int J Biometeorol*. 2012;56(2):243-252.
 37. Shaji S, Sejian V, Bagath M, Manjunathareddy GB, Kurien EK, Varma G, *et al.* Summer season-related heat and nutritional stresses on the adaptive capability of goats based on blood biochemical response and hepatic HSP70 gene expression. *Biol Rhythm Res*. 2017;48(1):65-83.
 38. Shan Q, Ma FT, Jin YH, Gao D, Li HY, Sun P. Chromium yeast alleviates heat stress by improving antioxidant and immune function in Holstein mid-lactation dairy cows. *Anim Feed Sci Technol*. 2020;269:114635.
 39. Shinde PL, Dass RS, Garg AK. Effect of vitamin E and selenium supplementation on haematology, blood chemistry and thyroid hormones in male buffalo (*Bubalus bubalis*) calves. *J Anim Feed Sci*. 2009;18(2):241-256.
 40. Son AR, Kim SH, Islam M, Oh SJ, Paik MJ, Lee SS, *et al.* Higher concentration of dietary selenium, zinc, and copper complex reduces heat stress-associated oxidative stress and metabolic alteration in the blood of Holstein and Jersey steers. *Animals*. 2022;12(22):3104.
 41. Srikandakumar A, Johnson EH. Effect of heat stress on milk production, rectal temperature, respiratory rate and blood chemistry in Holstein, Jersey and Australian Milking Zebu cows. *Trop Anim Health Prod*. 2004;36(7):685-692.

42. Tejaswi V, Balamurugan B, Samad HA, Sarkar M, Maurya VP, Singh G. Differential endocrine and antioxidant responses to heat stress among native and crossbred cattle. *J Vet Behav.* 2020;39:1-5.
43. Yadav B, Singh G, Wankar A, Dutta N, Chaturvedi VB, Verma MR. Effect of simulated heat stress on digestibility, methane emission and metabolic adaptability in crossbred cattle. *Asian J Anim Sci.* 2016;29(11):1585-1593.
44. Yari M, Nikkhah A, Alikhani M, Khorvash M, Rahmani H, Ghorbani GR. Physiological calf responses to increased chromium supply in summer. *J Dairy Sci.* 2010;93(9):4111-4120.
45. Zheng Y, Xie T, Li S, Wang W, Wang Y, Cao Z, *et al.* Effects of selenium as a dietary source on performance, inflammation, cell damage, and reproduction of livestock induced by heat stress: A review. *Front Immunol.* 2022;12:820853.
46. Sun B, Xun W, Wei X, Gao Y, Wang Y, Chen H. Effects of dietary selenium supplementation on antioxidant capacity and immune characteristics in late lactation dairy cows. *J Dairy Res.* 2018;85(4):400-403.
47. Kumbhar S, Khan RU, Parveen F, Alhidary IA, Alhotan RA. Effect of heat stress and the role of exogenous zinc and selenium on oxidative stress, lipid peroxidation, and apoptosis in rats. *J Anim Physiol Anim Nutr.* 2018;102(5):1134-1140.