

International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
IJABR 2025; SP-9(1): 653-655
www.biochemjournal.com
Received: 05-10-2024
Accepted: 09-11-2024

Dr. Parthkumar Nanda
M.V.Sc Scholar, Department of
Veterinary Medicine, Navsari
Veterinary College, Kamdhenu
University, Navsari, Gujarat,
India

Dr. Saurabh Parmar
Assistant Professor, Veterinary
Clinical Complex, College of
Veterinary Science and Animal
Husbandry, Kamdhenu
University, Navsari, Gujarat,
India

Dr. Nirav Patel
M.V.Sc Scholar, Department of
Veterinary Medicine, Navsari
Veterinary College, Kamdhenu
University, Navsari, Gujarat,
India

Dr. Jignesh Vala
Assistant Professor, Veterinary
Clinical Complex, College of
Veterinary Science and Animal
Husbandry, Kamdhenu
University, Navsari, Gujarat,
India

Dr. Manish Patel
Associate Professor & Head,
Department of Veterinary
Medicine, Navsari Veterinary
College, Kamdhenu University,
Navsari, Gujarat, India

Dr. Hitesh Parmar
Assistant Professor, Department
of Veterinary Clinical Complex,
College of Veterinary Science and
Animal Husbandry, Kamdhenu
University, Navsari, Gujarat,
India

Dr. Sudhir Mehta
Assistant Professor, Department
of Veterinary Medicine, Navsari
Veterinary College, Kamdhenu
University, Navsari, Gujarat,
India

Corresponding Author:
Dr. Parthkumar Nanda
M.V.Sc Scholar, Department of
Veterinary Medicine, Navsari
Veterinary College, Kamdhenu
University, Navsari, Gujarat,
India

Hematobiochemical profiles and cardiac biomarker assessment in Labrador retriever dogs

Parthkumar Nanda, Saurabh Parmar, Nirav Patel, Jignesh Vala, Manish Patel, Hitesh Parmar and Sudhir Mehta

DOI: <https://doi.org/10.33545/26174693.2025.v9.i1Si.3582>

Abstract

This study aimed to establish baseline hematobiochemical and cardiac biomarker parameters in clinically healthy Labrador Retrievers and assess the influence of body weight on these indices. Sixteen adult Labrador Retrievers (2–6 years, both sexes) were categorized into two groups based on body weight: Group-I (<30 kg) and Group-II (>30 kg). Comprehensive evaluations including hematological, biochemical, and cardiac biomarker analyses were performed following confirmation of the dogs' health status. Hematological parameters including hemoglobin, packed cell volume, RBC, WBC and platelet counts, as well as biochemical parameters such as AST, ALT, total protein, albumin, BUN and creatinine were within normal ranges with no significant intergroup differences. Cardiac biomarker analysis revealed cTnI levels of 0.0056 ± 0.0014 ng/mL in Group-I and 0.0081 ± 0.0018 ng/mL in Group-II, while NT-proBNP levels were 1.9731 ± 0.3085 pmol/L and 3.0789 ± 0.8485 pmol/L, respectively. These differences were not statistically significant indicating stable cardiac health across varying body weights. This study provides valuable baseline data on hematobiochemical and cardiac biomarker parameters in Labrador Retrievers emphasizing the minimal impact of body weight on these indices and contributing to improved cardiac health assessment in this breed.

Keywords: Labrador retrievers, hematobiochemical parameters, cardiac biomarkers, body weight

Introduction

The Labrador Retriever, renowned for its friendly temperament, intelligence, and versatility has become one of the most popular dog breeds globally. This increased popularity has brought heightened attention to the breed's specific health requirements particularly concerning cardiovascular health. Cardiovascular diseases are a growing concern in veterinary practice mirroring their significance in human medicine. Early detection and accurate assessment of cardiac function are essential for managing such conditions in companion animals. Haematological evaluations play a pivotal role in assessing the physiological and pathological states of animals. Complete blood cell counts (CBC) are critical diagnostic tools providing essential information that facilitates accurate diagnosis or the creation of differential diagnosis lists (Dixon, 1997; DeNicola, 2011) [14, 15]. Establishing breed-specific reference ranges for haematological parameters is crucial as these values are influenced by factors such as lifestyle, gender, age and environmental conditions (Olayemi and Ighagbon, 2011; Connolly *et al.*, 2020) [18, 17]. In veterinary cardiology, the advent of cardiac biomarkers such as cardiac troponins (cTn) and brain natriuretic peptides (BNP) has revolutionized diagnostic capabilities. These biomarkers provide non-invasive insights into cardiac stress and injury offering a reliable indication of myocardial damage and disease severity (Noszczyk and Nowak, 2011) [1]. Combined with advanced diagnostic methods like echocardiography, these tools enable a comprehensive evaluation of cardiac function. This study aims to assess haematological and cardiac parameters including biomarkers in healthy Labrador Retrievers. By establishing reference ranges and examining the influence of body weight and other factors this research contributes to enhancing diagnostic precision and advancing Veterinary cardiac care.

Materials and Methods

Animals and Inclusion Criteria

This study included clinically healthy adult Labrador Retriever dogs, aged 2–6 years, with no history of clinical conditions such as coughing, breathlessness or exercise intolerance. All included dogs had a history of routine preventive healthcare including vaccinations and deworming. The age, sex and body weight of each dog were recorded prior to the study.

Clinico-physiological Examination

A comprehensive clinical examination was conducted on each dog to assess various physiological parameters including heart rate, pulse rate, rectal temperature, respiratory rate, breathing pattern, capillary refill time (CRT) and the color of visible mucous membranes.

Haemato-biochemical Examination

Blood Collection

Approximately 5 mL of blood was collected aseptically and of this, 1 mL was transferred into a K₃EDTA vacutainer for haematological analysis and the remaining 4 mL was transferred into a clot activator vacutainer for serum separation. The serum samples were stored at -20 °C until further biochemical analysis.

Haematological and Biochemical analysis

Haematological parameters *viz.* Haemoglobin (Hb, g/dL), Total Erythrocyte Count (TEC, $\times 10^6/\mu\text{L}$), Total Leukocyte Count (TLC, $\times 10^3/\mu\text{L}$), Packed Cell Volume (PCV, %), Platelet Count ($\times 10^5/\mu\text{L}$), Differential Leukocyte Count (DLC, %) were measured using a fully automatic blood cell counter (MEK-6420P, Nihon Kohden India).

Biochemical parameters *viz.* Alanine Aminotransferase (ALT, IU/L), Aspartate Aminotransferase (AST, IU/L), Serum Creatinine (mg/dL), Blood Urea Nitrogen (BUN, mg/dL), Total Protein (TP, g/dL), Albumin (g/dL) were assessed using a semi-automated biochemical analyzer (Microlab 300) with commercially available diagnostic kits.

Cardiac Biomarkers

The cardiac biomarkers N-terminal pro-brain natriuretic peptide (NT-proBNP) and cardiac Troponin I (cTnI) were quantitatively estimated using FINETEST Canine NT-proBNP and FINETEST Canine hs-cTnI ELISA kits. The analysis was performed using an ELISA reader (I Gene Labserve).

Equipment Used

- Fully automatic blood cell counter (MEK-6420P, Nihon Kohden India)
- Semi-automated biochemical analyzer (Microlab 300)
- ECG machine (BPL CARDIART 6208)
- ELISA reader (I Gene Labserve)

Results and Discussion

Haematological Analysis

All assessed parameters, including haemoglobin (Hb), packed cell volume (PCV), total erythrocyte count (TEC), total leukocyte count (TLC), platelet count and differential leukocyte counts (neutrophils, lymphocytes, monocytes, and eosinophils) were within normal physiological ranges without any statistically significant differences between Group I (<30 kg) and Group II (>30 kg) (Table 1) suggesting that body weight does not significantly influence these hematological parameters in healthy Labrador Retriever dogs.

Table 1: Mean $\pm$ S.E. of various haematological parameters in Labrador Retriever dogs

Sr. No.	Parameters	Group-I (n=8)	Group-II (n=8)	p values
1	Hb (g/dl)	14.675 $\pm$ 0.83	15.463 $\pm$ 0.39	0.41
2	PCV (%)	47.53 $\pm$ 2.19	49.675 $\pm$ 1.74	0.46
3	TEC (m/cmm)	7.6900 $\pm$ 0.27	7.7038 $\pm$ 0.31	0.93
4	TLC (cmm)	11612.50 $\pm$ 736.65	12425.00 $\pm$ 1445.78	0.62
5	Neutrophils (%)	67.88 $\pm$ 3.01	64.38 $\pm$ 2.24	0.37
6	Lymphocytes (%)	25.88 $\pm$ 2.79	29.63 $\pm$ 2.42	0.33
7	Eosinophils (%)	3.00 $\pm$ 0.57	3.00 $\pm$ 0.59	1.00
8	Monocyte (%)	3.25 $\pm$ 0.42	3.00 $\pm$ 0.78	0.78
9	Platelets ($\times 10^5/\mu\text{L}$)	2.5975 $\pm$ 0.15	2.6325 $\pm$ 0.19	0.88

These findings align with previous studies, including those by Haritha *et al.* (2017) [2] and Ramteke *et al.* (2019) [2], which reported no significant differences in hematological parameters in healthy dogs between different breeds. Kara *et al.* (2024) [4] also found that gender differences did not significantly influence most haematological parameters. The normal haematological values in this study support the absence of anaemia, infection, or haematological abnormalities indicating that these factors are unlikely to affect cardiac function in the studied dogs.

Biochemical Analysis

The biochemical parameters assessed including AST, ALT, total protein, albumin, blood urea nitrogen (BUN) and creatinine showed no significant differences between the two groups (Table 2). All values were within the normal ranges reported by Kahn (2010) [5] and Kaneko *et al.* (2008) [6].

Table 2: Mean $\pm$ S.E. of various biochemical parameters in Labrador Retriever dogs

Sr. No.	Parameters	Group-I (n=8)	Group-II (n=8)	p values
1	AST(IU/L)	24.53 $\pm$ 1.36	21.18 $\pm$ 2.00	0.19
2	ALT (IU/L)	19.92 $\pm$ 0.41	19.96 $\pm$ 1.09	0.98
3	Total protein (g/dl)	8.33 $\pm$ 0.15	8.0138 $\pm$ 0.18	0.19
4	Albumin (g/dl)	3.36 $\pm$ 0.08	3.39 $\pm$ 0.68	0.77
5	BUN (mg/dl)	12.79 $\pm$ 0.34	12.23 $\pm$ 0.43	0.33
6	Creatinine (mg/dl)	0.98 $\pm$ 0.08	1.18 $\pm$ 0.07	0.08

These findings are consistent with studies by Haritha *et al.* (2017) [2] and Amaravathi *et al.* (2019) [7], which reported normal ALT, BUN, creatinine and total protein levels in healthy dogs and those with early cardiac diseases. In contrast, some studies, such as Martin *et al.* (2009) [8], have noted biochemical alterations including azotemia and elevated liver enzymes in dogs with advanced cardiac conditions.

The absence of significant differences in biochemical parameters in this study highlights the limited impact of body weight on routine biochemical markers in healthy

Labrador Retrievers.

Cardiac Biomarkers

Table 3: Mean $\pm$ S.E. of cardiac biomarkers in Labrador Retriever dogs

Sr. No.	Parameters	Group-I (n=8)	Group-II(n=8)	p values
1	Cardiac troponin-I (ng/ml)	0.0056 $\pm$ 0.0014	0.0081 $\pm$ 0.0018	0.28
2	NT-proBNP (pmol/L)	1.9731 $\pm$ 0.3085	3.0789 $\pm$ 0.8485	0.75

The levels of cardiac troponin I (cTnI) and NT-proBNP, shown in Table 3 were within the normal range for all dogs with no significant differences between the groups. The mean cTnI levels in Group I (0.0056 $\pm$ 0.0014 ng/mL) and Group II (0.0081 $\pm$ 0.0018 ng/mL) were well below the diagnostic threshold of 0.5 ng/mL suggested by Pelander *et al.* (2002) [9]. Similarly, NT-proBNP levels in Group I (1.9731 $\pm$ 0.3085 pmol/L) and Group II (3.0789 $\pm$ 0.8485 pmol/L) were comparable to baseline levels reported by Ettinger (2012) [10].

Sleeper *et al.* (2001) [11] also documented that cTnI levels in healthy dogs are typically 0.0–0.04 ng/mL, similar to the findings in this study. Elevated levels of cTnI and NT-proBNP have been associated with cardiac stress and myocardial injury in diseased animals (Barathan *et al.*, 2018) [12]. However, the absence of significant differences and the low biomarker levels in this study indicate that all dogs were free from cardiac stress or injury. These findings can also serve as baseline data for Veterinary cardiology research in India.

Conflict of Interest

None declared.

Acknowledgments

The authors wish to thank the staff of the Veterinary Clinical Complex and the Department of Veterinary Microbiology, Veterinary College, Kamdhenu University, Navsari, Gujarat, for their valuable assistance during the study.

Reference

- Noszczyk-Nowak A. NT-pro-BNP and troponin I as predictors of mortality in dogs with heart failure. *Polish Journal of Veterinary Sciences*. 2011;14(4).
- Haritha GS, Kumar KS, Ayodhya S, Kumar VVV. Prevalence of cardiac disorders in canines—a clinical study. *Intas Polivet*. 2017;18(1):148-152. Available from: <https://www.indianjournals.com/ijor.aspx?target=ijor:ip o&volume=18&issue=1&article=048&type=pdf>
- Ramteke SY, Deshmukh SG, Pawshe CH, Kuralkar SV, Thorat MG, Kudalkar SD. Comparative study of different haematological and biochemical parameters in normal German Shepherd and Labrador dogs. *Journal of Entomology and Zoology Studies*. 2019;7(5):972-975.
- Kara H, Şenel Y, Sayım AA, Güven M. Effects of gender on hematologic parameters in Kangal Shepherd Dogs. *Research and Practice in Veterinary and Animal Science (REPVAS)*. 2024;1(1):36-42. Available from: <https://app.repvas.com/uploads/makale-pdf/effects-of-gender-on-hematologic-parameters-in-kangal-shepherd-dogs-c3H37j.pdf>
- Kahn CM, Line S, Aiello SE. Reference guides. *The Merck Veterinary Manual*. 2010;1(9th ed.).
- Kaneko JJ. *Clinical Biochemistry of Domestic Animals*. Vol. 273. Academic Press; 2008.
- Amaravathi P, Prebavathy T, Rajesh K, Rao VV, Bharathi S, Raghunath M. Clinicopathological findings in dogs with cardiac diseases. *Health*. 2019;58(1):1-20. doi:10.22271/j.ento.2020.v8.i5t.7700
- Martin MWS, Stafford Johnson MJ, Celona B. Canine dilated cardiomyopathy: a retrospective study of signalment, presentation and clinical findings in 369 cases. *Journal of Small Animal Practice*. 2009;50(1):23-29. doi:10.1111/j.1748-5827.2008.00659.x
- Pelander L, Häggström J, Jones B. Troponin I: a possible marker for myocardial cell injury in dogs. *Swedish Veterinary Journal*. 2001;53(10):497–502. Available from: <https://www.cabidigitallibrary.org/doi/full/10.5555/20013118124>
- Ettinger SJ, Suter PF. Acquired valvular and endocardial heart disease. In: *Canine Cardiology*. Philadelphia, PA: WB Saunders; 1970. p. 324-325.
- Sleeper MM, Clifford CA, Laster LL. Cardiac troponin I in the normal dog and cat. *Journal of Veterinary Internal Medicine*. 2001;15(5):501-503. doi:10.1111/j.1939-1676.2001.tb01582.x
- Barathan P, Sumathi D, Nambi AP. Cardiac troponin—a cardiac biomarker for point of care testing in acute heart failure syndrome in canines. *Intas Polivet*. 2018;19(2):315-318. Available from: <https://www.indianjournals.com/ijor.aspx?target=ijor:ip o&volume=19&issue=2&article=041&type=pdf>
- Barger AM. The complete blood cell count: a powerful diagnostic tool. *Veterinary Clinics: Small Animal Practice*. 2003;33(6):1207-1222.
- Dixon LR. The complete blood count: physiologic basis and clinical usage. *The Journal of Perinatal & Neonatal Nursing*. 1997;11(3):1-18.
- DeNicola DB. Advances in hematology analyzers. *Topics in Companion Animal Medicine*. 2011;26(2):52-61.
- Etim NN, Williams ME, Akpabio U, Offiong EE. Haematological parameters and factors affecting their values. *Agricultural Science*. 2014;2(1):37-47.
- Connolly SL, Nelson S, Jones T, Kahn J, Constable PD. The effect of age and sex on selected hematologic and serum biochemical analytes in 4,804 elite endurance-trained sled dogs participating in the Iditarod Trail Sled Dog Race pre-race examination program. *PLoS One*. 2020;15(8):e0237706.
- Olayemi FO, Ighagbon FO. Haematology of the German Shepherd dog in a humid tropical environment. *Comparative Clinical Pathology*. 2011;20:61-64.