



ISSN Print: 2664-844X
 ISSN Online: 2664-8458
 IJAFS 2025; 7(1): 94-96
www.agriculturaljournals.com
 Received: 23-11-2024
 Accepted: 27-12-2024

Dr. Mansoor Nabi Mir
 Animal Husbandry,
 Department Kashmir, Jammu
 and Kashmir, India

The effect of excess green feeding on fat percentage and udder health in dairy cows

Mansoor Nabi Mir

DOI: <https://doi.org/10.33545/2664844X.2025.v7.i2b.284>

Abstract

Green fodder is a crucial dietary component for dairy cows, supplying essential nutrients that support milk production and overall health. However, excessive feeding of green forage can cause imbalances in nutrient intake, negatively affecting milk composition particularly fat percentage and increasing the risk of udder health issues, such as mastitis. This study investigates the impact of excess green feeding on milk fat percentage and udder health, with the aim of providing evidence-based recommendations for optimizing green fodder inclusion in dairy diets to enhance milk quality and maintain udder health.

Keywords: Excess green feeding, enhance milk quality and maintain udder health, dairy cows

1. Introduction

Green fodder is rich in fiber, moisture, and nutrients, making it an integral part of dairy cow diets (Elgersma, 2015) ^[5]. While moderate green fodder feeding supports optimal milk production, over-reliance on it can disrupt nutrient balance, leading to decreased milk fat content and compromised udder health. The high moisture content of excess green forage may influence rumen fermentation patterns and systemic metabolic functions, thereby affecting both milk composition and the immune response to udder pathogens (Negussie *et al.*, 2013) ^[8]. This study explores the dual impact of excess green feeding on milk fat percentage and udder health, aiming to provide practical guidelines for dairy farmers.

2. Milk Fat Percentage: Importance and Regulation

2.1 Role of Fat in Milk Quality

Milk fat significantly contributes to the nutritional and sensory qualities of dairy products, influencing market value and consumer acceptance. Its concentration is influenced by genetic, environmental, and dietary factors (Berry *et al.*, 2007) ^[2]. Dietary interventions, such as excessive green fodder feeding, can dilute the energy density of the diet, altering the volatile fatty acid (VFA) profile in the rumen, which in turn impacts milk fat synthesis.

2.2 Mechanisms Affecting Milk Fat Percentage

Excess green feeding impacts milk fat percentage through several mechanisms:

- **Reduced Fiber Intake:** High green fodder diets often displace fibrous dry matter, leading to lower rumen pH and decreased acetate production, a key precursor for milk fat synthesis.
- **Increased Moisture Content:** High-moisture diets reduce energy density, causing energy deficits that suppress milk fat output (Bostanova *et al.*, 2021) ^[3].

3. Udder Health and Excess Green Feeding

3.1 Impact on Mastitis Risk

Excessive green feeding can predispose cows to subacute ruminal acidosis (SARA), a metabolic disorder that impairs immune function and increases susceptibility to mastitis (Salama *et al.*, 2003) ^[9]. Altered rumen fermentation can lead to systemic inflammation, further compromising udder defenses.

Corresponding Author:
Dr. Mansoor Nabi Mir
 Animal Husbandry,
 Department Kashmir, Jammu
 and Kashmir, India

3.2 Moisture and Hygiene

High-moisture feces associated with green fodder diets can create unhygienic conditions, promoting bacterial growth on the udder and increasing mastitis risk.

4. Methodology

4.1 Study Design

A controlled study was conducted on 100 lactating Holstein Friesian cows over six months. The cows were divided into two groups: a control group fed a balanced diet with controlled green fodder levels and a test group fed a diet comprising 70% green fodder, exceeding recommended levels. Milk fat percentage was analyzed weekly using a milkPro Ultrasonic Analyser, and udder health was assessed bi-weekly through somatic cell count (SCC) and clinical mastitis evaluations. This study aimed to investigate the impact of a high green fodder diet on milk fat content and udder health in lactating cows.

5. Results and Discussion

5.1 Effect on Milk Fat Percentage

Cows in the test group showed a significant decline in milk fat percentage (3.2%) compared to the control group (3.8%). The reduction was attributed to lower acetate production and disrupted lipid metabolism, consistent with findings by Hashemzadeh-Cigari *et al.* (2014) ^[6] and Alyousef *et al.* (2022) ^[1].

5.2 Effect on Udder Health

- **Somatic Cell Count:** The test group had an average SCC of 450,000 cells/mL, higher than the control group's 250,000 cells/mL, indicating increased mastitis prevalence (Chandra *et al.*, 2015) ^[4].
- **Clinical Mastitis:** A 20% higher incidence of mastitis was observed in the test group, linked to poor hygiene due to wetter bedding and fecal contamination.




Fig 1: The effects of excess green feeding on dairy cows through four distinct graphs. Each graph highlights key metrics such as milk fat percentage, somatic cell count (SCC), mastitis incidence, and the diet composition effects

6. Practical Implications

6.1 Optimizing Green Fodder Feeding

A balanced approach that includes dry matter sources like hay and silage can mitigate the adverse effects of excessive green fodder on rumen health and milk fat synthesis.

6.2 Hygiene and Udder Care

Implementing strict hygiene protocols, such as regular cleaning of udders and bedding, is essential to prevent mastitis in cows fed high-moisture diets (McCarthy *et al.*, 2007) ^[7].

6.3 Monitoring and Intervention

Regular monitoring of milk composition and SCC allows for early detection of dietary imbalances and udder health issues, enabling timely management interventions.

7. Conclusion

Excessive green feeding negatively impacts both milk fat percentage and udder health in dairy cows, primarily through altered rumen fermentation patterns and increased mastitis susceptibility. A balanced diet, proper hygiene management, and regular health monitoring are critical for

maintaining high milk quality and ensuring udder health. These findings highlight the need for tailored feeding strategies to optimize dairy productivity and farm profitability.

8. References

1. Alyousef YM, *et al.* Influence of functional feed supplements on milk production efficiency and udder health. *Sustainability*. 2022;14(14):8444.
2. Berry DP, *et al.* Effect of strain of Holstein–Friesian and feed system on udder health and milking characteristics. *Livest Sci*. 2007;107(1):10-20.
3. Bostanova S, *et al.* Indicators of full value feeding rations for dairy cows. *Braz J Biol*. 2021.
4. Chandra G, *et al.* Effect of vitamin E and zinc supplementation on milk composition and udder health. *Indian J Anim Nutr Feed Technol*. 2015;15(1):75-80.
5. Elgersma A. Grazing increases the unsaturated fatty acid concentration of milk from grass-fed cows. *Eur J Lipid Sci Technol*. 2015;117(10):1555-69.
6. Cigari HF, *et al.* Effects of herbal supplementation on udder health in Holstein cows. *J Dairy Sci*. 2014;97(7):4110-9.
7. McCarthy S, *et al.* The impact of feed system on udder health and milk yield. *Livest Sci*. 2007;112(2-3):133-43.
8. Negussie E, *et al.* Genetic associations of fat-protein ratio with udder health in Nordic Red cattle. *J Dairy Sci*. 2013;96(12):7654-65.
9. Salama AAK, *et al.* Dietary zinc-methionine supplementation and udder health. *J Dairy Res*. 2003;70(1):9-17.