


International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
NAAS Rating: 5.29
IJABR 2025; 9(3): 341-347
www.biochemjournal.com
Received: 01-12-2024
Accepted: 06-01-2025

Vijayakumar Chinnathambi
Assistant Professor,
Department of Veterinary
Pharmacology and Toxicology,
Veterinary College and
Research Institute,
TANUVAS, Salem,
Tamil Nadu, India

Cholinergic modulation of contractile responses in caprine urinary bladder smooth muscle: Role of urothelium and atropine sensitivity

Vijayakumar Chinnathambi

DOI: <https://www.doi.org/10.33545/26174693.2025.v9.i3e.3955>

Abstract

The cholinergic system plays a pivotal role in urinary bladder function, primarily facilitating bladder emptying through the activation of muscarinic receptors in smooth muscle. This process can also be modulated by signaling molecules from the urothelium. In this study, we investigated the role of the urothelium in cholinergic-mediated bladder smooth muscle contraction.

Bladder smooth muscle contraction was induced using the muscarinic agonist acetylcholine (ACh). Our results demonstrated that ACh elicited a concentration-dependent contractile response in goat urinary bladder smooth muscle. Notably, the removal of the urothelium (detrusor muscle strips) resulted in a leftward shift of the ACh-induced contractile response curve, leading to a stronger and more pronounced contraction. The muscarinic response was significantly enhanced in the absence of the urothelium compared to intact bladder smooth muscle.

Furthermore, in the presence of atropine, a muscarinic receptor antagonist, ACh-mediated contractions were abolished. Minimal responses were observed in denuded caprine detrusor muscle strips, while no contraction occurred in detrusor muscle strips with an intact urothelium.

Our findings suggest that the caprine urinary bladder exhibits pharmacological properties similar to those of the human bladder, making it a potentially valuable model for evaluating new therapeutic agents for urinary incontinence.

Keywords: Acetylcholine, atropine, caprine detrusor muscle, cholinergic system

Introduction

Urinary incontinence refers to the involuntary leakage of urine, often resulting from an overactive bladder or other underlying conditions such as arthritis, prostatitis, weakened bladder or pelvic floor muscles, or nerve damage caused by diseases like multiple sclerosis, diabetes, or Parkinson's disease. It is more prevalent in older adults, particularly women. With advancements in medicine, there are now more treatment options and management strategies available than ever before. The urinary bladder's inner lining consists of a mucous membrane composed of transitional epithelium, which is continuous with that of the ureters. Beneath this lies the submucosa, a connective tissue layer containing elastic fibers that support the mucous membrane. The most prominent layer of the bladder wall is the muscularis, composed of smooth muscle fibers arranged in an interwoven pattern, collectively known as the detrusor muscle. Detrusor smooth muscle is made up of spindle-shaped cells containing thin actin and thick myosin myofilaments, which are interconnected both electrically and mechanically. Unlike skeletal and cardiac muscles, smooth muscle contraction follows a distinct mechanism. The regulation of bladder function involves multiple neurotransmitters, including acetylcholine, norepinephrine, dopamine, serotonin, excitatory and inhibitory amino acids, adenosine triphosphate, nitric oxide, and neuropeptides. These neurotransmitters play roles in both peripheral and central nervous system control of bladder activity (Andersson, 2013) [6].

Among the various neuronal inputs to the urinary bladder, parasympathetic innervation plays a crucial role in the micturition phase. Acetylcholine, acting on muscarinic M3 receptors, is responsible for the majority of contractile responses in the detrusor muscle across most species (Chess-Williams *et al.*, 2001; Matsui *et al.*, 2000) [13, 25]. However, the non-adrenergic, non-cholinergic (NANC) neurotransmitter ATP also contributes to

Corresponding Author:
Vijayakumar Chinnathambi
Assistant Professor,
Department of Veterinary
Pharmacology and Toxicology,
Veterinary College and
Research Institute,
TANUVAS, Salem,
Tamil Nadu, India

parasympathetically mediated contractions (Burnstock, 2008; D'Agostino *et al.*, 2012) ^[10, 15]. The extent to which the purinergic signaling system influences detrusor contraction, primarily via the P2X1 purinoceptor, varies among species (O'Reilly *et al.*, 2001; Vial and Evans, 2000) ^[29, 45]. In addition to its direct contractile role, ATP exerts paracrine functions, stimulating the release of other neuromodulators from the bladder urothelium (Winder *et al.*, 2014) ^[48]. These substances can modulate bladder function by acting directly on smooth muscle cells, suburothelial cells, afferent or efferent neurons, or by inducing the release of additional transmitters from the urothelium (Birder and Andersson, 2013) ^[6].

As previously mentioned, acetylcholine plays a crucial role in normal bladder function and can influence signaling at multiple levels of the micturition reflex arc (Kullmann *et al.*, 2008) ^[22]. Consequently, anticholinergic drugs are commonly used to treat overactive bladder (Robinson and Cardozo, 2012) ^[33]. However, their precise mechanism of action remains unclear. Studies suggest that anticholinergic drugs exert effects during both the voiding and storage phases of micturition. The involvement of the storage phase indicates a potential role for non-neuronal acetylcholine, such as that released from the urothelium or suburothelium (Andersson and Yoshida, 2003) ^[4]. This raises scientific interest in the interactions between the cholinergic transmitter system and other signaling pathways that may regulate non-neuronal acetylcholine release (Stenqvist *et al.*, 2020) ^[40].

The urothelium and lamina propria, which form the inner lining of the bladder and urethra, exhibit spontaneous phasic contractile activity and respond to neurokinin (Sadananda *et al.*, 2008) ^[34] and muscarinic receptor stimulation (Moro *et al.*, 2011) ^[28]. Although nitric oxide (NO) donors induce relaxation of this tissue, there is no evidence of endogenous NO release (Moro *et al.*, 2012) ^[27]. Crosstalk between the purinergic and cholinergic transmitter systems has been reported, wherein ATP appears to stimulate acetylcholine release, thereby supporting purinergic bladder contractions both *in vivo* and *in vitro* (Stenqvist *et al.*, 2017) ^[41] *in vitro* studies indicate that this atropine-sensitive component of ATP-induced contractions originates from the urothelium (Stenqvist *et al.*, 2017) ^[41]. This finding is further supported by transmitter release studies in both urothelial cells and urinary bladder tissue, suggesting the involvement of the P2Y6 purinoceptor in urothelial acetylcholine release (Silva *et al.*, 2015) ^[38]. However, the specific receptors functionally involved in this interaction remain unidentified, and potential neuronal involvement requires further investigation (Stenqvist *et al.*, 2020) ^[40].

To date, numerous pharmacological compounds have been identified, and a few have been marketed for human use following safety pharmacological studies conducted in laboratory animals. However, in certain safety pharmacology assessments, the use of slaughterhouse specimens instead of live laboratory animals could be highly beneficial, particularly for *in vitro* experiments such as isolated organ bath studies. This approach not only reduces the reliance on laboratory animals but also offers a viable alternative for specific experimental models. In this context, we have previously established the urinary bladder as a prototype smooth muscle model for ruminants, and the role of potassium channel openers in bladder hyperactivity has been well documented (Vijayakumar *et al.*, 2007) ^[46]. The

present study was conducted on the goat urinary bladder with the following objectives:

- To investigate the role of the mucous membrane in bladder function and
- To confirm that acetylcholine (ACh)-mediated contractile responses occur via muscarinic receptor activation.

Materials and Methods

Tissue mounting

The entire urinary bladder was collected immediately after slaughter from Black Bengal goats (2.0-2.5 years old) of either sex, weighing between 12.5 and 15.0 kg. The bladder was immediately flushed with Tyrode's solution and transported to the laboratory in chilled Tyrode's solution with the following composition (mM): NaCl, 120; KCl, 5.9; MgSO₄·7H₂O, 1.2; NaH₂PO₄·2H₂O, 1.2; CaCl₂·2H₂O, 2.5; and Glucose, 11.5. Each urinary bladder was weighed and dissected longitudinally along the midline of the ventral surface. Two transverse strips, approximately 8 mm in size, were obtained from the mid-region of the bladder. Additionally, the tissue was cut at the mid-dorsal section, yielding four strips of similar length for further experimentation.

Role of the Urothelium in Cholinergic-Mediated Contractile Responses: Non-Denuded Detrusor Muscle

For isometric contraction studies, urinary bladder smooth muscle with an intact mucous membrane (non-denuded detrusor muscle) was used. A tissue strip measuring approximately 2.5 cm × 8 mm was suspended in an organ bath containing 20 ml of Tyrode's solution, continuously bubbled with oxygen (pH 7.4) and maintained at 37±0.5 °C under a resting tension of 2 g. The tissue was equilibrated for 90 minutes, with washes performed every 15 minutes. Isometric contractions were recorded using a kymograph and fixated with a fixative fluid. To evaluate the cholinergic response, cumulative concentration-related contractions were induced by acetylcholine (ACh) in the range of 10⁻⁹ to 10⁻⁴ M, with increments of one log unit. After obtaining the control concentration-response curve, the tissue was allowed to fully relax. Subsequently, the urothelium (mucous membrane) was removed, and the tissue was remounted in the organ bath. Following an equilibration period, the denuded detrusor muscle was exposed to the same ACh concentrations (10⁻⁹ to 10⁻⁴ M), and the resulting cumulative contractile responses were recorded.

Antagonistic Effect of Atropine on Cholinergic-Mediated Contractile Responses in Denuded and Non-Denuded Detrusor Muscle

To evaluate the antagonistic effect of atropine (10⁻⁶ M) on cholinergic-mediated contractions in detrusor muscle with and without urothelium, cumulative concentration-related contractions were induced by acetylcholine (ACh) in the range of 10⁻⁹ to 10⁻⁴ M, with increments of one log unit. After obtaining the control concentration-response curve, tissues were treated with atropine (10⁻⁶ M) before eliciting a second dose-response curve, following complete relaxation of the initial response. In all experiments, tissues were pre-incubated with the cholinergic receptor blocker for 5 minutes prior to the addition of the agonist. Results were expressed as a percentage of the maximal response curve. The inhibitory effect of atropine on ACh-induced contractile

responses was assessed by comparing contractile responses before and after atropine treatment.

Drugs and Chemicals

Acetylcholine chloride (Sigma) and Atropine (Sigma) were used. The drug solutions used were either prepared fresh or were taken from the stock solution (10^{-1} M) stored at 4° C in a refrigerator.

Statistics


Results are presented as the mean $\pm$ standard error of the mean (SEM) or as means with 95% confidence limits. Statistical significance was determined using Student's t-test, with $p < 0.05$ considered statistically significant. The EC_{50} and IC_{50} values, along with their 95% confidence limits, were calculated using regression analysis based on the least squares method. Data analysis was performed using interactive non-linear regression with GraphPad Prism (GraphPad Prism Software, San Diego, CA, USA).

Results

Role of the urothelium on ACh induced concentration-related contractile response

Acetylcholine (ACh) administered cumulatively at concentrations ranging from 10^{-9} M to 10^{-4} M induced a concentration-dependent contractile response in caprine non-denuded detrusor muscle. The threshold concentration required to initiate contraction and the concentration eliciting the maximal response (E_{max}) were 10^{-6} M and 10^{-4} M, respectively. Following removal of the mucous membrane, isometric contraction studies revealed a 100-fold increase in the contractile response to ACh in denuded tissue compared to non-denuded tissue (Trace 1). In denuded detrusor muscle, the threshold concentration and E_{max} were found to be 10^{-9} M and 10^{-4} M, respectively. In the presence of the urothelium, ACh-mediated contractile responses at the highest concentration (10^{-4} M) reached an E_{Bmax} of $52.7 \pm 3.7\%$ ($n=6$), whereas the denuded detrusor muscle exhibited a maximal contractile response set at 100% (Table 1). The presence of the urothelium caused a

significant ($p < 0.001$) leftward shift in the ACh-induced concentration-response curve (Figure 1), with an increase in the mean $LogEC_{50}$ value (4.92 M) compared to the denuded detrusor response (6.06 M). All mean values were compared with 95% confidence limits.



Trace 1: ACh induced concentration-related contractile response Non Denuded (left) and Denuded Detrusor muscle (right).




Fig 1: ACh induced concentration-related contractile response Non Denuded and Denuded Detrusor muscle.

Table 1: ACh-induced dose-related contractile response in Non Denuded Detrusor Muscle vs Denuded Detrusor Muscle.


Exp	Response in percent	
	Non Denuded Detrusor Muscle	Denuded Detrusor Muscle
Ach 10^{-9} M	0	4.0 ± 0.9
Ach 10^{-8} M	0	15.7 ± 1.8
Ach 10^{-7} M	0	32.3 ± 6.9
Ach 10^{-6} M	11.8 ± 2.9	54.2 ± 7.2
Ach 10^{-5} M	25.3 ± 3.4	78.2 ± 4.2
Ach 10^{-4} M	52.5 ± 3.7	100

(Non Denuded Detrusor Muscle = 7 Nos.; Denuded Detrusor Muscle = 6 Nos.)

Antagonist effect of Atropine on ACh induced concentration-related contractile responses

Acetylcholine (ACh) administered cumulatively at concentrations ranging from 10^{-9} M to 10^{-4} M induced a concentration-dependent contractile response in goat detrusor muscle. However, in the presence of atropine (10^{-6} M), ACh-induced contractile responses were significantly inhibited (Trace 2). Atropine (10^{-6} M) nearly abolished the contractile response, with an E_{Bmax} of $12.8 \pm 3.9\%$ ($n = 6$) (Table 2). Atropine caused a significant ($p < 0.001$) leftward shift in the ACh-induced concentration-response curve, increasing the mean EC_{90} value to 2.6×10^{-4} M compared to

the control response (2.5×10^{-6} M). All mean values were compared with 95% confidence limits. Furthermore, in the presence of the urothelium, when tissue strips were pre-incubated with atropine (10^{-6} M), ACh-mediated contractile responses were completely abolished, and the E_{Bmax} value could not be determined. To confirm the cholinergic-mediated contractile response, the strips were thoroughly washed and subjected to isometric contraction studies. The results demonstrated a 100-fold increase in the mean EC_{50} value of ACh in denuded tissue compared to non-denuded tissue.



Trace 2: Impact of atropine (10^{-6} M) on ACh induced concentration-related contractile response in Denuded Detrusor muscle.

Table 2: Antagonist effect of atropine (10^{-6} M) on ACh-induced dose-related contractile response in goat urinary bladder.

Exp	Response in percent	
	Control	Atropine 10^{-6} M
Ach 10^{-9} M	2.9±1.5	0±0
Ach 10^{-8} M	6.5±2.4	0±0
Ach 10^{-7} M	15.4±3.0	1.3±0.8
Ach 10^{-6} M	41.5±3.0	2.8±1.1
Ach 10^{-5} M	73.7±1.7	3.7±2.0
Ach 10^{-4} M	100	12.8±3.1

(Control = 20 Nos.; Atropine = 6 Nos.)

Discussion

Bladder urothelial cells have been demonstrated to function as "sensor-transducers," capable of both releasing and responding to various signaling molecules (Birder and Andersson, 2013) [6]. These cells actively detect chemical, mechanical, and thermal stimuli, generating appropriate cellular responses (Chai *et al.*, 2016) [42]. Urothelial cells release a range of signaling molecules, including ATP, acetylcholine, substance P, nerve growth factor, nitric oxide, antiproliferative factor, and cytokines (Sui *et al.*, 2014; de Groat, 2013) [7, 16]. Additionally, urothelial cells express multiple classes of transducer receptors, including cholinergic (nicotinic and muscarinic), adrenergic (α and β), transient receptor potential channels (TRPV1, TRPV2, TRPV4, TRPM8), purinergic (P2X, P2Y), adenosine (A2a, A2b, and A3), and estrogen (ER α , ER β) receptors (de Groat, 2013; Birder *et al.*, 2002) [16, 9]. Among the mechanosensitive proteins expressed by urothelial cells are amiloride-sensitive ion channels (Du *et al.*, 2007) [17] and Piezo1, which has been implicated in stretch-evoked Ca²⁺ influx and ATP release in urothelial cell cultures (Miyamoto *et al.*, 2014) [26]. These mechanosensitive receptors respond to mechanical stretch by increasing intracellular calcium levels, leading to the subsequent release of ATP. The ATP released by urothelial cells can, in turn, signal neighboring urothelial cells, contributing to bladder sensory and functional regulation (Sun and Chai, 2006) [42].

Involvement of Urothelium in Caprine Bladder Smooth Muscle Contractility

In the present study, we examined the role of the urothelium in modulating cholinergic-mediated contractile responses in

caprine bladder smooth muscle. Our findings indicate that the presence of the urothelium significantly attenuates cholinergic-mediated contractions compared to bladder smooth muscle devoid of urothelium. This attenuation may be attributed to the release of relaxing molecules from the urothelium upon acetylcholine stimulation.

These findings align with previous studies demonstrating the inhibitory influence of urothelial-derived factors on bladder contractility. For instance, adenosine A1 receptors are known to be expressed in the uroepithelium of the bladder (Yu *et al.*, 2006) [12], and adenosine-induced relaxation has been primarily attributed to P1A1 receptor activation in normal rat bladders (Vesela *et al.*, 2011) [44]. Additionally, the impact of the mucosa on bladder strip contractility has been reported in various species. Maggi *et al.* (1987) [24] showed that guinea pig bladder strips lacking the mucosa exhibited enhanced contractility in response to substance P compared to full-thickness bladder strips, while responses to electrical field stimulation (EFS), histamine, or KCl remained unaffected by the presence or absence of the mucosa. Similarly, Pinna *et al.* (1992) [31] reported that bradykinin-induced contractions were significantly reduced in rat bladder strips when the mucosa was intact.

Further supporting these observations, Levin *et al.* (1995) [23] found that the feline bladder mucosa exerted an inhibitory effect on detrusor smooth muscle contractions. Likewise, studies in porcine bladder tissue suggested that the mucosa contributes to smooth muscle relaxation, hypothesizing that this effect originates from the urothelium or potentially other cellular components within the mucosal layer (Hawthorn *et al.*, 2000) [19]. Our findings reinforce the concept that the urothelium plays a critical role in modulating bladder smooth muscle function by releasing bioactive factors that influence detrusor contractility. Understanding this urothelial contribution could provide valuable insights into bladder physiology and potential therapeutic targets for conditions such as overactive bladder and urinary incontinence.

The Role of the Detrusor Muscle and Muscarinic Receptors in Bladder Function

Beyond the urothelium, the bladder wall primarily consists of the detrusor muscle, which plays a crucial role in bladder function. The ability of the detrusor muscle to contract over

a wide range of lengths and to relax appropriately is essential for the bladder's filling and micturition processes. These functions are regulated by intricate neural and hormonal control systems, and the detrusor muscle contains multiple receptor types and signaling pathways that mediate its activity.

Acetylcholine is the principal neurotransmitter governing bladder voiding, primarily acting through muscarinic receptors on smooth muscle cells (Wein, 2001) [47]. The distribution of muscarinic receptors within the bladder is not uniform; receptor density and sensitivity to muscarinic stimulation are highest in the bladder dome and lowest in the base. This gradient ensures efficient bladder emptying by optimizing contractile responses in different bladder regions.

Pharmacological, biochemical, and genetic studies have demonstrated that muscarinic receptors in the urinary bladder are heterogeneous in nature. At least three pharmacologically distinct muscarinic receptor subtypes—M1, M2, and M3—have been identified in the bladder (Akici *et al.*, 2000) [1]. These subtypes play distinct roles in modulating detrusor contractility, with M3 receptors primarily mediating contraction, while M2 receptors contribute to contractile responses under certain physiological and pathological conditions. Understanding the distribution and function of muscarinic receptors in the bladder is essential for developing targeted pharmacotherapies for disorders such as overactive bladder and urinary incontinence.

Acetylcholine-Mediated Contractile Response and Atropine Sensitivity in Caprine Bladder Smooth Muscle

Our study demonstrated that acetylcholine (ACh) induces a concentration-dependent contractile response in caprine detrusor muscle strips that lack the urothelium. These findings align with previous studies, where cholinergic muscarinic receptor agonists and electrical stimulation of intrinsic cholinergic nerves induced contractions in detrusor strips from healthy human bladders. Furthermore, contractile responses to muscarinic stimuli can be completely abolished by atropine in humans (Kinder and Mundy, 1985) [21], primates (Craggs *et al.*, 1986) [14], pigs, and rabbits (Sibley, 1984) [37].

Consistent with these observations, we found that the acetylcholine-mediated contractile response in caprine detrusor muscle was significantly attenuated in the presence of atropine. This confirms that the contractile response is mediated primarily via atropine-sensitive cholinergic mechanisms. Previous research has identified muscarinic receptor subtypes (M1, M2, and M3) in the human bladder, with M3 receptors predominating in detrusor muscle. In the sheep ureterovesical junction, contractions are mediated through M1, M2, and M3 receptor subtypes (Rivera *et al.*, 1992) [32]. Notably, co-activation of M2 receptors may enhance M3-mediated responses by inhibiting adenylate cyclase, suppressing sympathetically mediated detrusor relaxation, inactivating K⁺ channels, or activating nonspecific cation channels (Sellers and Chess-Williams, 2012) [36].

Stimulation of M3 receptors by ACh leads to phosphoinositol hydrolysis, which in turn releases intracellular Ca²⁺ and facilitates Ca²⁺ influx through L-type calcium channels. This process enhances contractile

sensitivity via myosin light chain phosphatase inhibition through Rho kinase activation (Andersson, 2002) [3].

Potential Non-Cholinergic Mechanisms in Caprine Bladder Contraction

Interestingly, despite atropine significantly reducing contractile responses, minimal ACh-induced contractions were still observed in caprine bladder smooth muscle. This suggests the involvement of a non-cholinergic system, possibly purinergic signaling, in ACh-mediated contractile responses.

Previous studies support this hypothesis. Atropine-resistant contractions have been observed in dysfunctional bladders, and these responses are not due to differential muscle sensitivity to ATP or cholinergic agonists (Bayliss *et al.*, 1999) [5]. Additionally, Sjögren *et al.* (1982) [39] reported atropine-resistant contractions in human bladder smooth muscle stimulated transmurally. Several studies propose that these responses originate from non-cholinergic nerves (Henderson & Roepke, 1934; Ambache & Zar, 1970; Dumsday, 1971; Burnstock *et al.*, 1972) [20, 2, 18, 10].

Atropine-resistant, nerve-mediated contractions occur in human detrusor muscle primarily from bladders with detrusor overactivity—whether idiopathic, neuropathic, or obstructive (Bayliss *et al.*, 1999) [5]—as well as in the detrusor muscle of small animals. These residual contractions can be abolished by pre-treatment with α , β -methylene ATP (ABMA), which desensitizes ionotropic purinergic receptors, suggesting that ATP acts as a co-transmitter alongside acetylcholine. Additionally, ATP release from nerves is independent of acetylcholine release (Pakzad *et al.*, 2016) [30], highlighting the potential for therapeutic strategies targeting ATP release to manage overactive bladder conditions.

However, atropine-resistant ACh-mediated contractile responses may occur through multiple mechanisms and vary across species and pathological conditions. While some studies attribute these responses to ATP-mediated purinergic signaling (Sjögren *et al.*, 1982) [39], others suggest direct smooth muscle stimulation due to increased excitability of the smooth muscle membrane (Tagliani *et al.*, 1997) [43]. Regardless of the underlying cause, alterations in detrusor muscle function in abnormal bladders present potential targets for drug development aimed at modulating bladder contractility.

Future Directions and Study Limitations

Our study suggests that an interaction between purinergic and cholinergic transmitter systems may exist at multiple levels within the micturition reflex arc. Further research is required to fully characterize this functional relationship.

Several limitations should be acknowledged

Sex-Based Differences: Our study was conducted exclusively on male caprine specimens. Male-specific organs, such as the penis and prostate, may influence bladder smooth muscle contraction, and these effects were not accounted for in our study.

Additional Influencing Factors: Various other physiological and pathological factors may contribute to bladder smooth muscle contraction, requiring further investigation.

Species-Specific Variations: While our findings align with data from human and animal studies, species-specific differences in bladder contractility mechanisms warrant further comparative research.

Future experiments should address these limitations by incorporating female specimens, exploring additional molecular pathways, and conducting comparative studies across different species.

Conclusion

Our findings demonstrate that acetylcholine-mediated contractile responses in caprine bladder smooth muscle occur via both atropine-sensitive and atropine-resistant pathways. Additionally, the presence of urothelium significantly modulates cholinergic-mediated contractile responses, likely through the release of relaxing factors. Moreover, our results align with previous studies on human urinary bladder smooth muscle, suggesting that the caprine bladder may serve as a relevant model for studying bladder physiology. Further investigations into caprine bladder smooth muscle function could enhance our understanding of urinary incontinence mechanisms and contribute to the development of more effective therapeutic strategies for human bladder dysfunction.

Acknowledgments

The author acknowledge Subash Chandra Parija, PhD, Post Doc (UK), Professor and Head, Department of Veterinary Pharmacology and Toxicology, Orissa University of Agriculture and Technology, Bhubaneswar-751 003, India, for his valuable guidance and support throughout the study.

References

1. Akici A, Karaalp A, Iskender E, Christopoulos A, El-Fakahany EE, Oktay S. Further evidence for the heterogeneity of functional muscarinic receptors in guinea pig gallbladder. *European Journal of Pharmacology*. 2000 Jan 24;388(1):115-123.
2. Ambache N, Zar MA. Non-cholinergic transmission by post-ganglionic motor neurones in the mammalian bladder. *Journal of Physiology*. 1970 Oct;210(3):761-783.
3. Andersson KE. Bladder activation: afferent mechanisms. *Urology*. 2002 May;59(5 Suppl 1):43-50.
4. Andersson KE, Yoshida M. Antimuscarinics and the overactive detrusor--which is the main mechanism of action? *European Urology*. 2003 Jan;43(1):1-5.
5. Bayliss M, Wu C, Newgreen D, Mundy AR, Fry CH. A quantitative study of atropine-resistant contractile responses in human detrusor smooth muscle, from stable, unstable and obstructed bladders. *Journal of Urology*. 1999 Nov;162(5):1833-1839.
6. Birder L, Andersson KE. Urothelial signaling. *Physiological Reviews*. 2013 Apr;93(2):653-680.
7. Sui G, Fry CH, Montgomery B, Roberts M, Wu R, Wu C. Purinergic and muscarinic modulation of ATP release from the urothelium and its paracrine actions. *American Journal of Physiology-Renal Physiology*. 2014 Feb 1;306(3):F286-298.
8. Birder LA, de Groat WC. Mechanisms of disease: involvement of the urothelium in bladder dysfunction. *Nature Clinical Practice Urology*. 2007 Jan;4(1):46-54.
9. Birder LA, Nealen ML, Kiss S, de Groat WC, Caterina MJ, Wang E, *et al.* Beta-adrenoceptor agonists stimulate endothelial nitric oxide synthase in rat urinary bladder urothelial cells. *Journal of Neuroscience*. 2002 Sep 15;22(18):8063-8070.
10. Burnstock G, Dumsday B, Smythe A. Atropine-resistant excitation of the urinary bladder: the possibility of transmission via nerves releasing a purine nucleotide. *British Journal of Pharmacology*. 1972 Mar;44(3):451-461.
11. Burnstock G. Purinergic cotransmission. *Experimental Physiology*. 2008;94:20-24.
12. Chai TC, Russo A, Yu S, Lu M. Mucosal signaling in the bladder. *Autonomic Neuroscience*. 2016 Oct;200:49-56.
13. Chess-Williams R, Chapple CR, Yamanishi T, Yasuda K, Sellers DJ. The minor population of M3-receptors mediate contraction of human detrusor muscle *in vitro*. *Journal of Autonomic Pharmacology*. 2001 Oct-Dec;21(5-6):243-248.
14. Craggs MD, Rushton DN, Stephenson JD. A putative non-cholinergic mechanism in urinary bladders of new but not old World primates. *Journal of Urology*. 1986 Dec;136(6):1348-1350.
15. D'Agostino G, Condino AM, Calvi V, Boschi F, Gioglio L, Barbieri A. Purinergic P2X3 heteroreceptors enhance parasympathetic motor drive in isolated porcine detrusor, a reliable model for development of P2X selective blockers for detrusor hyperactivity. *Pharmacological Research*. 2012 Jan;65(1):129-136.
16. de Groat WC. Highlights in basic autonomic neuroscience: contribution of the urothelium to sensory mechanisms in the urinary bladder. *Autonomic Neuroscience*. 2013 Oct;177(2):67-71.
17. Du S, Araki I, Mikami Y, Zakoji H, Beppu M, Yoshiyama M, *et al.* Amiloride-sensitive ion channels in urinary bladder epithelium involved in mechanosensory transduction by modulating stretch-evoked adenosine triphosphate release. *Urology*. 2007 Mar;69(3):590-595.
18. Dumsday B. Atropine-resistance of the urinary bladder innervation. *Journal of Pharmacy and Pharmacology*. 1971 Mar;23(3):222-225.
19. Hawthorn MH, Chapple CR, Cock M, Chess-Williams R. Urothelium-derived inhibitory factor(s) influences on detrusor muscle contractility *in vitro*. *British Journal of Pharmacology*. 2000 Feb;129(3):416-419.
20. Henderson VE, Roepke MH. The role of acetylcholine in bladder contractile mechanisms and in parasympathetic ganglia. *Journal of Pharmacology and Experimental Therapeutics*. 1934;51:97-111.
21. Kinder RB, Mundy AR. Atropine blockade of nerve-mediated stimulation of the human detrusor. *British Journal of Urology*. 1985 Aug;57(4):418-421.
22. Kullmann FA, Artim DE, Birder LA, de Groat WC. Activation of muscarinic receptors in rat bladder sensory pathways alters reflex bladder activity. *Journal of Neuroscience*. 2008 Feb 20;28(8):1977-1987.
23. Levin RM, Wein AJ, Krasnopolsky L, Atta MA, Ghoniem GM. Effect of mucosal removal on the response of the feline bladder to pharmacological stimulation. *Journal of Urology*. 1995 Apr;153(4):1291-1294.
24. Maggi CA, Santicioli P, Parlani M, Astolfi M, Patacchini R, Meli A. The presence of mucosa reduces the contractile response of the guinea-pig urinary

- bladder to substance P. *Journal of Pharmacy and Pharmacology*. 1987 Aug;39(8):653-655.
25. Matsui M, Motomura D, Karasawa H, Fujikawa T, Jiang J, Komiya Y, *et al*. Multiple functional defects in peripheral autonomic organs in mice lacking muscarinic acetylcholine receptor gene for the M3 subtype. *Proceedings of the National Academy of Sciences of the United States of America*. 2000 Aug 15;97(17):9579-9584.
 26. Miyamoto T, Mochizuki T, Nakagomi H, Kira S, Watanabe M, Takayama Y, *et al*. Functional role for Piezo1 in stretch-evoked Ca^{2+} influx and ATP release in urothelial cell cultures. *J Biol Chem*. 2014 Jun 6;289(23):16565-16575.
 27. Moro C, Leeds C, Chess-Williams R. Contractile activity of the bladder urothelium/lamina propria and its regulation by nitric oxide. *Eur J Pharmacol*. 2012 Jan 15;674(2-3):445-449.
 28. Moro C, Uchiyama J, Chess-Williams R. Urothelial/lamina propria spontaneous activity and the role of M3 muscarinic receptors in mediating rate responses to stretch and carbachol. *Urology*. 2011 Dec;78(6):1442.e9-15.
 29. O'Reilly BA, Kosaka AH, Knight GF, Chang TK, Ford AP, Rymer JM, *et al*. P2X receptors and their role in female idiopathic detrusor instability. *J Urol*. 2002;167:157-164.
 30. Pakzad M, Ikeda Y, McCarthy C, Kitney DG, Jabr RI, Fry CH. Contractile effects and receptor analysis of adenosine-receptors in human detrusor muscle from stable and neuropathic bladders. *Naunyn Schmiedebergs Arch Pharmacol*. 2016 Aug;389(8):921-929.
 31. Pinna C, Caratozzolo O, Puglisi L. A possible role for urinary bladder epithelium in bradykinin-induced contraction in diabetic rats. *Eur J Pharmacol*. 1992 Apr 22;214(2-3):143-148.
 32. Rivera L, Hernández M, Benedito S, Prieto D, García-Sacristán A. Mediation of contraction by cholinergic muscarinic receptors in the ureterovesical junction. *J Auton Pharmacol*. 1992 Jun;12(3):175-181.
 33. Robinson D, Cardozo L. Antimuscarinic drugs to treat overactive bladder. *BMJ*. 2012 Mar 27;344:e2130.
 34. Sadananda P, Chess-Williams R, Burcher E. Contractile properties of the pig bladder mucosa in response to neurokinin A: a role for myofibroblasts? *Br J Pharmacol*. 2008 Apr;153(7):1465-1473.
 35. Scarpero HM, Dmochowski RR. Muscarinic receptors: what we know. *Curr Urol Rep*. 2003 Dec;4(6):421-428.
 36. Sellers DJ, Chess-Williams R. Muscarinic agonists and antagonists: effects on the urinary bladder. *Handb Exp Pharmacol*. 2012;(208):375-400.
 37. Sibley GN. A comparison of spontaneous and nerve-mediated activity in bladder muscle from man, pig and rabbit. *J Physiol*. 1984 Sep;354:431-443.
 38. Silva I, Ferreirinha F, Magalhães-Cardoso MT, Silva-Ramos M, Correia-de-Sá P. Activation of P2Y6 receptors facilitates nonneuronal adenosine triphosphate and acetylcholine release from urothelium with the lamina propria of men with bladder outlet obstruction. *J Urol*. 2015 Oct;194(4):1146-1154.
 39. Sjögren C, Andersson KE, Husted S, Mattiasson A, Møller-Madsen B. Atropine resistance of transmurally stimulated isolated human bladder muscle. *J Urol*. 1982 Dec;128(6):1368-1371.
 40. Stenqvist J, Carlsson T, Winder M, Aronsson P. Functional atropine sensitive purinergic responses in the healthy rat bladder. *Auton Neurosci*. 2020 Sep;227:102693.
 41. Stenqvist J, Winder M, Carlsson T, Aronsson P, Tobin G. Urothelial acetylcholine involvement in ATP-induced contractile responses of the rat urinary bladder. *Eur J Pharmacol*. 2017 Aug 15;809:253-260.
 42. Sun Y, Chai TC. Augmented extracellular ATP signaling in bladder urothelial cells from patients with interstitial cystitis. *Am J Physiol Cell Physiol*. 2006 Jan;290(1):C27-34.
 43. Tagliani M, Candura SM, Di Nucci A, Franceschetti GP, D'Agostino G, Ricotti P, *et al*. A re-appraisal of the nature of the atropine-resistant contraction to electrical field stimulation in the human isolated detrusor muscle. *Naunyn Schmiedebergs Arch Pharmacol*. 1997 Dec;356(6):750-755.
 44. Vesela R, Aronsson P, Tobin G. Functional and morphological examinations of P1A1 purinoceptors in the normal and inflamed urinary bladder of the rat. *Auton Neurosci*. 2011 Jan 20;159(1-2):26-31.
 45. Vial C, Evans RJ. P2X receptor expression in mouse urinary bladder and the requirement of P2X(1) receptors for functional P2X receptor responses in the mouse urinary bladder smooth muscle. *Br J Pharmacol*. 2000 Dec;131(7):1489-1495.
 46. Vijayakumar C, Kathirvel K, Sardar KK, Parija SC. Effect of KATP channel openers on myogenic and neurogenic responses in goat urinary bladder. *Indian J Exp Biol*. 2007 Feb;45(2):185-193.
 47. Wein AJ. Pharmacological agents for the treatment of urinary incontinence due to overactive bladder. *Expert Opin Investig Drugs*. 2001 Jan;10(1):65-83.
 48. Winder M, Tobin G, Zupančič D, Romih R. Signalling molecules in the urothelium. *Biomed Res Int*. 2014;2014:297295.
 49. Yu W, Zacharia LC, Jackson EK, Apodaca G. Adenosine receptor expression and function in bladder uroepithelium. *Am J Physiol Cell Physiol*. 2006 Aug;291(2):C254-265.