



International Journal of Research in Agronomy

E-ISSN: 2618-0618
P-ISSN: 2618-060X
© Agronomy
NAAS Rating: 5.20
www.agronomyjournals.com
2025; 8(3): 383-385
Received: 13-01-2025
Accepted: 21-02-2025

Dr. Prashanthi Sandepogu
Department Faculty of Botany,
TSWRDC (W), Nirmal, Kakatiya
University, Telangana, India

Dr. Mamta J Patange
Assistant Professor,
Department of Agronomy,
VNMKV, Parbhani, Maharashtra,
India

Ashutosh Srivastava
Field Assistant, KVK-Pratapgarh,
Uttar Pradesh, India

Cennichon Khodang
Ph.D. Research Scholar,
Department of Agricultural
Economics, Kerala Agricultural
University, Vellanikkara, Thrissur,
Kerala, India

Dr. Satwinderjit Kaur
Associate Professor, Department of
Soil Science, PAU-KVK,
Gurdaspur, Punjab, India

Khushboo Gupta
Teaching Associate,
Department of Processing and
Agricultural Structure, Chandra
Shekhar Azad University of
Agriculture & Technology,
Kanpur, Uttar Pradesh

Dr. Santosh Kumar Vishwakarma
Senior Scientist and Head, Krishi
Vigyan Kendra Jamunabad,
Lakhimpur Kheri, Uttar Pradesh,
India

Shubhendu Singh
Ph.D. Research Scholar,
Department of Agronomy,
Banaras Hindu University,
Varanasi, Uttar Pradesh, India

Corresponding Author:
Dr. Prashanthi Sandepogu
Department Faculty of Botany,
TSWRDC (W), Nirmal, Kakatiya
University, Telangana, India

Crop residue management: Balancing agricultural sustainability and environmental impact

Prashanthi Sandepogu, Mamta J Patange, Ashutosh Srivastava, Cennichon Khodang, Satwinderjit Kaur, Khushboo Gupta, Santosh Kumar Vishwakarma and Shubhendu Singh

DOI: <https://www.doi.org/10.33545/2618060X.2025.v8.i3e.2688>

Abstract


Crop residue management (CRM) is a pivotal component of modern agricultural systems, influencing both agricultural sustainability and environmental health. Crop residues, the byproducts of crop production, can either serve as a resource for enhancing soil health and productivity or become a source of environmental pollution if mismanaged. This paper provides an in-depth analysis of the challenges and opportunities associated with CRM, focusing on its impact on soil fertility, greenhouse gas emissions, air quality, and biodiversity. It evaluates various CRM practices, including residue retention, incorporation, mulching, composting, and bioenergy production, and discusses their implications for sustainable agriculture. The paper concludes with recommendations for integrated CRM strategies that balance agricultural productivity with environmental conservation, supported by a comprehensive review of recent literature.

Keywords: Crop residue management (CRM), agricultural sustainability, environmental conservation, soil fertility and biodiversity

1. Introduction

Crop residues, including stalks, leaves, husks, and roots, are an inevitable byproduct of agricultural production. Globally, crop residues are generated in vast quantities, with estimates suggesting annual production exceeds 3.5 billion tons (Lal, 2005) ^[1]. Traditionally viewed as waste, these residues are increasingly recognized as a valuable resource for enhancing soil health, reducing erosion, and contributing to sustainable farming systems. However, improper management of crop residues, such as open burning or excessive removal, can lead to significant environmental degradation, including air pollution, soil nutrient depletion, and increased greenhouse gas emissions.

As global agricultural systems face the dual challenges of feeding a growing population and mitigating climate change, effective CRM has emerged as a key strategy for achieving sustainable development goals. This paper examines the role of CRM in balancing agricultural sustainability and environmental impact (fig. 1). It highlights the benefits and drawbacks of current CRM practices and proposes integrated approaches to optimize residue utilization while minimizing negative environmental consequences.



2. The Role of Crop Residues in Agricultural Systems

Crop residues are a vital component of agroecosystems, contributing to soil health, water conservation, and nutrient cycling. Their role in sustainable agriculture can be understood through the following key benefits:

2.1 Soil Fertility Enhancement

Crop residues decompose to release organic matter, which improves soil structure, water retention, and nutrient availability. The incorporation of residues into the soil increases soil organic carbon (SOC), which is critical for maintaining soil fertility and productivity (Blanco-Canqui & Lal, 2009) ^[2]. Residues also act as a slow-release fertilizer, providing essential nutrients such as nitrogen, phosphorus, and potassium as they decompose (Kumar & Goh, 1999) ^[3].

2.2 Erosion Control

Residue cover on the soil surface reduces soil erosion by wind and water. This protective layer minimizes the impact of raindrops, slows surface runoff, and prevents the detachment and transport of soil particles (Blanco-Canqui *et al.*, 2013) ^[4]. In regions prone to erosion, such as sloping lands and arid areas, residue retention is particularly effective in maintaining soil integrity and productivity.

2.3 Carbon Sequestration

Incorporating crop residues into the soil enhances carbon storage, mitigating climate change. Residues contribute to the formation of stable soil organic matter, which sequesters carbon for long periods (Lal, 2005) ^[1]. This process not only reduces atmospheric CO₂ levels but also improves soil health and resilience.

2.4 Biodiversity Support

Crop residues provide habitat and food for soil microorganisms and fauna, promoting ecosystem resilience. The decomposition of residues supports a diverse microbial community, which plays a crucial role in nutrient cycling and organic matter decomposition (Bastida *et al.*, 2008) ^[5]. Additionally, residues offer shelter for beneficial insects and other organisms that contribute to pest control and pollination.

Despite these benefits, the mismanagement of crop residues, particularly through open burning, has significant environmental consequences, including air pollution, loss of nutrients, and contributions to global warming.

3. Environmental Impacts of Crop Residue Mismanagement

The improper handling of crop residues poses several environmental challenges:

3.1 Air Pollution

Open burning of residues is a common practice in many regions, particularly in developing countries. This practice releases large quantities of particulate matter (PM_{2.5} and PM₁₀), carbon monoxide (CO), volatile organic compounds (VOCs), and other pollutants into the atmosphere (Gupta *et al.*, 2004) ^[6]. These emissions contribute to respiratory diseases, smog formation, and reduced visibility, posing significant public health risks.

3.2 Greenhouse Gas Emissions

Burning crop residues releases carbon dioxide (CO₂), methane

(CH₄), and nitrous oxide (N₂O), all of which are potent greenhouse gases (Singh *et al.*, 2014) ^[7]. The combustion of

residues also destroys organic matter that could otherwise contribute to soil carbon sequestration, exacerbating climate change.

3.3 Soil Degradation

Removing or burning residues depletes soil organic matter, reducing fertility and long-term productivity. The loss of organic matter leads to soil compaction, reduced water infiltration, and increased susceptibility to erosion (Blanco-Canqui & Lal, 2009)^[2]. Over time, this degradation can render agricultural lands less productive and more vulnerable to extreme weather events.

3.4 Water Pollution

Residue runoff can carry pesticides and fertilizers into water bodies, causing eutrophication and harming aquatic ecosystems. The accumulation of nutrients such as nitrogen and phosphorus in water bodies leads to algal blooms, oxygen depletion, and the death of aquatic organisms (Carpenter *et al.*, 1998)^[8].

4. Crop Residue Management Practices

Effective CRM requires a combination of practices tailored to local conditions. Key strategies include:

4.1 Residue Retention and Mulching

Leaving residues on the field as mulch conserves soil moisture, suppresses weeds, and enhances soil organic matter. Mulching is particularly beneficial in conservation agriculture systems, where it complements practices such as zero tillage and crop rotation (Kassam *et al.*, 2009)^[9].

4.2 Residue Incorporation

Ploughing residues into the soil accelerates decomposition, releasing nutrients and improving soil structure. However, excessive incorporation can lead to nitrogen immobilization, where microorganisms temporarily lock up nitrogen, making it unavailable to plants.

4.3 Bioenergy Production

Crop residues can be converted into biofuels, biogas, or biochar, providing renewable energy while reducing reliance on fossil fuels. Biochar, a carbon-rich product of pyrolysis, offers additional benefits for soil health and carbon sequestration (Lehmann *et al.*, 2011)^[10].

4.4 Composting

Composting residues with other organic waste produces nutrient-rich organic fertilizer, reducing the need for synthetic inputs and promoting circular agriculture (Bernal *et al.*, 2009)^[11].

4.5 Livestock Feed

In some regions, crop residues are used as fodder for livestock, providing an additional income stream for farmers while reducing waste (Giller *et al.*, 2021)^[12].

5. Challenges in Crop Residue Management

Despite the availability of various CRM practices, several barriers hinder their widespread adoption:

5.1 Lack of Awareness and Knowledge

Many farmers are unaware of the benefits of alternative CRM practices or lack the technical know-how to implement them (Jat *et al.*, 2020)^[13].

5.2 Economic Constraints

The cost of machinery for residue incorporation or bioenergy production can be prohibitive for smallholder farmers (Singh and Sidhu, 2014)^[14].

5.3 Policy and Institutional Gaps

Inadequate policies and incentives for sustainable CRM limit its adoption (Ali Mekouar, 2017)^[15].

5.4 Regional Variability

The effectiveness of CRM practices varies depending on climate, soil type, and cropping systems, necessitating context-specific solutions (Blanco-Canqui *et al.*, 2013)^[4].

6. Integrated Approaches to Crop Residue Management

To address these challenges, integrated CRM strategies are needed, combining technological, economic, and policy intervention (fig. 2) as:



Fig. 2: Integrated Approaches to Crop Residue Management

6.1 Promoting Conservation Agriculture

Encouraging practices such as zero tillage, crop rotation, and residue retention can enhance sustainability (Kassam *et al.*, 2009)^[9].

6.2 Subsidies and Incentives

Governments should provide financial support for the adoption of sustainable CRM practices and technologies (Singh and Sidhu, 2014)^[14].

6.3 Capacity Building

Training programs and extension services can equip farmers with the knowledge and skills needed for effective CRM (Jat *et al.*, 2020)^[13].

6.4 Research and Innovation

Investing in research to develop cost-effective and region-specific CRM technologies is essential (Blanco-Canqui *et al.*, 2013)^[4].

6.5 Policy Frameworks

Strengthening policies to regulate residue burning and promote sustainable practices is critical for long-term impact (FAO, 2017)^[15].

7. Conclusion

Crop residue management is a cornerstone of sustainable agriculture, offering significant benefits for soil health, climate mitigation, and environmental conservation. However, the mismanagement of residues poses serious environmental risks, underscoring the need for integrated and context-specific solutions. By adopting a holistic approach that combines technological innovation, policy support, and farmer engagement, it is possible to balance agricultural productivity with environmental sustainability. Effective CRM not only enhances farm resilience but also contributes to global efforts to combat climate change and preserve natural resources for future generations.

8. References

1. Lal R. World crop residues production and implications of its use as a biofuel. *Environ Int.* 2005;31(4):575-584.
2. Blanco-Canqui H, Lal R. Crop residue removal impacts on soil productivity and environmental quality. *Crit Rev Plant Sci.* 2009;28(3):139-163.
3. Kumar K, Goh KM. Crop residues and management practices: effects on soil quality, soil nitrogen dynamics, crop yield, and nitrogen recovery. *Adv Agron.* 1999;68:197-319.
4. Blanco-Canqui H. Crop residue removal for bioenergy reduces soil carbon pools: how can we offset carbon losses? *Bioenergy Res.* 2013;6:358-371.
5. Bastida F, Zsolnay A, Hernández T, García C. Past, present and future of soil quality indices: a biological perspective. *Geoderma.* 2008;147(3-4):159-171.
6. Gupta PK, Sahai S, Singh N, Dixit CK, Singh DP, Sharma C, *et al.* Residue burning in rice–wheat cropping system: Causes and implications. *Curr Sci.* 2004;1713-1717.
7. Singh A, Phogat VK, Dahiya R, Batra SD. Impact of long-term zero till wheat on soil physical properties and wheat productivity under rice–wheat cropping system. *Soil Tillage Res.* 2014;140:98-105.
8. Carpenter SR, Caraco NF, Correll DL, Howarth RW, Sharpley AN, Smith VH. Nonpoint pollution of surface waters with phosphorus and nitrogen. *Ecol Appl.* 1998;8(3):559-568.
9. Kassam A, Friedrich T, Shaxson F, Pretty J. The spread of conservation agriculture: justification, sustainability and uptake. *Int J Agric Sustain.* 2009;7(4):292-320.
10. Lehmann J, Rillig MC, Thies J, Masiello CA, Hockaday WC, Crowley D. Biochar effects on soil biota—a review. *Soil Biol Biochem.* 2011;43(9):1812-1836.
11. Bernal MP, Alburquerque JA, Moral R. Composting of animal manures and chemical criteria for compost maturity assessment: A review. *Bioresour Technol.* 2009;100(22):5444-5453.
12. Giller KE, Hijbeek R, Andersson JA, Sumberg J. Regenerative agriculture: an agronomic perspective. *Outlook Agric.* 2021;50(1):13-25.
13. Jat ML, Chakraborty D, Ladha JK, Rana DS, Gathala MK, McDonald A, *et al.* Conservation agriculture for sustainable intensification in South Asia. *Nat Sustain.* 2020;3(4):336-343.
14. Singh Y, Sidhu HS. Management of cereal crop residues for sustainable rice-wheat production system in the Indo-Gangetic plains of India. *Proc Indian Nat Sci Acad.* 2014;80(1):95-114.
15. Mekouar MA. Food and agriculture organization of the United Nations (FAO). *Yearbook of International Environmental Law.* 2017;28:506-520.