



ISSN Print: 2664-9926

ISSN Online: 2664-9934

NAAS Rating: 4.82

IJBS 2025; 7(3): 39-41

www.biologyjournal.net

Received: 12-01-2025

Accepted: 16-02-2025

A Abinaya

Department of Animal
Nutrition, Veterinary College
and Research Institute, Tamil
Nadu Veterinary and Animal
Sciences University, Tamil
Nadu, India

R Balamurugan

Department of Animal
Nutrition, Veterinary College
and Research Institute, Tamil
Nadu Veterinary and Animal
Sciences University, Tamil
Nadu, India

C Kadhivelan

Department of Animal
Nutrition, Veterinary College
and Research Institute, Tamil
Nadu Veterinary and Animal
Sciences University, Tamil
Nadu, India

Corresponding Author:

A Abinaya

Department of Animal
Nutrition, Veterinary College
and Research Institute, Tamil
Nadu Veterinary and Animal
Sciences University, Tamil
Nadu, India

Exploring the proximate nutrients of cotton gin trash: An alternate roughage feed to ruminants

A Abinaya, R Balamurugan and C Kadhivelan

DOI: <https://www.doi.org/10.33545/26649926.2025.v7.i3a.316>

Abstract

Cotton gin trash is a by-product of cotton ginning process and serves as an alternate source of roughage in dairy animal feed. Proximate analysis of coarse cotton gin trash samples showed the following nutrient composition: Moisture 11.2%, Crude protein 10.9%, Crude fibre 37.68%, Ether extract 3.2%, Total ash 7.25%, Acid insoluble ash (AIA) 2.3%. The high fibre and moderate protein content make cotton gin trash an important roughage source especially during times of fodder scarcity. Additionally, enhancing its nutrient profile could further increase its value as a beneficial agricultural byproduct for livestock nutrition.

Keywords: Cotton gin trash, proximate, fodder scarcity

Introduction

Cotton is used as a fabric dating back to 2000 BC in India. The industrial revolution brought significant advancements in cotton processing which fuelled the global expansion of cotton market. The processing of cotton produces a variety of by-products that include whole cottonseed, cottonseed meal, cottonseed hulls, cotton mote, textile mill waste, and cotton gin trash. Whole cottonseed is extracted during the ginning process and is primarily used for cottonseed oil production or as ruminant feed (Kunkle *et al.*, 2004) [7]. As it does not require further processing before feeding, which helps reduce costs (Poore and Rogers, 1995) [9]. However, two challenges associated with feeding whole cottonseed are its high fat content and the presence of gossypol, a toxic compound that can affect animal health. Cottonseed hulls, another by-product from cottonseed oil extraction, have relatively low nutritional value. They contain 45% Total Digestible Nutrients (TDN) and crude protein levels ranging from 4% to 5% (NRC, 2000; Waller, 2006) [5, 6]. Despite this, cottonseed hulls can still be used in animal feed, especially in ruminant diets where fiber content is crucial.

In addition to these, cotton mote and textile mill waste are also generated in the processing of cotton. These by-products, while not as widely studied, may offer alternative uses in animal feed or other agricultural applications. Cotton gin trash, in particular, has gained attention as a valuable source of roughage for livestock, especially during times of fodder scarcity [4].

The diverse by-products of cotton processing highlight the crop's versatility and its potential to contribute not only to the textile industry but also to agricultural and livestock sectors. Proper management and utilization of these by-products can help reduce waste and enhance the sustainability of cotton farming.

Cotton gin trash is a by-product of the cotton ginning process and serves as a valuable source of roughage in ruminant diets. It is composed of boll residues, leaves, stems, and lint, which results in a bulky feed that is high in fibre but low in energy. While this composition makes it an effective roughage source, its high fibre content can limit its energy value for livestock. To improve its utilization, reducing the particle size by grinding cotton gin trash makes it easier to handle and increases its intake by animals.

Palatability of cotton gin trash is generally good, provided that it does not exceed 40% of the total diet, as higher inclusion levels may lead to reduced feed intake (Lalor *et al.*, 1975) [10]. Unlike whole cottonseed, gossypol is not considered a significant issue in cotton gin trash, as it contains only a small number of immature seeds, which minimizes the risk of toxicity in livestock.

The current study aims to analyze the proximate composition of cotton gin trash to assess its suitability as an alternative roughage source for ruminant feeding. This research is crucial for determining the nutritional value of cotton gin trash, as it has the potential to supplement traditional feed resources, especially during periods of feed scarcity. Additionally, with proper processing and formulation, cotton gin trash could serve as an affordable and sustainable feed option in livestock nutrition.

Further research could explore methods to enhance the digestibility and nutritional content of cotton gin trash, such as through fermentation, enzyme supplementation, or combining it with other feed ingredients. By improving its nutritional profile, cotton gin trash could become a more widely adopted feed resource, contributing to the sustainability of livestock production and reducing agricultural waste.

Materials and Methods

Six coarse-type cotton gin trash samples were collected from various locations within the Tirupur district. The samples were carefully dried and ground to pass through a 2-mm screen to ensure uniformity in size for analysis. The proximate composition of these samples was determined using standard procedures as outlined by the AOAC (2012) [1]. All results were expressed as percentages on a dry matter basis to ensure accuracy and consistency in comparison. By exploring these aspects, the study could contribute to expanding the use of cotton gin trash as a cost-effective and environmentally friendly feed resource in animal agriculture.

Table 1: Proximate Composition of Cotton Gin Trash (on Dry Matter Basis)

Proximate Nutrients	Mean $\pm$ SE
Moisture (%)	11.23 $\pm$ 0.43
Crude Protein (%)	10.9 $\pm$ 0.67
Crude Fibre (%)	37.6 $\pm$ 0.47
Ether Extract (%)	3.2 $\pm$ 0.28
Total Ash (%)	7.2 $\pm$ 0.67
Acid Insoluble Ash (%)	2.3 $\pm$ 0.20
Nitrogen Free Extract (%)	40.06 $\pm$ 0.71
Each value is a mean of three observations	

Results and Discussion

The results of the proximate analysis of coarse-type cotton gin trash samples revealed the following composition (Table 1): moisture 11.2%, crude protein 10.9%, crude fiber 37.68%, ether extract 3.2%, total ash 7.25%, and acid-insoluble ash (AIA) 2.3%. These values are consistent with findings from previous studies, such as Bader *et al.* (1998) [2], who noted that the nutrient composition of cotton gin trash can vary based on factors such as the variety of cotton, geographical location, harvest method, and the presence of impurities. This variability is further highlighted by Myer *et al.* (2008) [3], who reported that crude protein in cotton gin trash ranges from 2% to 16%, moisture content from 2% to 60%, and crude fiber between 28% and 45%.

Moisture content in cotton gin trash can be highly variable, depending on the time of harvest and environmental conditions. For example, cotton gin trash harvested in different seasons or under varying climatic conditions may have significant differences in moisture, which could affect its storage and use in animal feed. Cotton gin trash also

contains a moderate to high amount of ash, with levels ranging from 8% to 28% (Lalor *et al.*, 1975) [10], due to the presence of non-combustible minerals like silica, calcium, potassium, and magnesium.

In terms of its nutritional profile, cotton gin trash is considered to have moderate protein and energy content, making it comparable to low-quality forage such as bermudagrass hay. Lalor *et al.* (1975) [10] found that the palatability of gin trash is generally acceptable as long as it does not exceed 40% of the total diet, beyond which feed intake may decrease. Notably, gossypol, a toxic compound found in cottonseed, does not pose a significant concern in cotton gin trash due to the limited presence of immature seeds. This makes it a safer alternative to whole cottonseed, particularly in ruminant diets.

The low cost of cotton gin trash is one of its key advantages, as it is often available at little more than the transportation cost from the gin (Kunkle *et al.*, 2004) [7]. This affordability, combined with its fiber content, makes cotton gin trash an attractive option as a roughage source, especially during periods of feed scarcity.

Recent studies have also explored the potential benefits of cotton gin trash in livestock feeding. For example, cotton gin trash has been successfully used as an unconventional feedstuff in the feeding of Mecheri sheep, where it was found to improve weight gain and feed efficiency (Nagarajan *et al.*, 2023) [8]. The chemical composition of cotton gin trash in this study was found to contain 15.7% crude protein and 31.9% crude fiber, further supporting its potential as a useful feed resource.

One of the major challenges in utilizing cotton gin trash as livestock feed is its low bulk density, which can make transportation difficult and costly (Kunkle *et al.*, 2004) [7]. Despite this, the low cost and potential benefits of incorporating cotton gin trash into animal diets make it a promising option. Further research is needed to explore methods for increasing the protein content of cotton gin trash, such as through fermentation or supplementation, to enhance its nutritional value and expand its potential as a valuable agricultural by-product for livestock nutrition.

The variability in protein content, which is influenced by factors such as the number of immature seeds and environmental conditions during the growing season (Lalor *et al.*, 1975) [10], underscores the importance of further research into optimizing the use of cotton gin trash in livestock feeding. Such efforts could lead to better utilization of this by-product, reducing waste and improving the sustainability of livestock production systems.

Conclusion

In conclusion, cotton gin trash is a valuable alternative roughage source, containing a moderate level of protein and a high level of fiber. These characteristics make it suitable for use as a primary roughage source, particularly during periods of fodder scarcity. Given its high fiber content, cotton gin trash can effectively replace conventional roughages in ruminant diets, providing a cost-effective and sustainable solution for livestock feeding.

However, further research is needed to optimize its use, particularly regarding its digestibility, protein content, and potential improvements in its palatability and nutrient composition. Enhancing the protein content of cotton gin trash through processing or supplementation could increase its nutritional value and make it an even more viable option

for livestock nutrition. Additionally, exploring efficient methods for managing its low bulk density and improving its handling and transportation will further facilitate its adoption as a widespread feed resource, reducing waste and contributing to the sustainability of livestock production systems.

References

1. AOAC. Official methods of analysis. Washington, D.C., USA: Association of Official Analytical Chemists; 2012.
2. Bader MJ, Bramwell RK, Stewart RL, Hill GM. Gin trash studies conducted in Georgia. In: Beltwide Cotton Production Research Proceedings. 1998;2:1698-1699.
3. Myer RO. Cotton Gin Trash: Alternative Roughage Feed for Beef Cattle: AN177/AN189, 11/2007, EDIS series. Institute of Food and Agricultural Science, University of Florida; 2008.
4. Hayavadana GR, Sangam P, N Jaishankar, KP Chethan, MC Shivakumar, B Ramachandra, R Guruprasad, K Gowda H. Utilization of perennial fodder COFS-29 as sole roughage source in Hassan sheep lambs under intensive rearing system. *Int J Vet Sci Anim Husbandry*. 2024;9(1):76-78.
5. Waller JC. By-products and unusual feedstuffs. *Feedstuffs Reference Issue and Buyer's Guide*. 2006;77:18.
6. NRC. Nutrient Requirements of Beef Cattle. 7th rev. ed. Washington, DC: National Academy Press; 2000.
7. Kunkle WE, Stewart RL, Brown WF. Using by-product feeds in beef supplementation programs. 2004.
8. Nagarajan SB, Ramakrishnan S, Muralidharan J, Vasani P, Sivakumar K, Thiruvankadan AK. Effect of cotton gin trash supplementation as unconventional feedstuff on feed intake and production characteristics of Mecheri sheep of India. *Sustainability*. 2023;15(13):10500.
9. Poore MH, Rogers G. Feeding whole cottonseed and other cotton by-products to beef cattle; 1995.
10. Lalor WF, Jones JK, Slater GA. Cotton gin trash as a ruminant feed. Cotton Gin and Oil Mill Press. 1975;28.