



International Journal of Veterinary Sciences and Animal Husbandry



ISSN: 2456-2912
NAAS Rating: 4.61
VET 2025; 10(4): 148-150
© 2025 VET
www.veterinarypaper.com
Received: 20-02-2025
Accepted: 27-03-2025

Keerthi Nallareddy
Interns at College of Veterinary
Science, Tirupati, Andhra
Pradesh, India

Revathi Reddy Sadu
Interns at College of Veterinary
Science, Tirupati, Andhra
Pradesh, India

Jaya Prakash Tellapati
Veterinary Assistant Surgeon,
Veterinary Poly Clinic, Nellore,
Andhra Pradesh, India

Use of hydrogen peroxide and normal saline flushing in clearing feline urethral obstruction: A retrospective study of 30 cases

Keerthi Nallareddy, Revathi Reddy Sadu and Jaya Prakash Tellapati

DOI: <https://www.doi.org/10.22271/veterinary.2025.v10.i4c.2190>

Abstract

Urethral obstruction is a life-threatening urological emergency in male cats. This study investigates the efficacy of hydrogen peroxide and normal saline (1:1) flushing via penile catheter as a treatment modality for clearing urethral obstruction in tom cats. A retrospective study of 30 intact male cats with urethral obstruction was conducted, and success rate of 96.7% was recorded. The procedure is minimally invasive, with no complications recorded.

Keywords: Urethral obstruction, tom cats, mucus plug, urethral catheterization, hydrogen peroxide, flushing

1. Introduction

Feline urethral obstruction is a common and severe urological emergency in male cats, often resulting in acute kidney injury, bladder rupture, and death if not addressed promptly. The obstruction is frequently caused by a variety of factors, including urethral mucus plugs, calculi, stricture, or neoplasia (Pachtinger, 2020) [3]. Male cats are particularly prone to urethral obstructions due to their longer and narrower urethra compared to females (George and Grauer, 2017) [1]. In addition, diet plays a significant role in the development of urethral obstructions, with dry food diet contributing to a reduced water intake and increased risk of urolithiasis (Segev *et al.*, 2011) [4]. Traditional management of feline urethral obstruction generally involves catheterization, but there is growing interest for exploring less invasive techniques for clearing obstructions in feline practitioners. Hydrogen peroxide has been suggested as a potential aid for breaking down obstructions in the urinary tract due to its effervescent property, which help to mechanically dislodge and dissolve debris (Xu *et al.*, 2020) [5]. The combination of hydrogen peroxide and normal saline for flushing is being explored as a less invasive, effective treatment for feline urethral obstruction. This study aims to evaluate the safety, effectiveness, and complications associated with hydrogen peroxide and normal saline flushing for clearing urethral obstruction in male cats.

2. Materials and Methods

30 male cats presented with symptoms of urinary obstruction to Veterinary Poly Clinic, Nellore, Andhra Pradesh during the period from May 2024 to September 2024 were included in the study. The details of the cases are presented in Table.1. The inclusion criteria for the study were male cats presenting with clinical signs of urethral obstruction such as anuria, stranguria, excessive perineal licking, and distended urinary bladder. Exclusion criteria included cats with a history of bladder rupture, neoplastic conditions, or prior urethral surgeries. Each animal was sedated using xylazine @1mg/kg b.wt I/M and atropine @0.045 mg/kg b.wt I/M to facilitate intervention, with additional ketamine @ 22 mg/kg b.wt I/M administered for analgesia as and when required. The animals were positioned in lateral recumbency, and the preputial hair was clipped and prepped for intervention.

Corresponding Author:
Keerthi Nallareddy
Interns at College of Veterinary
Science, Tirupati, Andhra
Pradesh, India

A 22 G I.V. cannula lubricated with sterile 2% lignocaine gel was inserted into the tip of the urethra (Figure 1). The needle was withdrawn and the urethra was flushed with a 1:1 mixture of 3% hydrogen peroxide and sterile normal saline, allowing the solution to dwell for one minute before further advancement. In cases where hydrogen peroxide and saline infusion failed to relieve the obstruction, urethral catheterization was performed using no 3 tom catheter (Figure 2). The urethral catheter was secured with stay sutures to prevent dislodging. After the obstruction was cleared, the urinary bladder was flushed multiple times with Normal

Saline till returning fluid was normal in color. Diluted Metronidazole was injected into the urinary bladder at the end of flushing. Post-procedural care included isotonic crystalloid fluid therapy @ 10ml/kg b.wt I/V, Inj.meloxicam @ 0.2 mg/kg b.wt s/c for 3 days, Inj.amoxicillin and sulbactam @15mg/kg I/M for 5 days and advised to use Reno protective syrup Advaren @ 0.5ml/kg b.wt

¹Syp.Advaren, SAVAVET, Lalwani Plaza, Viman Nagar, Pune, India, 411014

Table 1: Details of cases included in the study

S. No	Breed	Age	Body weight	Food habit	Site of mucosal plug	Clearance of obstruction with Hydrogen peroxide
1	Persian	2 years	4 kgs	Dry food	Penile orifice	Yes
2	Persian	3 years	4.3 kgs	Dry food	Penile orifice	Yes
3	Persian	2.5 years	3.2 kgs	Dry food	Penile orifice	Yes
4	Persian	3 years	3.8 kgs	Dry food	Not visible	Yes
5	Persian	4 years	3.6 kgs	Dry food	Penile orifice	Yes
6	Persian	3.2 years	4.3 kgs	Dry food	Penile orifice	Yes
7	Persian	4 years	5.6 kgs	Dry food	Penile orifice	Yes
8	Persian	3 years	4.6kgs	Dry food	Penile orifice	Yes
9	Persian	2.8 years	4.9 kgs	Dry food	Penile orifice	Yes
10	Persian	3 years	5.1 kgs	Dry food	Not visible	No
11	Persian	3.5 years	3.4 kgs	Dry food	Penile orifice	Yes
12	Persian	2 years	4.8 kgs	Dry food	Not visible	Yes
13	Persian	2.6 years	4.1 kgs	Dry food	Penile orifice	Yes
14	Persian	3.5 years	4.7 kgs	Dry food	Penile orifice	Yes
15	Persian	3 years	3.9 kgs	Dry food	Penile orifice	Yes
16	Persian	2 years	3.3 kgs	Dry food	Not visible	Yes
17	Persian	4 years	4.5 kgs	Dry food	Penile orifice	Yes
18	Persian	2.4 years	2.8 kgs	Dry food	Penile orifice	Yes
19	Persian	3 years	3.3 kgs	Dry food	Not visible	Yes
20	Persian	3.8 years	2.5 kgs	Dry food	Penile orifice	Yes
21	Persian	3.2 years	2.9 kgs	Dry food	Penile orifice	Yes
22	Persian	3.6 years	5 kgs	Dry food	Penile orifice	Yes
23	Persian	2.7 years	4.2 kgs	Dry food	Not visible	Yes
24	Persian	2.5 years	3.4 kgs	Dry food	Penile orifice	Yes
25	Persian	3.2 years	2.9 kgs	Dry food	Not visible	Yes
26	Persian	2.3 years	3.1 kgs	Dry food	Penile orifice	Yes
27	Persian	3.7 years	4.2 kgs	Dry food	Penile orifice	Yes
28	Persian	2 years	1.8 kgs	Dry food	Penile orifice	Yes
29	Persian	2.8 years	3.6 kgs	Dry food	Penile orifice	Yes
30	Persian	3.5 years	3.8 kgs	Dry food	Not visible	Yes



Fig 1: Lubricated I/V cannula being inserted into the tip of the urethra for flushing



Fig 2: Urethral catheterization using no 3 tom catheter

3. Results

The 30 cats treated with hydrogen peroxide and saline flushing, 96.7% (N=29) showed successful clearance of the obstruction, with immediate urine outflow following the procedure (Figure 4). During the study, the most common

finding was the presence of hematuria in 26 cases, suggesting hemorrhagic cystitis. In 22 cases, pasty white plug was observed at the penile orifice, which was identified as the primary cause of the obstruction (Figure 5). In the other seven cases, the mucosal plug was not visible. In all 29 cases, the

obstruction was cleared without the need for further surgical intervention, and the cats were monitored for any complications and none were recorded. All cases recovered uneventfully without recurrence for a period of six months.



Fig 4: Normal urine flow after relieving the obstruction, note hematuria



Fig 5: Pasty material at penile orifice

4. Discussion

This study demonstrated a high success rate (96.7%) of clearing urethral obstruction in male cats using hydrogen peroxide and saline flushing, which is consistent with Xu *et al.* (2020) [5]. Murphy and Friedman (2019) ascribed the mechanical disruption of obstructions to hydrogen peroxide's effervescence, which aligns with our observations of relieving obstruction. Pachtinger (2020) [3] recommended the use of saline to aid in the flushing and removal of debris, which was corroborated by our results. The presence of pasty material at the urethral opening observed in 22 cases is consistent with Pachtinger (2020) [3], who informed that mucus plugs are a significant contributor to urethral obstruction in male cats. This pasty material, likely a combination of mucus and cellular debris, is known to block the urethra and can be effectively dislodged by the effervescence created by hydrogen peroxide (Murphy and Friedman, 2019). Despite causing mild irritation to mucous membranes, hydrogen peroxide facilitates the breakdown of obstructions through its effervescent property as observed in this study. The success rate observed in this study is promising and suggests that hydrogen peroxide and saline flushing could be a viable, minimally invasive alternative to traditional methods. Future studies with larger sample sizes are needed to confirm these findings and explore the long-term outcomes of this technique.

5. Conclusion

Use of 3% hydrogen peroxide and normal saline (1:1), flushing via penile catheterization offers a safe, effective, and minimally invasive approach to manage urethral obstruction in male cats. With a high success rate and minimal complications, this technique is a valuable alternative to more invasive surgical interventions. Further studies with larger sample sizes are necessary to validate these findings and explore the long-term outcomes of this treatment.

Conflict of Interest

Not available

Financial Support

Not available

6. References

1. George CM, Grauer GF. Feline urethral obstruction: Diagnosis and management. *Veterinary Clinics of North America: Small Animal Practice*. 2017;47(6):1101-1123.
2. Murphy EC, Friedman AJ. Hydrogen peroxide and cutaneous biology: Translational applications, benefits and risks. *Journal of the American Academy of Dermatology*. 2019;81(6):1379-1386.
3. Pachtinger G. Urinary catheter placement for feline urethral obstruction. *Journal of Feline Medicine and Surgery*. 2020;22(9):772-781.
4. Segev G, Livne H, Ranen E, Lavy E. Urethral obstruction in cats: predisposing factors, clinical, clinicopathological characteristics and prognosis. *Journal of Feline Medicine and Surgery*. 2011;13(2):101-108.
5. Xu M, Jin L, Shan Y, Zhu J, Xue B. A simple and effective method for bladder blood clot evacuation using hydrogen peroxide. *Journal of International Medical Research*. 2020;48(5):1-7.

How to Cite This Article

Nallareddy K, Sadu RR, Tellapati JP. Use of hydrogen peroxide and normal saline flushing in clearing feline urethral obstruction: A retrospective study of 30 cases. *International Journal of Veterinary Sciences and Animal Husbandry*. 2025;10(4):148-150.

Creative Commons (CC) License

This is an open-access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.