



International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
NAAS Rating: 5.29
IJABR 2025; 9(4): 1004-1007
www.biochemjournal.com
Received: 12-01-2025
Accepted: 16-02-2025

Prachi Jain

M.Sc. (Ag.), Department of
Genetics & Plant Breeding,
S.V. Agricultural College,
Tirupati, Andhra Pradesh,
India

M Sreevalli Devi

Scientist (Genetics & Plant
Breeding), Agricultural
Research Station,
Perumallapalle, Andhra
Pradesh, India

K John

Principal Scientist (Genetics &
Plant breeding), Regional
Agricultural Research Station,
Tirupati, Andhra Pradesh,
India

P Arunasri

Assistant Professor (Plant
Pathology), S.V. Agricultural
College, Tirupati, Andhra
Pradesh, India

Selection indices for yield, yield components and resistance to late leaf spot in groundnut (*Arachis hypogaea* L.)

Prachi Jain, M Sreevalli Devi, K John and P Arunasri

DOI: <https://www.doi.org/10.33545/26174693.2025.v9.i4l.4218>

Abstract

Thirty one diverse genotypes of groundnut were evaluated in a randomized block design with three replications for the study of selection indices during *kharif* 2023. Sixty three selection indices involving pod yield/plant (X_1) and five yield components viz., number of primary branches/plant (X_2), dry haulm yield/plant (X_3), kernel yield/plant (X_4), late leaf spot at 75 DAS (X_5) and late leaf spot at 90 DAS (X_6) were constructed using the discriminant function technique. Discriminant function analysis indicated that selection efficiency of the function was improved by increasing the number of characters in the index. Among the single character indices, dry haulm yield/plant exhibited higher genetic advance and relative efficiency over straight selection for pod yield/plant. The index based on six characters viz., pod yield/plant (X_1) + number of primary branches/plant (X_2) + dry haulm yield/plant (X_3) + kernel yield/plant (X_4) + late leaf spot at 75 DAS (X_5) and late leaf spot at 90 DAS (X_6) recorded the highest genetic advance as well as relative efficiency and selection efficiency. However, in practice, the plant breeder might be interested in maximum gain with minimum number of characters. In the present study, improvement of pod yield in groundnut could be achieved by selecting the parents with these four characters; number of primary branches/plant + dry haulm yield/plant + late leaf spot at 75 DAS + late leaf spot at 90 DAS which is more desirable and practically possible to use by breeder than the index that includes more number of characters.

Keywords: Discriminant function, relative efficiency, selection indices and groundnut

Introduction

Groundnut (*Arachis hypogaea* L.) is one of the most economic oilseed crops of the world. It is considered as the world's fourth largest source of edible oil and third most important source of vegetable protein (Desai *et al.*, 1999) [4]. Yield in crops is a quantitative trait and has a complex genetic control mechanism and hence, direct selection is not much effective on it. Since the economic part of groundnut known as pod is developed under the soil, the prediction of its performance based on aerial morphological characters is almost difficult (Weiss, 2000) [15]. The most desirable approach to improve characteristics such as pod yield is simultaneous selection based on related traits (Bos and Caligri, 2007) [1]. This can be done using selection index, which is multiple regressions of genotypic values on phenotypic values of several traits (Falconer, 1996) [6]. The use of selection index is superior in improving complex traits (Hasel and Lush, 1942) [9]. Furthermore, the selection indices approach aimed at determining the most suitable combination of traits with the intention of indirectly improving the pod yield in groundnut was well documented (Dobariya *et al.*, 2008) [5].

Certain desired plant characteristics are considered while selecting for particular genotype with varying weightage given to different traits for arriving on decisions. The better way of exploiting genetic correlations with several traits having high heritability is to construct an index which combines information on all the characters associated with yield. This suggests the use of selection index, which gives proper weight to each of the two or more characters to be considered. Selection index was proposed for the first time by Smith (1936) [14] on the basis discriminant function of Fisher (1936) [7]. Hazel and Lush (1943) [10] and Robinson *et al.* (1951) [13] showed that the selection based on such an index is more efficient than selecting individually for the various characters. Few studies on selection indices in

Corresponding Author:**Prachi Jain**

M.Sc. (Ag.), Department of
Genetics & Plant Breeding,
S.V. Agricultural College,
Tirupati, Andhra Pradesh,
India

groundnut have been carried out earlier by Dobariya *et al.* (2008) ^[5]; Babariya *et al.* (2014) ^[2] and Gupta *et al.* (2015) ^[8]. However, in order to have a more comprehensive knowledge about genetic variability for yield and its attributing traits, it is important to find out a suitable selection indices for the improvement of pod yield in groundnut. Keeping these facts in view, the present study was undertaken in order to construct selection indices for efficient selection in groundnut breeding programme.

Materials and Methods

Thirty one genotypes of groundnut were sown in a Randomized Block Design (RBD) with three replications during *kharif* 2023. Each genotype was sown in 2 rows of 3 m length with a spacing of 30 cm between rows and 10 cm between plants within the row. The crop was raised under irrigated conditions and recommended dose of chemical fertilizers at the rate of 20 kg of N, 40 kg of P₂O₅ and 50 kg K₂O ha⁻¹ in the form of urea, single super phosphate, murate of potash, respectively were applied during final ploughing operation and gypsum was added at the rate of 500 kg ha⁻¹ at peak flowering stage followed by one hand weeding to remove weeds and to mix the gypsum properly in the soil. Apart from these cultural operations, need based plant protection measures were taken to control insect pests. Observations were recorded for all the genotypes separately on randomly chosen five competitive plants in each genotype in each replication for all the characters except for days to 50% flowering and days to maturity which were recorded on plot basis. Data were recorded for 15 characters *viz.*, days to 50% flowering, days to maturity, plant height at harvest, number of primary branches/plant, number of secondary branches/plant, number of mature pods/plant, number of immature pods/plant, pod yield/plant, dry haulm yield/plant, harvest index percent, shelling percent, kernel yield/plant, sound mature kernel percent, hundred kernel weight, disease scoring of late leaf spot disease at 75 DAS and 90 DAS.

Discriminant function analysis described by Dabholkar (1992) ^[3] was used to construct the selection indices involving six characters, pod yield/plant (X₁), number of primary branches/plant (X₂), 100-kernel weight (X₃), biological yield/plant (X₄), harvest index (X₅) and days to maturity (X₆). For computing selection indices, pod yield/plant was considered as the dependent variable with the relative efficiency of 100%. The model suggested by Robinson *et al.* (1951) ^[13] was used for the construction of genetic advance as well as selection indices and development of a required discriminant function using six characters along with pod yield/plant.

Results and Discussion

High genetic advance coupled with high heritability estimates offers the most suitable condition for selection which indicates the presence of additive genes in the trait and further suggest reliable crop improvement through selection of traits (Ogunniyan and Olakojo, 2014) ^[11]. Therefore, total six characters out of fifteen characters under study with high genetic advance coupled with high heritability were used for construction of selection indices. A total of sixty three selection indices (Table 1) based on six

characters constructed in all possible combinations revealed that the selection efficiency was high over straight selection when selection was based on individual components. Dry haulm yield/plant (g) showed a genetic advance of 23.17%, which was higher than those calculated for other characters including pod yield/plant suggested that dry haulm yield/plant (g) proved to be better selection index based on one character.

The highest genetic gain (Table 1) of 34.45% was obtained when selection was simultaneously based on discriminant function of two characters, e.g. pod yield/plant (X₁) and dry haulm yield/plant (X₃). When three characters, e.g. number of primary branches/plant (X₂), dry haulm yield/plant (X₃) and kernel yield/plant (X₄) were taken together, the genetic advance increased to 38.14%. Index based on combination of four characters, *i.e.* number of primary branches/plant (X₂), dry haulm yield/plant (X₃), late leaf spot at 75 DAS (X₅) and late leaf spot at 90 DAS (X₆) recorded high genetic gain of 46.82%. Index based on combination of five characters, *i.e.* number of primary branches/plant (X₂), dry haulm yield/plant (X₃), kernel yield/plant (X₄), late leaf spot at 75 DAS (X₅) and late leaf spot at 90 DAS (X₆) recorded high genetic gain of 49.01%. The maximum gain of 52.04% was achieved by taking six characters at a time, *i.e.* pod yield/plant (X₁), number of primary branches/plant (X₂), dry haulm yield/plant (X₃), kernel yield/plant (X₄), late leaf spot at 75 DAS (X₅) and late leaf spot at 90 DAS (X₆).

Thus, the current study revealed that the index which includes more than one character, gave high genetic advance, suggesting the utility of constructing of selection indices for effecting simultaneous improvement in several characters. Hazel and Lush (1943) ^[10] stated that the superiority of selection based on index increases with an increase in the number of characters under selection. Smith (1936) ^[14], Rao (1974) ^[12], Dobariya *et al.* (2008) ^[5], Babariya *et al.* (2014) ^[2] and Gupta *et al.* (2015) ^[8] also with same opinion that inclusion of characters one by one in the function resulted in increasing genetic advance and that the selection indices improve the efficiency than the straight selection for yield alone.

The relative efficiency (RE%) of various selection indices presented in Table 1 indicated that when relative efficiency of single character index was measured over straight selection for pod yield/plant, the efficiency was declined to less than 100%. This observation indicated that the indirect selection through individual traits over straight selection for pod yield/plant alone would not be effective. Therefore, an increase in number of characters in combination with yield would be effective.

It is interesting to note that selection efficiency (Table 2) improved with an increase in number of characters in combination with yield. For example, average selection efficiency of 108.74%, when one character was included in selection function. Similarly, the selection efficiency was 236.27% for two characters, 341.16% for three characters, 458.11% for four characters, 488.40% for five characters and 532.01% for six characters selection indices improve the selection efficiency than the straight selection for yield alone with an increase in the number of characters under selection.

Table 1: Selection index, discriminant function, expected genetic advance in yield and relative efficiency from the use of different selection indices in groundnut

S. No.	Selection Index	Discriminant Function	Expected Genetic Advance (%)	Relative Efficiency (%)
1	X ₁ Pod yield/plant (g)	0.7018 X ₁	9.22	100
2	X ₂ Number of primary branches/plant	0.8657 X ₂	6.12	80.71
3	X ₃ Dry haulm yield/plant (g)	0.9107 X ₃	23.17	204.12
4	X ₄ Kernel yield/plant (g)	0.6379 X ₄	8.17	93.12
5	X ₅ Late leaf spot at 75 DAS	0.8306 X ₅	6.42	87.83
6	X ₆ Late leaf spot at 90 DAS	0.8428 X ₆	7.93	86.63
7	X ₁ .X ₂	0.6976 X ₁ + 1.1436 X ₂	16.78	190.02
8	X ₁ .X ₃	0.6984 X ₁ + 0.9975 X ₃	34.45	360.92
9	X ₁ .X ₄	0.7521 X ₁ + 0.8637 X ₄	17.14	197.91
10	X ₁ .X ₅	0.6838 X ₁ + 0.6833 X ₅	15.12	184.03
11	X ₁ .X ₆	0.6812 X ₁ + 0.7350 X ₆	15.95	185.42
12	X ₂ .X ₃	0.9468 X ₂ + 0.9125 X ₃	30.15	331.84
13	X ₂ .X ₄	0.9729 X ₂ + 0.6301 X ₄	14.12	180.09
14	X ₂ .X ₅	0.8207 X ₂ + 0.8015 X ₅	12.04	175.84
15	X ₂ .X ₆	0.7565 X ₂ + 0.7876 X ₆	14.43	181.71
16	X ₃ .X ₄	0.9446 X ₃ + 0.6441 X ₄	33.43	349.04
17	X ₃ .X ₅	0.9111 X ₃ + 0.8711 X ₅	26.21	316
18	X ₃ .X ₆	0.9046 X ₃ + 0.8118 X ₆	30.62	336.71
19	X ₄ .X ₅	0.6021 X ₄ + 0.6782 X ₅	15.63	185.12
20	X ₄ .X ₆	0.6306 X ₄ + 0.7777 X ₆	16.23	188.12
21	X ₅ .X ₆	0.9096 X ₅ + 0.8467 X ₆	14.23	181.23
22	X ₁ .X ₂ .X ₃	0.6940 X ₁ + 1.1691 X ₂ + 0.9957 X ₃	36.12	375.04
23	X ₁ .X ₂ .X ₄	0.7591 X ₁ + 1.1217 X ₂ + 0.8356 X ₄	30.12	331.01
24	X ₁ .X ₂ .X ₅	0.6843 X ₁ + 1.0815 X ₂ + 0.7141 X ₅	26.73	316.34
25	X ₁ .X ₂ .X ₆	0.6852 X ₁ + 1.0189 X ₂ + 0.7333 X ₆	27.23	320.42
26	X ₁ .X ₃ .X ₄	0.7705 X ₁ + 0.9913 X ₃ + 0.8159 X ₄	37.15	381.14
27	X ₁ .X ₃ .X ₅	0.6799 X ₁ + 1.0008 X ₃ + 0.7937 X ₅	34.12	358.65
28	X ₁ .X ₃ .X ₆	0.6812 X ₁ + 0.9942 X ₃ + 0.7574 X ₆	33.23	348.94
29	X ₁ .X ₄ .X ₅	0.7450 X ₁ + 0.8303 X ₄ + 0.6527 X ₅	27.13	320.03
30	X ₁ .X ₄ .X ₆	0.7285 X ₁ + 0.8742 X ₄ + 0.7607 X ₆	30.14	331.42
31	X ₁ .X ₅ .X ₆	0.6677 X ₁ + 0.7930 X ₅ + 0.8177 X ₆	26.94	316.89
32	X ₂ .X ₃ .X ₄	1.0306 X ₂ + 0.9462 X ₃ + 0.6296 X ₄	38.14	380.04
33	X ₂ .X ₃ .X ₅	0.9214 X ₂ + 0.9133 X ₃ + 0.8667 X ₅	35.12	372.81
34	X ₂ .X ₃ .X ₆	0.8352 X ₂ + 0.9073 X ₃ + 0.7741 X ₆	36.03	374.02
35	X ₂ .X ₄ .X ₅	0.8988 X ₂ + 0.5994 X ₄ + 0.6657 X ₅	26.42	316.22
36	X ₂ .X ₄ .X ₆	0.8425 X ₂ + 0.6283 X ₄ + 0.7391 X ₆	23.43	304.91
37	X ₂ .X ₅ .X ₆	0.7451 X ₂ + 1.0056 X ₅ + 0.7230 X ₆	24.52	310.62
38	X ₃ .X ₄ .X ₅	0.9437 X ₃ + 0.6204 X ₄ + 0.7533 X ₅	32.64	342.43
39	X ₃ .X ₄ .X ₆	0.9371 X ₃ + 0.6416 X ₄ + 0.7699 X ₆	34.52	365.07
40	X ₃ .X ₅ .X ₆	0.9080 X ₃ + 1.2573 X ₅ + 0.5777 X ₆	31.83	340.83
41	X ₄ .X ₅ .X ₆	0.6215 X ₄ + 0.5957 X ₅ + 0.9952 X ₆	26.73	316.34
42	X ₁ .X ₂ .X ₃ .X ₄	0.7770 X ₁ + 1.1559 X ₂ + 0.9899 X ₃ + 0.7861 X ₄	44.23	456.72
43	X ₁ .X ₂ .X ₃ .X ₅	0.6766 X ₁ + 1.1446 X ₂ + 1.0006 X ₃ + 0.8382 X ₅	42.61	421.51
44	X ₁ .X ₂ .X ₃ .X ₆	0.6838 X ₁ + 1.0580 X ₂ + 0.9930 X ₃ + 0.7627 X ₆	43.01	436.41
45	X ₁ .X ₂ .X ₄ .X ₅	0.7548 X ₁ + 1.0497 X ₂ + 0.8087 X ₄ + 0.6673 X ₅	39.01	382.94
46	X ₁ .X ₂ .X ₄ .X ₆	0.7420 X ₁ + 1.0014 X ₂ + 0.8495 X ₄ + 0.7481 X ₆	39.84	396.61
47	X ₁ .X ₂ .X ₅ .X ₆	0.6720 X ₁ + 1.0178 X ₂ + 0.8855 X ₅ + 0.7519 X ₆	41.43	410.03
48	X ₁ .X ₃ .X ₄ .X ₅	0.7578 X ₁ + 0.9930 X ₃ + 0.8000 X ₄ + 0.7451 X ₅	45.92	502.31
49	X ₁ .X ₃ .X ₄ .X ₆	0.7492 X ₁ + 0.9883 X ₃ + 0.8278 X ₄ + 0.7662 X ₆	45.95	505.01
50	X ₁ .X ₃ .X ₅ .X ₆	0.6611 X ₁ + 1.0032 X ₃ + 1.1870 X ₅ + 0.5726 X ₆	45.01	481.03
51	X ₁ .X ₄ .X ₅ .X ₆	0.7127 X ₁ + 0.8795 X ₄ + 0.5942 X ₅ + 0.9821 X ₆	39.71	385.03
52	X ₂ .X ₃ .X ₄ .X ₅	0.9801 X ₂ + 0.9460 X ₃ + 0.6108 X ₄ + 0.7578 X ₅	46.12	516.01
53	X ₂ .X ₃ .X ₄ .X ₆	0.9044 X ₂ + 0.9396 X ₃ + 0.6335 X ₄ + 0.7433 X ₆	46.74	516.45
54	X ₂ .X ₃ .X ₅ .X ₆	0.8204 X ₂ + 0.9113 X ₃ + 1.3538 X ₅ + 0.4708 X ₆	46.82	516.94
55	X ₂ .X ₄ .X ₅ .X ₆	0.8342 X ₂ + 0.6224 X ₄ + 0.6909 X ₅ + 0.8903 X ₆	43.14	444.23
56	X ₃ .X ₄ .X ₅ .X ₆	0.9392 X ₃ + 0.6417 X ₄ + 0.9675 X ₅ + 0.7352 X ₆	45.72	500.42
57	X ₁ .X ₂ .X ₃ .X ₄ .X ₅	0.7643 X ₁ + 1.1130 X ₂ + 0.9926 X ₃ + 0.7777 X ₄ + 0.7729 X ₅	44.11	450.04
58	X ₁ .X ₂ .X ₃ .X ₄ .X ₆	0.7615 X ₁ + 1.0410 X ₂ + 0.9869 X ₃ + 0.8028 X ₄ + 0.7605 X ₆	45.21	480.43
59	X ₁ .X ₂ .X ₃ .X ₅ .X ₆	0.6638 X ₁ + 1.0557 X ₂ + 1.0025 X ₃ + 1.2784 X ₅ + 0.5146 X ₆	44.62	460.83
60	X ₁ .X ₂ .X ₄ .X ₅ .X ₆	0.7252 X ₁ + 1.0014 X ₂ + 0.8581 X ₄ + 0.6787 X ₅ + 0.9122 X ₆	45.24	495.83
61	X ₁ .X ₃ .X ₄ .X ₅ .X ₆	0.7225 X ₁ + 0.9964 X ₃ + 0.8479 X ₄ + 0.9731 X ₅ + 0.7346 X ₆	48.52	520.13
62	X ₂ .X ₃ .X ₄ .X ₅ .X ₆	0.8908 X ₂ + 0.9420 X ₃ + 0.6369 X ₄ + 1.0607 X ₅ + 0.6427 X ₆	49.01	523.12
63	X ₁ .X ₂ .X ₃ .X ₄ .X ₅ .X ₆	0.7337 X ₁ + 1.0393 X ₂ + 0.9954 X ₃ + 0.8261 X ₄ + 1.0563 X ₅ + 0.6723 X ₆	52.04	532.01

Table 2: Average selection efficiency of different combination of characters in groundnut

No. of characters in the index	Selection efficiency (%)
One	108.74
Two	236.27
Three	341.16
Four	458.11
Five	488.4
Six	532.01

Table 3: Highest selection efficiency with character combinations in groundnut

S. No.	Character	Selection Efficiency (%)
1	Dry haulm yield/plant	204.12
2	Pod yield/plant + Dry haulm yield/plant	360.92
3	Pod yield/plant + Dry haulm yield/plant + Kernel yield/plant	381.14
4	Number of primary branches/plant + Dry haulm yield/plant + Kernel yield/plant	380.04
5	Number of primary branches/plant + Dry haulm yield/plant + Late leaf spot at 75 DAS + Late leaf spot at 90 DAS	516.94
6	Number of primary branches/plant + Dry haulm yield/plant + Kernel yield/plant + Late leaf spot at 75 DAS + Late leaf spot at 90 DAS	523.12
7	Pod yield/plant + Number of primary branches/plant + Dry haulm yield/plant + Kernel yield/plant + Late leaf spot at 75 DAS + Late leaf spot at 90 DAS	532.01

Some of the selection indices with high relative efficiency listed in Table 1 indicated that the highest efficiency was observed with a combination of six characters (532.01%). Selection indices with six characters, i.e. pod yield/plant (X_1), number of primary branches/plant (X_2), dry haulm yield/plant (X_3), kernel yield/plant (X_4), late leaf spot at 75 DAS (X_5) and late leaf spot at 90 DAS (X_6), therefore, appear to be more useful. It can be seen that number of primary branches/plant (X_2), dry haulm yield/plant (X_3) and kernel yield/plant (X_4) were the characters being commonly involved in more number of the combinations, the next being pod yield/plant (X_1), late leaf spot at 75 DAS (X_5) and late leaf spot at 90 DAS (X_6) in order (Table 3).

Keeping in view, the basic idea of saving time and labour in a selection programme, it would be desirable to base the selection of few characters. In the present study, selection index based on six characters gave maximum genetic gain and high efficiency over straight selection, but practically it is more cumbersome to use in the selection exercise. However, in practice, the plant breeder might be interested in maximum gain with minimum number of characters. In the present study, selection index based on four characters (number of primary branches/plant + dry haulm yield/plant + late leaf spot at 75 DAS + late leaf spot at 90 DAS) showing genetic gain (46.82%) and selection efficiency (516.94%) comparable to some extent of those based on five or more characters, which is more desirable and practically possible to use breeder than the index that includes more number of characters.

Conclusion

From this investigation, it is concluded that improvement of pod yield in groundnut could be achieved by selecting the parents with these four characters; number of primary branches/plant + dry haulm yield/plant + late leaf spot at 75 DAS + late leaf spot at 90 DAS.

References

- Bos I, Caligari P. Selection methods in plant breeding. Springer Science & Business Media; 2007.
- Babariya CA, Dobariya KL, Sapovadiya MH, Vavdiya PA. Discriminant function method of selection in groundnut (*Arachis hypogaea* L.). Electronic Journal of Plant Breeding. 2014;5(1):112-116.
- Dabholkar AR. Elements of Biometrical Genetics. Concept Publishing Company, New Delhi; 1992. p. 444-478.
- Desai BB, Kotecha PM, Salunkhe DK. Science and Technology of Groundnut: Biology, Production, Processing and Utilization. Naya Prokash, Calcutta; 1999. p. 2-4.
- Dobariya KL, Odedra RK, Jivani LL, Ponkia HP, Khanpara MD. Selection indices for yield improvement in groundnut (*Arachis hypogaea* L.). International Journal of Bioscience Reporter. 2008;6(2):299-330.
- Falconer DS. Introduction to Quantitative Genetics. Pearson Education India; 1996.
- Fisher RA. The use of multiple measurements in taxonomic problems. Annals of Eugenics. 1936;7(2):179-188.
- Gupta RP, Vachhani JH, Kachhadia VH, Vaddoria MA, Praveen Kumar. Selection indices in Virginia groundnut (*Arachis hypogaea* L.). Electronic Journal of Plant Breeding. 2015;6(2):560-565.
- Hazel LN, Lush JL. The efficiency of three methods of selection. Journal of Heredity. 1942;33(11):393-399.
- Hazel LN, Lush JL. The genetic basis for constructing selection indexes. Genetics. 1943;28(6):476-490.
- Ogunniyan DJ, Olakojo SA. Genetic variation, heritability, genetic advance and agronomic character association of yellow elite inbred lines of maize (*Zea mays* L.). Nigerian Journal of Genetics. 2014;28:24-28.
- Rao L. Analysis of genetic variability and formulation of selection indices for yield in soybean (*Glycine max* (L.) Merrill). The Mysore Journal of Agricultural Sciences. 1974;8(1):156.
- Robinson HF, Comstock RE, Harvey VH. Genotypic and phenotypic correlations in corn and their implications in selection. Agronomy Journal. 1951;41:353-359.
- Smith HF. A discriminant function for plant selection. Annals of Eugenics. 1936;7:240-250.
- Weiss EA. Oilseed Crops. 2nd ed. Blackwell Science Ltd., London; 2000. p. 53-59.