


International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
NAAS Rating: 5.29
IJABR 2025; 9(5): 31-35
www.biochemjournal.com
Received: 05-03-2025
Accepted: 08-04-2025

Ankita Chib

Division of Post Harvest and Management, SKUAST-Jammu, Chatha, Jammu and Kashmir, India

Jagmohan Singh

Division of Post Harvest and Management, SKUAST-Jammu, Chatha, Jammu and Kashmir, India

Anju Bhat

Division of Post Harvest and Management, SKUAST-Jammu, Chatha, Jammu and Kashmir, India

Shivani Verma

Division of Post Harvest and Management, SKUAST-Jammu, Chatha, Jammu and Kashmir, India

Corresponding Author:**Ankita Chib**

Division of Post Harvest and Management, SKUAST-Jammu, Chatha, Jammu and Kashmir, India

Development and quality evaluation of Harad-Ber blended leather

Ankita Chib, Jagmohan Singh, Anju Bhat and Shivani Verma

DOI: <https://www.doi.org/10.33545/26174693.2025.v9.i5a.4291>

Abstract

The present investigation was carried out to develop blended fruit leather using Harad and Ber in six different combination T₁ (100:0::HP:BP), T₂ (70:30::HP:BP), T₃ (60:40::HP:BP), T₄ (50:50::HP:BP), T₅ (40:60::HP:BP), and T₆ (30:70::HP:BP) and evaluated for proximate composition during storage at room temperature. Physicochemical attributes (moisture, crude protein, crude fibre, fat, and ash) were analysed during six months of storage at an interval of 0,2,4,6 months. Results indicated that treatment T₁ (100:0 HP:BP) exhibited the highest crude protein (3.48 %), crude fibre (5.15%), crude fat (1.72%), and ash content (4.07%) at initial day. Whereas, the treatment T₆ (70:30 HP:BP) recorded the lowest values. A significant decline ($p \leq 0.05$) in all nutritional parameters was observed over the analysis attributed to enzymatic activity, moisture loss, and Maillard reactions. During storage there was decreased in moisture content from 10.49% to 9.33%, crude protein 2.88% to 2.48%, crude fibre 4.45 to 4.22, crude fat 1.19 to 0.98 and ash content from 3.37 to 3.08% respectively. Overall blending and storage had a significant impact on the nutritional profile of leather.

Keywords: Terminalia chebula, Zizyphus mauritiana, fruit leather, storage, proximate composition, nutritional quality

1. Introduction

Medicinal plants have been a valuable source of both curative and preventive medical therapeutic preparations for humans, as well as for the extraction of key bioactive substances (Thirumalai *et al.*, 2009) [37]. Nearly 80% of people worldwide rely on traditional medicine and goods for their medical requirements, particularly in third-world nations. *Terminalia chebula*, also known as Harad or Chebulic Myrobalan, is one of the most highly valued medicinal herbs and wide range of phytoconstituents. Harad fruits are known as drupes, and they are 2.5-4 cm long, glabrous to ellipsoidal or obovoid in shape with a broad base, and a yellowish green more or less 5-ribbed shape when dried. It is mostly produced in the northwestern Indian states of Himachal Pradesh, Jammu and Kashmir, Punjab, Haryana, Uttar Pradesh, and Uttarakhand (Sankanur *et al.*, 2015) [27]. It is highly regarded as the 'King of medicines' in the Ayurvedic Medicines. It is very nutritious containing essential vitamins, minerals, and proteins. It contains potassium, iron, copper, manganese, selenian, and vitamin C. It is used to treat a wide range of symptoms, including neurological problems, hepatomegaly, diabetes, rheumatism, tumours, intermittent fever, paralysis, memory loss, and constipation (Gupta, 2012) [12].

One of India's oldest fruits is ber (*Zizyphus mauritiana* Lamk.). Ber, a member of the Rhamnaceae family, is known as the "king of arid zone fruits". In India, ber has been known as a beneficial edible fruit from ancient times (Take and Bhotmange, 2012) [36]. It is widely cultivated in different states of India, i.e. Madhya Pradesh, Rajasthan, Gujarat, Punjab, Haryana, Uttar Pradesh, Maharashtra and to limited extent in several other states (Samant *et al.*, 2008) [26]. Although Indian jujube is grown primarily in India, it is found throughout the world's warm climates. It contains carbohydrates, moisture, protein, fat and fiber. This fruit also contains minor elements including magnesium, copper, iron, zinc, and sodium (Sareen *et al.*, 2020) [28].

Demand for nutritious and safe food is rising around the world because eating a balanced diet is the recommended approach to avoid health issues like obesity, diabetes, and nutritional deficiencies, which are mostly caused by dietary mistakes.

The current trend of eating nutritious, inventive and healthful cuisine has resulted in a steady increase in the fruit bar market (Saleem *et al.*, 2023) ^[25]. Thus, there is a need for diversity in commercial utilization of fruits. Fruits can be utilized in numerous ways by processing into juice, squashes, jams, concentrates, pulp, dehydrated products, jellies and fruit leather. Fruit leathers are restructured fruits created from fresh fruit pulp or a mixture of fruit juice concentrates and other components following an innovative dehydration procedure (Maskan *et al.* 2002) ^[15]. Different drying methods used in processing fruit leather. Therefore, the current research was designed to developed harad fruit leather in combination with ber fruit for preservation of nutrients and analysis its physico-chemical properties of harad-ber blended leather.

2. Materials and Methods

Blended leathers were prepared using the Chavan *et al.* (2016) ^[39] method. The Harad pulp was blended with ber pulp in different proportions T₁ (100:00::HP:BP), T₂ (70:30::HP:BP), T₃ (60:40::HP:BP), T₄ (50:50::HP:BP), T₅ (40:60::AP:BP) and T₆ (30:70::HP:BP) for the preparation of leather. Blended pulp was heated along with desired quantities of sugar and citric acid with constant stirring on a uniform flame. The pulp or puree was spread on oil-smeared stainless steel trays. The pulp/puree was dried at 50 °C for 7-10 hrs in cabinet tray drier. After that, the semi surface dried pulp sheets were cut into desired size usually of rectangular shape and again dried for 4-5 hrs. Then desired cutting was done and dried under fan for 2-3 hrs. The blended leather was stored in laminated pouches under ambient conditions (30-35 °C) for a period of six months. The samples were analyzed for physico-chemical parameters during six months of storage at an interval of 0, 2, 4 and 6 months. The moisture content was estimated using the AOAC method (2012) ^[3]. Kjeldahl method was used to determined crude protein using Sadasivam and Manickam, (2008) ^[24] methodology. Crude fibre was estimated by AOAC (2005) ^[4] Ash content was measured using a weighing balance (GM 1501), an analytical balance (Shimadzu ATX 224), and an electric muffle furnace (Thermolyne 1400 Furnace), as reported by AACC (2000) ^[1]. Crude fat was determined by the Soxhlet extraction technique (AOAC, 2012) ^[3]. Factorial completely randomized design (FCRD) was used to statistically analyze the data (Gomez & Gomez, 1984) ^[11].

3. Results and Discussion

Table 1: Physico-chemical composition of raw materials

Characteristics	Harad Pulp	Ber Pulp
Moisture (%)	59.62	81.27
Titratable acidity (%)	3.10	0.41
Crude protein (%)	2.62	1.21
Crude fibre (%)	4.36	2.31
Fat (%)	1.59	0.16
Ash (%)	3.22	1.60

3.1 Proximate composition

The data pertaining to the physical properties of the raw materials in Table 1 revealed that the moisture and titratable acidity in fresh harad pulp were represented as 59.62% and 3.10 percent which were in close compliance to the findings of Navhale *et al.*, (2011) ^[18] and Bhutia *et al.*, (2018) ^[7]. The

crude protein, crude fibre, fat and ash in fresh harad pulp were found to be 2.62%, 4.36%, 1.59% and 3.22%, respectively which concurred with the outcomes of Singh *et al.*, (2017) ^[32].

The moisture content, and titratable acidity in fresh ber pulp were recorded as 81.27% and 0.41%, respectively which coincide by Pandey *et al.*, (2019) ^[19] and Kavitha *et al.*, (2014) ^[14]. The crude protein, crude fibre, fat and ash were found to be 1.21%, 2.31%, 0.16% and 1.60% which are in accordance with the findings of Dewan (2021) ^[9], Sharma and Rana (2017) ^[30], Sharif *et al.*, (2022) ^[29] and Pareek (2013) ^[20] in fresh ber pulp.

3.2 Moisture Content

The mean moisture content of blended leather (Table 2) ranged from 9.19 to 10.41% and varied significantly with varying levels of harad pulp and ber pulp. The highest mean moisture content of 10.41 was observed in treatment T₆ (30:70:: HP : BP) whereas the lowest mean moisture content of 9.19 was recorded in treatment T₁ (100:00:: HP : BP). The decrease in moisture content might be due to evaporation of water from leather during storage (Bhatt and Jha, 2015) ^[6]. Kaushal *et al.* (2017) ^[13] also reported decrease in moisture content during storage in ginger plum leather. Ahmed *et al.* (2021) ^[2] observed similar results, indicating a decrease in moisture content of pear and grapes blended leather after three months of storage.

3.3 Crude protein

It was apparent from Table 3 that treatment exerted a considerable effect on the crude protein of blended leather. The highest and lowest treatment mean crude protein of 3.26% and 2.24% were found in treatment T₁ (100:00::HP:BP) and T₆ (70:30::HP:BP), respectively. There was a significant decrease in the mean value of blended leather from initial value of 2.88 to 2.49%, during 6 months of storage duration. The decrease in protein content during storage might be due to hydrolysis of peptide bond by the help of protease enzyme that causes splitting of protein molecules and denaturation and degradation of protein into amino acid during storage. The crude protein of the blended leather decreases with the advent of storage period which are quite similar with the findings of Kaushal *et al.* (2015) ^[33] and Patil *et al.* (2017) ^[22] in leather variant prepared from the seabuck leather and peach-soy fruit leather, respectively.

3.4 Crude fibre

The maximum and minimum mean value of crude fibre (Table 4) of 5.01 and 3.72 were recorded in treatment T₁ (100:00::HP:BP) and T₆ (30:70::HP:BP), respectively. Significant decrease in crude fibre was observed from 4.55 to 4.22 during storage period in the blended leather. The decrease in crude fibre might be due to the degradation of hemicelluloses and other structural polysaccharides during storage. Also, heat and moisture solubilizers degraded pectic substances leading to the decrease in the fibre content (Sharon and Usha, 2006) ^[31]. Decrease in crude fibre with storage of leather was also reported by Rahman (2021) ^[23] in the mulberry aloe vera blended leather. Similar decline in crude fibre content was also reported by Ghimire and Ojha (2016) ^[10] trend in papaya-soy fruit leather.

3.5 Crude fat

The maximum and minimum mean crude fat of 1.61 and 0.71% were found in treatment T₁ (100:00::HP:BP) and T₆ (30:70::HP:BP), respectively. There was decrease in crude fat (Table 5) of blended leather from initial mean value 1.91 to 0.98% during six months of storage. The decrease in the crude fat content might be because of enhanced lipolytic activity of enzymes like lipase and lipoxygenase in response to increased relative humidity of storage environment causing conversion of fats into esters and glycerol (Zahra *et al.* 2020) [38]. The finding of Bhat *et al.* (2014) [5], who similarly reported a decrease in the crude fat content of soy peach blending leather.

3.6 Ash

The data pertaining to ash content of different treatments in Table 6 depicted a decrease in ash content. At initial day of storage the highest mean ash content of 3.91% was recorded in treatment T₁(100:00:HP::BP) and lowest by 2.63% in treatment T₆ (70:30:HP::BP). During storage the mean ash content of blended leather decreased significantly from 3.37 to 3.08%. The decrease in ash content of blended leather might be attributed to mineral losses from binding of minerals by Maillard reaction products during storage (Nadarajah and Mahenran, 2015) [17]. Similar decrease in ash content is in line with orange leather fortified with orange peel by Parida, (2016) [21]. These findings were also reported by Das and Das (2025) [8] in Kinnow Mandarin fruit leather.

Table 2: Effect of treatments and storage period on moisture content (%) of harad and ber blended leather

Treatments	Storage period(months)				
	0	2	4	6	Mean
T ₁ (100:0 :: HP: BP)	9.80	9.36	8.90	8.69	9.19
T ₂ (70:30:: HP: BP)	10.20	9.68	9.29	8.97	9.54
T ₃ (60:40:: HP: BP)	10.42	9.97	9.62	9.31	9.83
T ₄ (50:50:: HP: BP)	10.65	10.17	9.80	9.52	10.03
T ₅ (40:60:: HP: BP)	10.83	10.39	9.94	9.65	10.20
T ₆ (30:70:: HP: BP)	11.08	10.58	10.13	9.88	10.41
Mean	10.49	10.02	9.61	9.33	
Effects	C.D. ($p \leq 0.05$)				
Treatment (T)	0.03				
Storage (S)	0.03				
T × S	0.06				

Table 3: Effect of treatments and storage period on crude protein (%) of harad and ber blended leather

Treatments	Storage period(months)				
	0	2	4	6	Mean
T ₁ (100:0 :: HP: BP)	3.48	3.31	3.16	3.09	3.26
T ₂ (70:30:: HP: BP)	3.05	2.89	2.77	2.68	2.84
T ₃ (60:40:: HP: BP)	2.90	2.78	2.59	2.49	2.69
T ₄ (50:50:: HP: BP)	2.75	2.60	2.47	2.36	2.54
T ₅ (40:60:: HP: BP)	2.63	2.44	2.35	2.23	2.41
T ₆ (30:70:: HP: BP)	2.49	2.31	2.14	2.05	2.24
Mean	2.88	2.72	2.58	2.49	
Effects	C.D. ($p \leq 0.05$)				
Treatment (T)	0.02				
Storage (S)	0.02				
T × S	0.05				

Table 4: Effect of treatments and storage period on crude fibre (%) of harad and ber blended leather

Treatments	Storage period(months)				
	0	2	4	6	Mean
T ₁ (100:0 :: HP: BP)	5.15	5.03	4.97	4.89	5.01
T ₂ (70:30:: HP: BP)	4.80	4.71	4.65	4.57	4.68
T ₃ (60:40:: HP: BP)	4.52	4.43	4.37	4.29	4.40
T ₄ (50:50:: HP: BP)	4.34	4.25	4.18	4.12	4.22
T ₅ (40:60:: HP: BP)	4.11	4.02	3.95	3.87	3.98
T ₆ (30:70:: HP: BP)	3.82	3.76	3.69	3.62	3.72
Mean	4.45	4.36	4.30	4.22	
Effects	C.D. ($p \leq 0.05$)				
Treatment (T)	0.03				
Storage (S)	0.03				
T × S	0.06				

Table 5: Effect of treatments and storage period on crude fat (%) of harad and ber blended leather

Treatments	Storage period(months)				
	0	2	4	6	Mean
T ₁ (100:0 :: HP: BP)	1.72	1.64	1.57	1.52	1.61
T ₂ (70:30:: HP: BP)	1.39	1.31	1.26	1.15	1.25
T ₃ (60:40:: HP: BP)	1.25	1.18	1.12	1.07	1.16
T ₄ (50:50:: HP: BP)	1.09	1.02	0.95	0.89	0.98
T ₅ (40:60:: HP: BP)	0.93	0.86	0.79	0.71	0.82
T ₆ (30:70:: HP: BP)	0.81	0.74	0.68	0.62	0.71
Mean	1.19	1.12	1.06	0.98	
Effects	C.D. ($p \leq 0.05$)				
Treatment (T)	0.02				
Storage (S)	0.02				
T × S	0.04				

Table 6: Effect of treatments and storage period on ash (%) of harad and ber blended leather

Treatments	Storage period(months)				
	0	2	4	6	Mean
T ₁ (100:0 :: HP: BP)	4.07	3.95	3.84	3.79	3.91
T ₂ (70:30:: HP: BP)	3.67	3.54	3.42	3.28	3.47
T ₃ (60:40:: HP: BP)	3.45	3.32	3.25	3.17	3.29
T ₄ (50:50:: HP: BP)	3.23	3.16	3.07	2.99	3.11
T ₅ (40:60:: HP: BP)	3.01	2.91	2.83	2.75	2.87
T ₆ (30:70:: HP: BP)	2.79	2.67	2.59	2.50	2.63
Mean	3.37	3.25	3.16	3.08	
Effects	C.D. ($p \leq 0.05$)				
Treatment (T)	0.04				
Storage (S)	0.03				
T × S	0.08				

4. Conclusion

In the development of blended fruit leather, varying proportions of harad pulp (HP) and ber pulp (BP) significantly influenced the proximate composition. Treatments with higher harad pulp ratios (T₁: 100:0 HP:BP) exhibited higher levels of moisture, crude protein, fibre, fat, and ash. However, storage for six months resulted in a significant decrease in all parameters across treatments. These reductions can be attributed to factors such as enzymatic degradation, Maillard reactions, moisture loss, and structural polysaccharide breakdown. Overall, blending and storage had a significant impact on the nutritional profile of the leather, with harad pulp contributing more robustly to its nutritional quality.

5. Acknowledgement

The authors are thankful to Division of PHM, SKUAST-Jammu for providing laboratory facility to conduct this study.

6. References

1. AACC. Approved methods of the AACC. 10th ed. St. Paul, MN: American Association of Cereal Chemists; 2000.
2. Ahmad I, Riaz A, Khan A, Shah SS, Shah FN, Zeeshan M. Preparation and evaluation of pear and grapes blended leather. Biological Sciences-PJSIR. 2021;64(2):182-191.
3. AOAC. Official methods of analysis. 19th ed. Gaithersburg, MD: Association of Official Analytical Chemists; 2012.
4. AOAC. Official methods of analysis. 17th ed. Washington, DC: Association of Official Analytical Chemists; 2005.
5. Bhat A, Kumari KR, Anand V, Anjum MA. Preparation, quality evaluation and storage stability of peach-soy fruit leather. SAARC Journal of Agriculture. 2014;12(1):73-88.
6. Bhatt DA, Jha AA. Study of incorporation of therapeutic values of wood apple in fruit bar. International Journal of Pharmaceutical Sciences and Research. 2015;6(10):4398-4405.
7. Bhutia DK, Suresh PC, Pala AN, Shukla G, Chakravarty S. Nutraceutical potential of some wild edible fruits of Sikkim, Himalaya, India. Ethno-Medicine. 2018;12:106-120.
8. Das P, Das A. Process optimization for development of kinnow leather coupled with storage studies. Ecology, Environment & Conservation. 2025;31:405-410.
9. Dewan A. Development and nutritional evaluation of value added products from ber (*Zizyphus mauritiana*). Research and Reviews: Journal of Food and Dairy Technology. 2021;9(2):45-52.
10. Ghimire R, Ojha P. Preparation and physicochemical evaluation of papaya-soy fruit leather. Journal of Science and Technology. 2016;21(1):23-28.
11. Gomez KA, Gomez AA. Statistical procedures for agricultural research. 2nd ed. New York: Wiley-Interscience Publication; 1984.
12. Gupta PC. Biological and pharmacological properties of *Terminalia chebula* Retz. (Haritaki). International Journal of Pharmacy and Pharmaceutical Sciences. 2012;4(3):62-68.
13. Kaushal M, Dhiman A, Vaidya D. Formulation, acceptability and storage stability of appetized ginger plum leather. International Journal of Environment, Agriculture and Biotechnology. 2017;2(1):389-396.
14. Kavitha C, Kuna A, Sagar SB, Supraja T, Padmavathi TVN, Prabhakar BN. Evaluation of physicochemical and antioxidant properties of ber (*Zizyphus mauritiana* Lam.) and RTS ber beverage. Biochemical and Cellular Archives. 2014;14(1):217-222.
15. Maskan A, Kaya S, Maskan M. Effect of concentration and drying processes on color change of grape juice and leather (pestil). Journal of Food Engineering. 2002;54(1):75-80.
16. Meena AK, Rao MM, Sharma K, Yadav A, Singh U. Physicochemical and preliminary phytochemical studies on the fruit of *Terminalia chebula* Retz. Asian Journal of Research in Chemistry. 2010;3(4):844-846.
17. Nadarajah S, Mahendran T. Influence of storage conditions on the quality characteristics of wheat-defatted coconut flour biscuits packed in metalized polypropylene. International Journal of Engineering Research and Technology. 2015;4(7):948-951.
18. Navhale CV, Sonone GN, Jangam SP, Jadhav TS, Bhav GS. Genetic variability and selection of candidate plus trees in chebulic myrobalan (*Terminalia chebula* Retz.). Electronic Journal of Plant Breeding. 2011;2(1):157-163.
19. Pandey S, Poonia A, Rai S. Optimization of spray drying conditions for the production of quality ber (*Zizyphus mauritiana* Lamk.) fruit powder. Nutrition & Food Science. 2019;49(6):1088-1098.
20. Pareek S. Nutritional composition of jujube fruit. Emirates Journal of Food & Agriculture. 2013;25(6):463-476.
21. Parida S. Effect of drying on nutrition of orange leather fortified with orange peel. Journal of Agricultural Engineering and Food Technology. 2016;3(3):203-209.
22. Patil SH, Shere DM, Sadawarte SK. Development of date-mango leather and evaluation of its physicochemical and sensory properties. Trends in Biosciences. 2017;10(22):4355-4358.
23. Rahman R. Utilization of unexplored mulberry fruit into blended value added products [Doctoral dissertation]. Jammu: Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu; 2021.
24. Sadasivam S, Manickam A. Antinutritional factors. In: Biochemical methods. New Delhi: New International Pvt. Ltd; 2008. p.215-216.
25. Saleem A, Ahmad T, Yasir A, Tahir F, Ali E, Ahmad H, et al. Development and evaluation of fruit leather from guava and jujube blend. International Journal of Health Sciences. 2023;6(S7):6893-6914.
26. Samant D, Mishra NK, Singh AK, Lal RL. Effect of micronutrient sprays on fruit yield and quality during storage in ber cv. Umran under ambient conditions. Indian Journal of Horticulture. 2008;65(4):399-404.
27. Sankanur M, Singh NB, Thakur S, Suresh NV, Verma A. Phenological behavior and reproductive characters of *Terminalia chebula* Retz. an important tree species for sustainable development of sub-tropical Himalayas. Journal of International Academic Research for Multidisciplinary. 2015;2:5075-5083.
28. Sareen A, Gupta RC, Bansal G, Singh V. Comparison of key mineral elements in wild edible fruits of *Zizyphus mauritiana* and *Z. nummularia* using atomic absorption spectrophotometer (AAS) and flame photometer. International Journal of Fruit Science. 2020;20(2):S987-S994.
29. Sharif M, Naz A, Farooq U, Razzaq K, Sher MA, Khan MA, et al. A comprehensive investigation of novel ber (*Zizyphus mauritiana*) products from South Punjab, Pakistan. Journal of Bioresource Management. 2022;9(2):3-11.
30. Sharma R, Rana JC. Nutritional composition and value-added products of Chinese ber (*Zizyphus jujuba* Mill.) growing in Northern hill regions of India. Indian Journal of Natural Products and Resources. 2017;7(4):323-327.

31. Sharon C, Usha V. Effect of storage on nutritional and sensory qualities of breadfruit flour. *Journal of Food Science and Technology*. 2006;43(3):256-258.
32. Singh NB, Thakur S, Sharma K, Sankanur M. Genetic improvement of Harar (*Terminalia chebula* Retz). National Seminar on Smart Farming for Enhancing Input Use Efficiency, Income and Environmental Security. 2017;265:68.
33. Soni S, Sharma HK, Kaushal P, Singh C. Effect of process parameters on the antioxidant activities of bioactive compounds from Harad (*Terminalia chebula* Retz.). *Agricultural Engineering International: CIGR Journal*. 2015;17(2):205-220.
34. Sultan MT, Anwar MJ, Imran M, Khalil I, Saeed F, Neelum S, *et al*. Phytochemical profile and pro-healthy properties of *Terminalia chebula*: A comprehensive review. *International Journal of Food Properties*. 2023;26(1):526-551.
35. Sundriyal M, Sundriyal DC. Wild edible plants of the Sikkim Himalaya: Nutritive values of selected species. *Economic Botany*. 2001;55(3):377-390.
36. Take AM, Bhotmange MG. Preparation of candy from ber-a value addition. *Food Science Research Journal*. 2012;3(2):217-220.
37. Thirumalai T, Kelumalai E, Senthilkumar B, David E. Ethnobotanical study of medicinal plants used by the local people in Vellore District, Tamil Nadu, India. *Ethnobotanical Leaflets*. 2009;(10):165-172.
38. Zahra SM, Hussain S, Mahmood S, Kausar T, Din GMU, Tufail T, *et al*. Shelf stable iron fortified fruit bars development, proximate estimation and organoleptic characterization. *International Journal of Biosciences*. 2020;16(4):111-135.
39. Chavan RB, Thipparaboina R, Kumar D, Shastri NR. Co amorphous systems: A product development perspective. *International journal of pharmaceuticals*. 2016 Dec 30;515(1-2):403-415.