



ISSN Print: 2664-844X
ISSN Online: 2664-8458
NAAS Rating: 4.97
IJAFS 2025; 7(5): 55-61
www.agriculturaljournals.com
Received: 24-03-2025
Accepted: 25-04-2025

Kandarp Parmar
Research Scholar, Department
of Geography, Faculty of
Science, The Maharaja
Sayajirao University of
Baroda, Baroda, Gujarat,
India

Bindu Bhatt
Professor, Department of
Geography, Faculty of Science,
The Maharaja Sayajirao
University of Baroda, Baroda,
Gujarat, India

Corresponding Author:
Kandarp Parmar
Research Scholar, Department
of Geography, Faculty of
Science, The Maharaja
Sayajirao University of
Baroda, Baroda, Gujarat,
India

Evaluating agricultural patterns and crop shifts in Vadodara-Chhotaudepur District of Gujarat, India

Kandarp Parmar and Bindu Bhatt

DOI: <https://www.doi.org/10.33545/2664844X.2025.v7.i5a.380>

Abstract

India is an agrarian country, and agriculture is one of the major sources of food and income in India. Around 20% of Gujarat's economy comes from agriculture sectors. There are observable shifts in Gujarat's crop patterns, land use, and productivity. Many districts in Gujarat are experiencing administrative and environmental changes. The study assesses long-term agriculture patterns from the year 1997-98 to 2022-23 in five-year intervals. The analysis focuses on ten major crops of the region, representing both Kharif and Rabi seasons. Parameters such as agricultural area, production, and yield were examined to evaluate the crop performance. Further, relative yield gap analysis was done to compare the state yield to the study region, which revealed insights into the crop selection and potential of the study area as an agricultural region. The results indicate a significant decline in the area and yield of food crops, while there are significant increases in high-value cash crops such as cotton, tobacco, and oilseeds. These patterns reflect a growing emphasis on commercial agriculture, driven by urban expansion, climate variability, and changing market incentives.

Keywords: Agricultural pattern, agricultural shifts, crop diversification, yield gap, cash crops

Introduction

During the period from 10,000 to 4,000 years ago, agriculture independently emerged at least seven times worldwide (Taiz, 2013) ^[14]. A distinct group of crop species were cultivated in each of these seven primary areas of domestication. For instance, the primary domesticated crops in Asia were rice and millet; in Africa, sorghum; in Mexico, maize, squash, and beans; in South America, the potato and sweet potato and in North America, sunflowers (Taiz, 2013) ^[14]. All of these different crops travelled through historical trade, cultural exchanges & migrants and diversified the agriculture production of particular locality. India also received some of its major crops such as Maize, Potato, Tea, etc from travellers of other countries. Increased reproductive output, non-dehiscent fruits, self-pollination, bigger seeds and fruits, loss of seed dormancy, quick and uniform germination, uniform ripening, annual life cycle, and palatability were all genetic features that were usually selected during domestication (Taiz, 2013) ^[14].

Since its domestication to the age of industrialization agriculture has remain primary source of food and income to the majority of population. In India Agriculture is still the primary source of earning for the rural population and this sector individually represents 23% of India's Gross National Product (GNP), which is playing an important role in the country's socioeconomic growth (Patel *et al.*, 2020) ^[9].

While the world's population grew from 3 to 7 billion people between 1960 and 2010, agricultural productivity per person increased as well, despite only 10% more agricultural land being available for production (Van Ittersum & Cassman, 2013) ^[16]. According to a study by Smil (2001) ^[13] and Von Braun (2005) ^[20] food production has kept up with rising demand over the last few decades. Modern day's major concerns are its distribution and access (Pingali, 2006) ^[10], and there are growing worries about threats to the world's food supply from growing competition from the production of animal feed, biofuels, and climate change (Verhagen *et al.*, 2007) ^[11]. To satisfy the rising demand, yields must be increased and pre- and post-harvest losses must be decreased (Roetter and van Keulen, 2007, Meerburg *et al.*, 2009) ^[11, 8].

Crop yield refers to the amount of harvested product (like grains, fruits, or vegetables) produced per unit area of land. It's a key measure of agricultural productivity, indicating how much output is generated for a given area. Most discussions on crop yield gaps have two main objectives (Van Ittersum & Cassman, 2013) ^[16]. To determine the possible scope for increasing average yields by management adjustments, the first step is to estimate the extent of the yield gap and the second step is to determine the main reasons of the yield gap (Lobell, 2013) ^[7].

In certain areas, the gap between farmers actual yields and what may be theoretically obtained under optimal management has reduced to such an extent that it is both theoretically challenging and economically less viable to increase farmers yields further (Van Ittersum & Cassman, 2013) ^[16]. However, because of their favourable climate, high-quality soil, and occasionally access to irrigation, some areas have a considerably greater potential than others to sustain higher yields in a sustainable way (Van Ittersum & Cassman, 2013) ^[16].




Fig 1: Evolution of population, arable land area, world average wheat and rice yields, fertilizer N use and irrigated area between 1960 and 2010 Source: (Van Ittersum, 2011) ^[15]

Study area

In the central region of Gujarat's mainland, between 21°49'19" and 22°48'37" N latitudes and 72°51'05" and 74°16'55" E longitudes, with the area of 7548-square-kilometer is Vadodara district (Bhowmik & Bhatt, 2023) ^[1]. Fertile ground in Vadodara district is mostly found in agricultural areas and plains, particularly in the district's eastern and central regions (Bhowmik & Bhatt, 2024) ^[2]. Primarily, the Vadodara district area has a sub-tropical climate with moderate humidity because it is situated south of the Tropic of Cancer and in the area between the parched North Gujarat plains and the South Gujarat regions that have substantial rainfall. The coldest and hottest month in Vadodara is typically January and May. The average temperature of January is around 21.5°C (70.7°F) while the temperature of hottest month in Vadodara reaches up to around 40 °C. Vadodara district experience tropical savannah climate. The district experiences high summer temperatures, cool winters, moderate rainfall, and mild humidity, all of which are characteristics of a sub-tropical climate. Rainfall averages 832 mm per year (1981-2002). Many taluka areas in the district have previously experienced mild to severe drought (Sabhaya *et al.*, 2018) ^[12]. According to the 2011 statistics, there are around 4.17 million population living inside the Vadodara-Chhotaudepur district encompassing total area of 7527 km², of which 777.8 km² is forest area (Chandole & Joshi, 2023) ^[3]. There

were 12 Talukas in Vadodara-Chhotaudepur district in 2011, but for better administration and growing population a separate Chhotaudepur district was carved out in year 2014. This explains why two distinct districts were chosen as the entire study area since the available agricultural data from before year 2014 is aggregated for both districts (Vajinath Kantiram Chavan *et al.*, 2024) ^[19].

The district's two main rivers, the Narmada and the Mahi, run along its northwest and southern borders, respectively, while the Dhadhar, a distinct small river system with many tributaries, runs through the district's south-central region. In general, the entire district is separated into three river basins: The Dhadhar, the Mahi, and the Narmada (Gupte, 2011) ^[6]. There are various other rivers also which drains the district like, Dhandhar, Orsang, Vishwamitri and Jambua. Mahi and Narmada Doab, an alluvial plain region with gently sloping terrain, make up the western and southern parts of the Vadodara district. Three general categories can be used to classify the Vadodara district's soils. These Three types of soils are red or hilly, alluvial, brown and black. The agricultural crops grown in the district are diverse and include both food and non-food crops. Food grains such as rice, wheat, jowar, bajra, maize, etc., as well as pulses make up the main food crops. Fruits, vegetables, and sugarcane are other food crops Cotton, Ground nuts, castor, tobacco and many other are example of non-food crops (Gupte, 2011) ^[6].




Fig 2: Map of study area India




Fig 4: Soil Map of Vadodara district (Gupte, 2011) [6]

Relative Yield Gap

Relative yield gap is an approach used to assess the crop production performance of a particular region (i.e., the Vadodara-Chhotaudepur district) by comparing it to the actual observed yield of a broader reference area (i.e., Gujarat State). This approach focuses on intraregional differences to highlight the performance of a particular region to observe the standard of a larger area and can be calculated by the equation (1). Relative yield gap helps policymakers to identify crops that are performing better in that particular state. It also helps to find out the suitable climatic, geographic, and irrigation conditions, infrastructure, and technology adoption for particular crops to perform better.

$$\text{Relative yield gap} = \left(\frac{\text{Yield of District} - \text{Yield of Gujarat State}}{\text{Yield of Gujarat State}} \right) * 100 \quad (1)$$

The relative yield gap results indicate district's yield performance in relation to state average yield. When the yield gap reaches 0% the district matches the state performance, while district produces greater yields than the state average, the percentage yield gap is positive which indicates its higher agricultural performance at a regional level. The district shows signs of high performance based on its favorable agro-climatic situation and effective farming techniques and efficient input use. When the district yield falls beneath state average the yield gap becomes negative

which signals poor production performance. The district's yield below the state average shows an existing yield gap indicating hidden potential of production which could be enhanced through attentive approaches and technological improvement with enhanced agronomic support.

Data

Table 1: Crop-wise agriculture area ('00 ha) of Vadodara-Chhotaudepur District

	1997-98	2002-03	2007-08	2012-13	2017-18	2022-23
Rice	547	472	614	446	446	498.18
Maize	492	538	518	687.22	528.52	407.94
Tur	1054	1000	757	641	675.03	421.65
Wheat	139	105	301	365	205.82	286.28
Groundnuts	130	55	171	32	17.75	10.04
Castor	150	81	166	325	407	425
Cotton	1498	1584	1734	2038	1592.06	1854.79
Tobacco	258	151	116	40	53.98	100.09
Jowar	260	128	67	41	10.59	2.2
Bajra	182	137	113	131	76.14	70.22

Source: (Directorate_Agriculture, 2024)

Table 2: Crop-wise production ('00 MT) of Vadodara-Chhotaudepur District

	1997-98	2002-03	2007-08	2012-13	2017-18	2022-23
Rice	547	380	836	632	709	913
Maize	863	899	915	1493	1228	910
Tur	826	745	858	808	991	526
Wheat	297	240	777	1250	551	814
Groundnuts	208	80	280	43	42	25
Castor	202	105	227	746	822	964
Cotton	2461	2438	4562	8447	8170	8294
Tobacco	314	197	146	68	67	162
Jowar	276	138	67	49	14	3
Bajra	284	180	194	315	168	174

Source: (Directorate_Agriculture, 2024)

Table 3: Crop-wise yield (kg/ha) of Vadodara-Chhotaudepur District

	1997-98	2002-03	2007-08	2012-13	2017-18	2022-23
Rice	1000	807	1362	1417	1579	1690
Maize	1752	1670	1765	2173	1945	2312
Tur	783	745	1133	1261	1542	1207
Wheat	3140	2298	2577	3424	2874	3057
Groundnuts	1600	1472	1637	1327	2094	2514
Castor	1346	1285	1370	2293	2019	2269
Cotton	279	262	447	704	868	776
Tobacco	1221	1301	1261	1700	618	807
Jowar	1062	1075	1000	1195	1316	1526
Bajra	1560	1317	1717	2404	2559	2714

Source: (Directorate_Agriculture, 2024)

Results and Discussion

Trends of crop-wise agricultural area in Vadodara District (1997-2023)

The agricultural practices in Vadodara district's farming areas underwent substantial changes between the years 1997-98 to 2022-23. Foodgrain production area decreased substantially from its first measurement of 2909 thousand hectares in 1997-98 until it dropped to 1786.34 thousand hectares in 2022-23. Foodgrain cultivation area declined year after year because urban development and land use transition forced farmers to choose profit-generating cash

crops. During the kharif season cultivation territory of both cereal and pulse crops reduced drastically. The cultivated area of kharif cereal farming decreased by 598 thousand hectares from 1997-98 until the harvest reached 610.42 thousand hectares in 2022-23. During this corresponding period the kharif pulse cultivation decreased by 767.73 thousand hectares. Two traditional cereal varieties tur and jowar along with bajra have become less popular in cultivation since their farmed land decreased from 1054 to 421.65 thousand hectares respectively (Swagata Bera, 2024) [18]. The reduction in preference or suitability of these coarse grains demonstrates the main reason behind this decline. The cereal group within rabi crops elevates its planting areas consistently and controls its production figures at current levels. The rabi food grain regions managed to restore previous crop swinging patterns and expanded to 545.49 thousand hectares following their initial 384 thousand hectares. The wheat stock for the 2022-23 season expanded its cultivation areas to 286.28 thousand hectares.

The production of oilseeds along with cotton has demonstrated increasing trends as cash crops. Over time the area dedicated to oilseed cultivation increased from 297 to 682.69 thousand hectares simultaneously with a cotton farming area grew from 1498 to 1854.79 thousand hectares. The agricultural policy experiences widespread transformation coming from high returns and increasing market requirements and possible governmental support for cash crops. Pulses witnessed a massive decrease in their cultivated land between 1279 and 521.5 thousand hectares throughout both seasons even though they are nutritionally important and economically valuable crop.

Statistics showed Vadodara has shifted agricultural production toward more profitable crops like cotton alongside oilseeds because kharif food grain farming has taken a decline. Farmers make such decisions considering agroclimatic conditions and follow policy recommendations while considering economic benefits in each specific period.

Trends of crop production in Vadodara District (1997-2023)

The major crop production records of Vadodara district from 1997-1998 to 2022-2023 give insights into both farming pattern adjustments and technological progress as well as climate variability. The constant production levels of coarse grains and pulses together with yield improvements in certain crops shape an overall rising trend. The 2022-23 foodgrain production in Vadodara district came to 3429.70 ('00 MT) while maintaining its maximum level in 2012-2013 at 4788 ('00 MT) starting from an initial value of 3224 ('00 MT) during 1997-98. High levels of production stemmed from enhanced monsoons combined with better agricultural methods and increased rabi harvests during that time. Field production numbers appear to have decreased because farmers decreased their land cultivation as erratic rain patterns combined with constant harvest output rates. The kharif harvests produced varying results of rice and maize throughout successive years. Rice cultivation area decreased to 913.43 during 2022-23 yet production efficiency along with yield grew from 547 to 913.43. The area used to grow maize decreased to reach 909.6 during 2022-23 since yield fluctuations and land reduction occurred after maximum levels at 1493 during 2012-13. Planned tur cultivation decreased by about three hundred percent during

2017-18 to 2022-23 since it started at 990.82 hectares but ended at 525.57 hectares.

The commercial demand for oilseeds and cotton together with their steady performance led to strong growth rates. The overall cotton cultivation area experienced a minor modification yet the production numbers surged past triple from 2461 in 1997-98 to reach 8294.05 in 2022-23 as farmers adopted better crop management techniques and hybrid cultivars. The agricultural strategy of farmers shifted towards oilseed cultivation because minimum support measures and improved market prices prompted this vast jump from 421 to 1325.69 in production. Traditional cereals such as bajra and jowar now generate much less output with reduced significance in the agricultural sector. The agricultural abandonment of Jowar became apparent as its output declined from 276 to only 3.42. Bajra producers maintained only low production quantities during the recent years to match the same results. The alteration in food preferences coupled with budgetary limitations aligns with reduced dietary adequacy. Rabi foodgrain output measured 512 in 2005 but rose significantly up to its current level of 1489.01 due mainly to wheat and other rabi cereal cultivation. Production numbers in the rabi pulse sector substantially increased from 2017-18 to 2022-23 without surpassing the scale of other categories. The production system of Vadodara exhibits a clear transition from basic staple crops to profitable commercial crops. Overall foodgrain production stability is maintained through rabi season success while declining kharif foodgrain and pulse production levels demand interrogation about food security and proper nutrition. Crop diversity together with sustainable agriculture practices and specialized support for essential yet weak producing categories prove important based on these measurements.

Trends of Crop Yield in Vadodara District (1997-2023)

Principal crop yields in Vadodara district from 1997-98 to 2022-23 show that agricultural development took a definite turn due to advanced practices which created improved crop output across the region. Improvements were made to various irrigation and farming technologies and crop management methods which brought significant development in this time period. The output values of cereal crops specifically rice and maize and wheat continue to show steady growth increments. Producers cultivated rice yields at a steady rate from 1000 kg/ha in 1997-98 until they reached 1690 kg/ha in 2022-23. During this time period maize yield production escalated from its initial 1752 kg/ha to 2312 kg/ha. Prior to its peak of 3424 kg/ha in 2012-13 wheat production levels reached 3057 kg/ha because farmers used agricultural inputs effectively to achieve this advancement from 3140 kg/ha. There was significant growth noted in the production levels of oilseed crops. The implementation of crop diversification alongside better seed varieties led India to achieve total oilseed yield of 1810 kg/ha alongside groundnut cultivation yield of 2514 kg/ha. Cotton production in the yearly assessment increased remarkably from 279 kg/ha through 2017-18 reaching its maximum level at 868 kg/ha then dropped slightly to 776 kg/ha during 2022-23 though exceeding traditional historical records. All pulse crops follow their own evaluation patterns differently during this observation phase. Hectare yields of rabi pulse crops increased from 783 to 1608 kg per hectare so kharif pulses demonstrated inconsistent results by

showing 1375 kg per hectare in 2017-18 and measuring 1133 kg per hectare presently. The local climate together with minimal technological advancements have made these production numbers irregular because pulse crops respond strongly to variability in monsoons. The downward movement of total pulse yields starting from 2017 requires special attention from experts who need to focus on improving pulse productivity levels.

The rabi foodgrains production increased by 1799 kg/ha from 1333 kg/ha to 2721 kg/ha during the period while kharif foodgrains experienced a lower growth from 1052 kg/ha to 1477 kg/ha. The rising gap demonstrates how rabi cultivation with irrigation support has replaced traditional kharif cultivation which now faces severe rainfed challenges. Agricultural modernization became evident through total foodgrain yield growth from 1108 to 1890 kg/ha which happened despite changes in climate along with socioeconomic conditions. The performance evaluation shows that crop yields steadily increased for most agricultural assets including cereals and oilseeds although pulses exhibit slower and less reliable yield growth. The empirical data shows why specific actions need to be directed toward weak production areas while confirming that agricultural initiatives and farmer adoption strategies have been successful within Vadodara's specific agricultural environment.

Relative yield gap analysis

Table 4: Relative yield gap analysis of Vadodara-Chhotaudepur District with comparison to state of Gujarat

	1997-98	2002-03	2007-08	2012-13	2017-18	2022-23
Rice	-35.48	-34.92	-29.90	-33.66	-28.55	-33.20
Maize	6.44	-2.11	28.36	29.73	8.84	8.09
Tur	4.54	18.25	2.07	6.41	24.05	3.78
Wheat	32.32	16.83	-14.47	19.05	-1.88	-3.69
Groundnut	17.82	173.10	-7.83	123.78	-13.54	-2.18
Castor	-31.71	10.02	-30.74	15.28	-20.54	1.29
Cotton	-21.63	49.71	-23.06	105.25	32.52	16.34
Tobacco	-26.31	-24.67	-26.77	-0.93	-79.30	-65.20
Jowar	31.76	37.82	-26.58	-11.35	-4.22	-63.95
Bajra	14.29	36.48	21.09	37.92	5.31	7.19

Positive (%) = District is yielding more than the state → District is overperforming

Negative (%) = District is yielding less than the state → District is underperforming → yield gap exists

The yield of major ten crops of Vadodara district compare to Gujarat state shows a distinct crop-specific and time-based inequalities. Rice consistently shows negative yield gap across the study period, indicating Vadodara's rice yield is significantly lower than the state. The yield gaps of rice in Vadodara continue to be negative in all years between 29.90% and -35.48% due to its lower production than the state average, which highlights Vadodara's proportional disadvantage in rice production, owing to climatic conditions and improper irrigation facilities.

Data indicates that maize and Tur (pigeon pea) exhibit positive yield gaps in most recorded years achieving maximum rates of 29.73% for maize in 2012-13 and 24.05% for Tur in 2017-18. The production results show a comparative advantage for pulses and coarse cereals mainly because of favourable soil conditions and local agricultural practices together with climatic adaptations.




Fig 5: Relative Yield Gap analysis of Vadodara-Chhotaudepur district

The performance of wheat has exhibited an overall negative trend which might stem from deteriorating irrigation methods coupled with declining soil fertility and unfavourable market conditions. The initial 32.32% advantage in 1997-98 wheat supply evolved into a -3.69% disadvantage in 2022-23 thus demonstrating deteriorating relative market performance. The production data shows significant fluctuations in Groundnut because the positive variation between, 2002-2003 reached 173.10% and again reached 123.78% in 2012-2013. The crop production experiences sudden large decreases in different years suggesting irregular production patterns which could result from rain fed cultivation and pests' behaviour.

The yield gap performance of Castor fluctuates having brief positive years in 2002-03 and 2012-13 but demonstrating significant underperformance during recent times (-20.54% in 2017-18). The Vadodara district reveals significant growth in cotton production which is indicated by two large positive results of 105.25% in 2012-13 and 32.52% in 2017-18 representing strong production capabilities possibly resulting from modified seed variety adoption, improved pest management and suitable soil types. The negative yield gaps recorded in tobacco cultivation for recent years reached their worst levels at -79.30% and -65.20% which demonstrates significant decline when compared to statewide performance. The yield gaps of jowar and Bajra reveal significant changes in the data since moving from early advantage to major negative levels especially jowar reached -63.95% in 2022-23. The poor adaptation together with inadequate crop management practices characterizes the farming practices in the district's ecological zone. This analysis can be used into policy formulation, and district level agricultural planning aiming at increasing overall production and sustainability of the region.

Conclusion

Throughout a period from 1997-98 until 2022-23 Vadodara district underwent a noticeable shift in its agricultural sector as farmers began focusing on commercial crops instead of foodgrains. The reduction of kharif foodgrain and pulse cultivation areas happened primarily because of shifting land use patterns, altering food preferences, policy incentives and economic profitability changes. The farming of wheat in Rabi season maintained satisfactory foodgrain production levels but commercial farming of cotton and

oilseed grew much higher because of strong market demands backed by technological advancements. Statistics from crop yields reveal better productivity in cereal and oilseed production however pulse and coarse grain yields remain inconsistent because of deficient irrigation and unreliable monsoon patterns. Data from yield gap analysis demonstrates that the rice and wheat and bajra production in Vadodara district remains below state averages while the district achieves better results in maize, Tur and cotton results when compared against state levels during particular years.

Various sustainability programs for food production require strategic measures for specific crops in designated areas. Strategic investments into irrigation systems along with pulse crop productivity improvement projects combined with proper crop rotation schemes will both eliminate existing negative yield gaps and optimize all land resources across the district.

Acknowledgement

This research is financially supported by University Grants Commission, India as UGC-NET/JRF to Kandarp Parmar. Additionally we would like to thank the Directorate of Gujarat, Department of Agriculture, farmer's welfare and co-operation, Government of Gujarat for providing necessary data.

References

1. Bhowmik S, Bhatt B. Spatiotemporal analysis of land surface temperature owing to NDVI: A case study of Vadodara District, Gujarat. *J Geomatics*. 2023;17(1):83-90. DOI: 10.58825/jog.2023.17.1.83.
2. Bhowmik S, Bhatt B. Drought Monitoring Using MODIS Derived Indices and Google Earth Engine Platform for Vadodara District, Gujarat. *J Indian Soc Remote Sens*. 2024 Apr;52:1-16. DOI: 10.1007/s12524-024-01922-1.
3. Chandole V, Joshi GS. Case study of rainfall and temperature assessment through trend and homogeneity analyses in Vadodara and Chhotaudepur district of Gujarat State, India. *Environ Monit Assess*. 2023 May;195(5):589. DOI: 10.1007/s10661-023-11089-w.
4. Vermeulen SJ, Meerburg BG, Schils RLM, Verhagen J, Kuikman PJ. Feeding the world's increasing population while limiting climate change impacts: linking N2O and

- CH₄ emissions from agriculture to population growth. *Environ Sci Policy*. 2010 Feb;13(2):89-96. DOI: 10.1016/j.envsci.2009.11.001.
5. Directorate of Agriculture, Gujarat. Area, Production and Yield. 2024 [cited 2025 Apr 29]. Available from: <https://dag.gujarat.gov.in/Home/AreaProductionAndYield>.
 6. Gupte P. Vadodara District Report-CGWB-WCR 2011. 2011 [cited 2025 Apr 29]. Available from: https://www.researchgate.net/publication/234012681_Vadodara_District_Report_-_CGWB-WCR_2011.
 7. Lobell DB. The use of satellite data for crop yield gap analysis. *Field Crops Res*. 2013 Mar;143:56-64. DOI: 10.1016/J.FCR.2012.08.008.
 8. Meerburg BG, Singleton GR, Leirs H. The year of the rat ends-time to fight hunger! *Pest Manag Sci*. 2009 Apr;65(4):351-352.
 9. Patel MP, Gami B, Patel A, Patel P, Patel B. Climatic and anthropogenic impact on groundwater quality of agriculture dominated areas of southern and central Gujarat, India. *Groundwater Sustain Dev*. 2020 Jan;10:100306. DOI: 10.1016/j.gsd.2019.100306.
 10. Pingali P. Economic reforms and food security: the impact of trade and technology in South Asia. *Agric Econ*. 2006 Dec;35 Suppl 1:115-128.
 11. Roetter RP, van Keulen H. Food security. In: Roetter RP, van Keulen H, Kuiper M, Verhagen J, Van Laar HH, editors. *Science for Agriculture and Rural Development in Low-Income Countries*. Dordrecht (Netherlands): Springer, 2007, p. 27-56.
 12. Sabhaya M, Solanki PM, Pancholi V, Singh P. Geotechnical Characteristics of soils to identify substratification of Vadodara City, Gujarat, Western India. *J Indian Geophys Union*. 2018;21(2):105-115.
 13. Smil V. *Feeding the World: A Challenge for the TwentyFirst Century*. Cambridge (MA): MIT Press; c2001.
 14. Taiz L. Agriculture, plant physiology, and human population growth: past, present, and future. *Theor Exp Plant Physiol*. 2013 Oct;25(3):167-178. DOI: 10.1590/s2197-00252013000300001.
 15. Van Ittersum MK. Future harvest the fine line between myopia and utopia. Wageningen (Netherlands): Wageningen University, 2011 [cited 2025 Apr 29]. Available from: <https://edepot.wur.nl/169680>.
 16. Van Ittersum MK, Cassman KG. Yield gap analysis-Rationale, methods and applications-Introduction to the Special Issue. *Field Crops Res*. 2013 Mar;143:1-3. DOI: 10.1016/J.FCR.2012.12.012.
 17. Verhagen J, Woosten H, Jager DA. Agriculture and environment. In: Roetter RP, van Keulen H, Kuiper M, Verhagen J, Van Laar HH, Editors. *Science for Agriculture and Rural Development in Low-Income Countries*. Dordrecht (Netherlands): Springer, 2007, p. 57-82.
 18. Bera S. Spatiotemporal analysis of land use/land cover changes and vegetation health in south Andaman district using remote sensing and GIS. *Int J Geogr Geol Environ*. 2024;6(2):57-65. DOI: 10.22271/27067483.2024.v6.i2a.289
 19. Chavan VK, Patil PK. A study of agriculture land use efficiency of Dhule district: A geographical analysis. *Int J Geogr Geol Environ*. 2024;6(1):170-173. DOI: 10.22271/27067483.2024.v6.i1c.21
 20. Braun VJ. Agricultural economics and distributional effects. *Agric Econ*. 2005 Jan;32(1):1-20.