

International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
NAAS Rating: 5.29
IJABR 2025; 9(6): 1092-1095
www.biochemjournal.com
Received: 21-03-2025
Accepted: 22-04-2025

Subhi Dwivedi
M.Sc. Scholar, Rama
University, Mandhana,
Kanpur, Uttar Pradesh, India

Vinay Joseph Silas
Assistant Professor, Faculty of
Agricultural Sciences and
Allied Industries Rama
University, Mandhana,
Kanpur, Uttar Pradesh, India

Jitendra Kumar
Assistant Professor, Rama
University, Mandhana,
Kanpur, Uttar Pradesh, India

Deepak Lall
Assistant Professor, SHUATS,
Prayagraj, Uttar Pradesh,
India

Maseera Anjum
M.Sc. Scholar, Rama
University, Mandhana,
Kanpur, Uttar Pradesh, India

Corresponding Author:
Vinay Joseph Silas
Assistant Professor, Faculty of
Agricultural Sciences and
Allied Industries Rama
University, Mandhana,
Kanpur, Uttar Pradesh, India

A study on the impact of INM on yield and quality parameter of strawberry (*Fragaria x ananassa*) cv. sweet Charlie under protected cultivation

Subhi Dwivedi, Vinay Joseph Silas, Jitendra Kumar, Deepak Lall and Maseera Anjum

DOI: <https://www.doi.org/10.33545/26174693.2025.v9.i6m.4664>

Abstract

An experiment was conducted to study the “Impact of Integrated nutrient Management on growth, yield and quality of strawberry (*Fragaria x ananassa*)” at Department of Horticulture, Institute of Agriculture Sciences, Rama university Kanpur the experiment was laid out according to Randomized Blocks Design with eight treatments replicated thrice. The result reveals that the maximum plant height (21.38), plant spread (24.51 cm), number of leaves (18.69), leaf area (82.42cm²), marketable fruit (83.04%), number of harvest (5.87), yield/plant (227.74 g), TSS (9.62oBrix), TSS/acid ratio (14.77), ascorbic acid (61.25 mg/100 gm), were recorded in plants treated with treatment T₈(Integrated Nutrient + Azotobacter + other biofertilizers).The minimum days taken in flowering (54.77 days) and minimum titratable acidity (0.62%) were also recorded in treatment T₈ Azotobacter 50% + Aspergillum 50% + NPK 50%. Considering overall performances, the study revealed that the combined application of Integrated Nutrient could be a good source of plant nutrients for organic strawberry cultivation.

Keywords: Strawberry, sweet Charlie, INM, Azotobacter, PSB, aspergillums, organic

Introduction

The strawberry (*Fragaria x ananassa* Duch.), a member of the Rosaceae family, is a delicious fruit crop known for its bright color, taste and aroma. It is rich in Vitamin C, minerals, and antioxidants, making it an important source of human nutrition. (Kumar *et al.*, 2018) ^[6]. In India strawberry cultivation is increasing rapidly along with Maharashtra. With increasing demand and health awareness, its cultivation is growing in India. However, higher dependency on chemical fertilizers harms the soil and environmental economy. Integrated Nutrient Management (INM) involves the use of both organic and inorganic nutrients along with biofertilizers. The cultivation of strawberries is experiencing rapid growth, particularly in regions such as Maharashtra, Himachal Pradesh, and Uttar Pradesh, attributed to the higher economic returns per unit area. To achieve successful strawberry cultivation, it is crucial to manage nutrients effectively. The overuse of chemical fertilizers results in environmental pollution and degradation of soil quality (Chadha *et al.*, 2018) ^[1]. Integrated Nutrient Management (INM) represents an environmentally friendly approach that integrates chemical fertilizers, organic manures such as FYM and vermicompost, along with biofertilizers like Azotobacter and Azospirillum. This method enhances nutrient availability, promotes soil health, and boosts microbial activity. Numerous studies have indicated that INM not only promotes vegetative growth but also enhances both the yield and quality of fruits. Biofertilizers are vital for nitrogen fixation, hormone production, and enhancing disease resistance. The application of organic amendments improves the soil's water retention capacity and nutrient levels, leading to superior plant performance. Consequently, this study was undertaken to assess the effects of various INM combinations on the growth, yield, and quality of strawberries (cv. Sweet Charlie) under conditions of protected cultivation.

Material and Methods

The study was conducted at the Department of Horticulture, Rama University, Kanpur,

during the Rabi season of 2024-25. Kanpur is located in central region of Uttar Pradesh which lies between 25°30'44" North latitude and 78°32'31" East longitude at an elevation of 285 meters above mean sea level. The experimental design employed was Randomized Block Design (RBD) featuring three replications and nine treatments.

The strawberry variety utilized for this research was Sweet Charlie. Each treatment plot comprised 16 plants with a spacing of 30 × 25 cm on raised beds within a polyhouse. Before transplanting the raised beds are covered with black polyshets to conserve moisture. And planting material of strawberry sweet Charlie were purchased from a private organization Pune by Amazon.

Replication 3

Plot size 2.0×1.2 m

Plant Population 240 plants (10 per treatment and 3 Replication)

The treatment combinations included

T₀: Control

T₁: Recommended dose of NPK

T₂: Azotobacter (100%)

T₃: Azospirillum (100%)

T₄: FYM @ 20 t/ha

T₅: Azotobacter (50%) + Azospirillum (50%)

T₆: Azotobacter (100%) + NPK (50%)

T₇: Azospirillum (100%) + NPK (50%)

T₈: Azotobacter (50%) + Azospirillum (50%) + NPK (50%)

Observations were made on various growth parameters at 30,60,90,120 days after planting growth characters (plant height, spread, number of leaves, leaf area), yield parameters (fruit number, average fruit weight, yield/plant), and quality parameters (TSS, acidity, and ascorbic acid content). The data gathered were analyzed using analysis of variance (ANOVA) at a 5% significance level.

Results and Discussion

The findings revealed notable variations among the treatments concerning all growth, yield, and quality traits. T₈ (Azotobacter 50% + Azospirillum 50% + NPK 50%) surpassed all other treatments.

Growth Parameters

At 120 days post-planting, T₈ achieved the greatest plant height (21.38 cm), followed by T₆ and T₇. The lowest plant height (18.07 cm) was recorded in T₀. Additionally, T₈ exhibited the largest plant spread (24.51 cm) and leaf area (118.2 cm²). The improved growth may be attributed to the balanced nutrient supply and the stimulation of plant hormones by biofertilizers.

Table 1: Impact of INM on Plant Height, Spread, Number of leaves per plant and Leaf Area

Treatments	Plant Height	Number of leaves per plant	Leaf Area	Plant Spread n-s direction
T ₀	18.07	16.79	78.49	22.70
T ₁	19.45	17.57	79.87	23.14
T ₂	20.33	17.53	79.45	22.96
T ₃	20.31	17.80	79.60	22.92
T ₄	20.39	17.86	79.63	23.78
T ₅	20.43	17.93	79.76	23.04
T ₆	20.62	18.73	80.74	23.73
T ₇	20.86	18.71	80.67	23.37
T ₈	21.38	19.16	82.42	24.51

Yield Parameters

Treatment T₈ recorded the highest average marketable percentage (85.12%), the lowest unmarketable percentage (17.38%), fruit yield per plant (227.74g), Number of harvest per plant 5.87, The lowest yield (128.26 g/plant) and

number of harvest of plant 4.04 was noted in the control (T₀). The combination of biofertilizers and partial NPK likely enhanced nutrient use efficiency and fruit development.

Table 2: Impact of INM on Yield Parameters:

Treatments	Marketable%	Unmarketable %	Yield per Plant	No. of Harvest
T ₀	74.11	28.39	128.26	4.04
T ₁	78.39	24.11	178.16	4.95
T ₂	77.14	25.36	156.39	4.89
T ₃	77.02	25.48	153.19	4.33
T ₄	81.07	21.43	206.82	5.06
T ₅	77.94	24.56	174.24	4.92
T ₆	81.04	21.46	202.94	5.05
T ₇	78.62	23.88	187.42	4.96
T ₈	85.12	17.38	227.74	5.87
CD	0.13	0.14	0.02	0.42
Sem±	0.05	0.05	0.01	0.15

Quality Parameters

T₈ also demonstrated the best quality metrics with the highest TSS (9.65 °Brix), the lowest acidity (0.79%), and the highest ascorbic acid content (57.1 mg/100 g). This supports the beneficial impact of INM on enhancing fruit

flavor and nutritional quality.

The integrated application of NPK with azotobacter and Azospirillum likely enhanced nutrients availability and assimilation, thus improving overall crop production.

These Findings are consistent with previous research (Negi *et al.*, 2021,) [8], emphasizing the synergistic effects of organic and biological sources with chemical fertilizers.

Table 3: Impact of INM on Quality Parameters:

Treatment	TSS (° Brix)	Acidity (%)	Ascorbic Acid (mg/100 g)
T ₀	7.88	0.83	56.21
T ₁	8.65	0.77	57.89
T ₂	8.30	0.79	57.77
T ₃	8.23	0.80	57.88
T ₄	9.32	0.69	58.98
T ₅	8.54	0.78	57.81
T ₆	9.25	0.70	58.90
T ₇	8.86	0.75	58.03
T ₈	-	0.65	61.25
CD	0.74	0.46	0.88
Sem±	0.26	0.16	0.31

Summary

The study titled “Evaluation of Integrated Nutrient Management on Growth, Yield, and Quality of Strawberry (*Fragaria x ananassa* Duch.)” was conducted during 2024-2025 at the Department of Horticulture, Rama University, Kanpur, Uttar Pradesh. The key findings from this research are summarized below.

Plant Height: The tallest plants (21.38 cm) were observed in the treatment combining integrated nutrient management (INM) with Azotobacter and other biofertilizers, while the shortest (18.79 cm) were in the control treatment.

Plant Spread: The widest plant spread (24.51 cm) was recorded in the INM + Azotobacter + other biofertilizers treatment, with the narrowest (21.67 cm) in the control.

Number of Leaves: The highest leaf count (18.69) was achieved in the INM + Azotobacter + other biofertilizers treatment, compared to the lowest (15.67) in the control.

Leaf Area: The largest leaf area (82.42 cm²) was recorded in the INM + Azotobacter + PSB treatment, while the smallest (78.49 cm²) was in the control.

Yield per Plant: The highest yield per plant (227.74 g) was observed in the INM + Azotobacter + other biofertilizers treatment, with the lowest (128.26 g) in the control.

Marketable Fruit Percentage: The INM + Azotobacter + other biofertilizers treatment produced the highest percentage of marketable fruit, while the control had the highest percentage of unmarketable fruit.

Number of Harvests: The greatest number of harvests (5.87) was recorded in the INM + Azotobacter + other biofertilizers treatment, compared to the fewest (4.04) in the control.

Total Soluble Solids (TSS): The highest TSS (9.62 °Brix) was found in the Azospirillum + Azotobacter treatment, while the lowest (7.88 °Brix) was in the control.

TSS/Acid Ratio: The highest TSS/acid ratio (14.77) was observed in the INM + Azotobacter + other biofertilizers treatment, with the lowest (9.14) in the control.

Ascorbic Acid: The highest ascorbic acid content (61.25 mg/100 g) was recorded in the INM + Azotobacter + organic fertilizers treatment, while the lowest (56.21 mg/100 g) was in the control.

Conclusion

The results of this study indicate that the application of various organic nutrient sources significantly enhanced vegetative growth, yield, and quality parameters of strawberry crops. The treatment combining integrated

nutrient management with Azotobacter and other biofertilizers yielded the best outcomes, including maximum plant height (21.38 cm), plant spread (24.51 cm), leaf count (18.69), leaf area (82.42 cm²), marketable fruit percentage (83.04%), yield per plant (227.74 g), number of harvests (5.87), TSS (9.62 °Brix), TSS/acid ratio (14.77), ascorbic acid content (61.25 mg/100 g), and minimum titratable acidity (0.62%). Therefore, integrating INM with Azotobacter and other biofertilizers in strawberry cultivation can significantly improve yield and economic returns.

References

- Chadha KL. Handbook of Horticulture. New Delhi: ICAR; 2018.
- Sharma V, Gaur GS. Effect of biofertilizers and inorganic fertilizers on growth and yield of strawberry. *Progressive Horticulture*. 2009;41(2):221-224.
- Singh AK, Ram RB, Yadav PK. Effect of integrated nutrient management on growth, yield and quality of strawberry cv. Chandler. *Indian Journal of Horticulture*. 2011;68(2):210-213.
- Kumar R, Verma V, Meena VS. INM for enhancing productivity of horticultural crops. *Journal of Eco-friendly Agriculture*. 2018;13(1):21-26.
- Meena RK, Patel JC, Jain MP. Role of *Azospirillum* and *Azotobacter* on vegetative growth and yield of strawberry. *Asian Journal of Horticulture*. 2020;15(2):267-270.
- Yadav RK, Sharma A, Verma V. Impact of integrated nutrient management on horticultural crops. *Journal of Horticultural Science*. 2021;18(2):110-114.
- Jat RK, Singh R, Meena VS. Role of biofertilizers in sustainable crop production. *International Journal of Current Microbiology and Applied Sciences*. 2020;9(1):1250-1257.
- Negi Y, Sharma A, Yadav S. Nutrient dynamics in strawberry under protected conditions. *Indian Journal of Horticulture*. 2021;78(4):564-570.
- Sharma P, Verma V. Use of *Azotobacter* and *Azospirillum* in fruit crops: A review. *Plant Archives*. 2018;18(2):1115-1119.
- Singh SP, Patel SR, Ghosh S. Organic and inorganic nutrient effect on quality in fruit crops. *Asian Journal of Horticultural Science*. 2020;9(3):203-209.
- Meena VS, Verma J. Role of INM in enhancing soil fertility. *Agro Journal*. 2020;10(2):55-59.
- Ghosh P, Singh A, Kumari N. Organic and inorganic nutrient management in strawberry (*Fragaria x ananassa* Duch.) cv. Chandler. *International Journal of Chemical Studies*. 2019;7(4):2023-2027.
- Patel NL, Parmar BR, Desai DT. Integrated nutrient management in fruit crops: A review. *Agricultural Reviews*. 2017;38(3):218-222.
- Bairwa HL, Meena ML. Effect of INM on quality and shelf life of strawberry. *International Journal of Agriculture Sciences*. 2017;9(4):3701-3703.
- Pathak RA, Shukla AK. Influence of INM on productivity of strawberry cv. Sweet Charlie. *Plant Archives*. 2019;19(2):4029-4032.
- Sharma RR, Singh R. Influence of biofertilizers on fruit quality of strawberry. *Indian Journal of Agricultural Sciences*. 2009;79(8):613-617.
- Singh G, Lal S. INM effects on soil health and strawberry yield under protected cultivation.

- International Journal of Agricultural Innovations and Research. 2020;8(6):132-136.
18. Khan A, Alam M. Biofertilizers in fruit production: Mechanism and benefits. *Journal of Crop and Weed*. 2018;14(3):130-137.
 19. Lal R, Kumawat BL. INM practices for strawberry under low tunnel conditions. *The Pharma Innovation Journal*. 2020;9(9):314-317.
 20. Tomar M, Yadav P. Effect of INM on growth, yield and shelf life of strawberry. *Journal of Horticultural Research*. 2021;29(1):77-84.
 21. Naik D, Mishra HN. INM effect on strawberry under drip irrigation. *Journal of Water and Soil Conservation*. 2022;21(1):59-64.
 22. Sharma VK, Singh K. Influence of organic and inorganic sources on strawberry cv. Sweet Charlie. *HortFlora Research Spectrum*. 2021;10(3):145-148.
 23. Kaur G, Gill KS. Performance of strawberry with biofertilizers under polyhouse. *Agricultural Science Digest*. 2019;39(2):122-126.
 24. Kumar R, Rattan RK. Integrated nutrient management for improving crop productivity and soil health. *Journal of Soil and Water Conservation*. 2021;20(3):112-117.
 25. Das B, Mondal T. Biofertilizers and their role in sustainable horticulture. *Environment & Ecology*. 2018;36(2):493-498.