



ISSN Print: 2664-9926
 ISSN Online: 2664-9934
 IJBS 2024; 6(1): 172-175
www.biologyjournal.net
 Received: 06-01-2024
 Accepted: 11-02-2024

R Vikesh Iyer
 Department of Medicinal Plant
 Research, Kavita Memorial
 Medical College, Pune,
 Maharashtra, India

Role of plant secondary metabolites in biotic interactions and defense

R Vikesh Iyer

DOI: <https://www.dx.doi.org/10.33545/26649926.2024.v6.i1c.428>

Abstract

Plants, being sessile organisms, have evolved complex biochemical strategies to cope with biotic pressures from herbivores, pathogens, and competing plants. Among these strategies, the production of secondary metabolites plays a central role in mediating plant interactions with other organisms and in fortifying defense mechanisms. These metabolites—including alkaloids, phenolics, terpenoids, and glucosinolates—are not directly involved in primary metabolism but confer evolutionary advantages by deterring predators, attracting beneficial organisms, and modulating plant-plant competition. This paper reviews the biosynthetic pathways, ecological significance, and defensive roles of plant secondary metabolites in various biotic contexts, emphasizing experimental findings and recent advances in metabolomics and molecular biology. Understanding these interactions provides valuable insights into natural defense systems and offers promising avenues for sustainable agriculture and biocontrol applications.

Keywords: Plant secondary metabolites, biotic interactions, defense mechanisms, herbivores, pathogens

1. Introduction

Plants interact continuously with a diverse range of organisms, including herbivorous insects, fungal and bacterial pathogens, nematodes, and neighboring plant species. These interactions have shaped plant evolution, prompting the development of sophisticated defense mechanisms. One of the most effective strategies involves the production and release of plant secondary metabolites (PSMs)—organic compounds that are not essential for growth or reproduction but confer survival advantages under stress. These metabolites often function as chemical signals, toxins, repellents, or antimicrobials. They are central to the plant's dynamic response to biotic challenges and are tightly regulated both spatially and temporally.

Plant secondary metabolites not only serve in direct defense (e.g., poisoning herbivores or pathogens) but also in indirect defense by attracting predatory insects or activating systemic resistance responses. Additionally, allelopathic compounds influence plant-plant competition and community composition. This review explores the defensive roles of these compounds across multiple biotic interactions, their biosynthetic diversity, and the underlying molecular mechanisms.

2. Classification and Biosynthesis of Secondary Metabolites

Plant secondary metabolites are an exceptionally diverse group of organic compounds categorized based on their biosynthetic origin and structural features. Though not directly involved in the fundamental physiological processes of growth, development, or reproduction, their ecological significance lies in mediating plant interactions with other organisms and facilitating defense. The three major classes of plant secondary metabolites include terpenoids, phenolic compounds, and nitrogen-containing compounds (including alkaloids and glucosinolates). Each group arises from distinct metabolic pathways and contributes uniquely to plant defense.

Terpenoids

Terpenoids (also known as isoprenoids) form the largest class of secondary metabolites, derived from the five-carbon precursor isopentenyl diphosphate (IPP), produced through the

Corresponding Author:
R Vikesh Iyer
 Department of Medicinal Plant
 Research, Kavita Memorial
 Medical College, Pune,
 Maharashtra, India

mevalonate (MVA) and methylerythritol phosphate (MEP) pathways. They range from simple monoterpenes to complex tetraterpenes like carotenoids. Many terpenoids act as feeding deterrents to herbivores, antimicrobials against pathogenic fungi and bacteria, and attractants for pollinators or predatory insects. For instance, volatile terpenes released upon herbivore attack serve as airborne signals that recruit natural enemies of herbivores—an indirect defense mechanism.

Phenolic Compounds

Phenolics are synthesized primarily via the phenylpropanoid pathway and include compounds such as flavonoids, tannins, lignins, and coumarins. They are known for their antioxidant properties, UV-protective roles, and structural functions. In defense, phenolic compounds fortify plant cell walls through lignification and act as antimicrobial agents. Flavonoids, for example, exhibit broad-spectrum antifungal and antibacterial activity, while tannins deter herbivory by binding to proteins and reducing digestibility.

Alkaloids and Other Nitrogen-Containing Compounds

Alkaloids are a chemically diverse class containing nitrogen atoms, often synthesized from amino acids such as tryptophan, tyrosine, or ornithine. They include pharmacologically significant molecules like morphine, nicotine, and quinine. Alkaloids are typically toxic to insects and mammals, acting through interference with neurotransmitter function or inhibition of metabolic enzymes. Glucosinolates, prevalent in the Brassicaceae family, yield toxic isothiocyanates upon enzymatic hydrolysis, which deter generalist herbivores and microbial invaders.

Biosynthesis Regulation

The biosynthesis of secondary metabolites is tightly regulated by environmental cues and phytohormonal signaling networks, particularly jasmonic acid, salicylic acid, and ethylene. These hormones mediate transcriptional activation of genes involved in biosynthetic pathways upon detection of herbivory, infection, or wounding. The spatial distribution of biosynthetic enzymes—localized to specific tissues such as trichomes or laticifers—further supports targeted chemical defense.

Tissue-Specific and Inducible Production

Unlike primary metabolites, the production of secondary compounds is often tissue-specific and inducible. For example, glandular trichomes on the leaf surface may accumulate high concentrations of terpenoids, while root exudates may contain allelopathic flavonoids or saponins. Upon pathogen recognition, certain alkaloids or phytoalexins may be synthesized *de novo* as part of localized or systemic acquired resistance responses.

3. Defensive Roles in Plant-Herbivore Interactions

Plants have co-evolved with herbivores over millions of years, developing intricate defense mechanisms to deter feeding and minimize tissue damage. Among the most crucial of these defenses is the synthesis of secondary metabolites that serve as chemical barriers against herbivory. These compounds can act through diverse modes—deterrence, toxicity, anti-digestive properties, and

behavioral disruption—to reduce the attractiveness or nutritional value of plant tissues to herbivores.

One of the best-known examples is the use of alkaloids like nicotine in tobacco plants (*Nicotiana spp.*), which interfere with the central nervous system of insects, leading to paralysis or death. Similarly, pyrrolizidine alkaloids found in *Senecio* species are toxic to many generalist herbivores but tolerated by some specialist insects, underscoring the evolutionary arms race between plants and herbivores. Terpenoids such as limonoids in citrus and cucurbitacins in cucumbers serve as feeding deterrents due to their bitter taste and disruption of insect hormonal balance.

Phenolic compounds, especially tannins and flavonoids, contribute to defense by reducing the digestibility of plant tissues. Tannins can precipitate dietary proteins in the herbivore gut, limiting nutrient absorption and growth. In some cases, phenolics are stored in inactive forms and rapidly activated upon tissue damage, ensuring an efficient but economical defense system. The release of phenolics also generates reactive oxygen species (ROS), creating an oxidative environment harmful to invading organisms.

Furthermore, secondary metabolites often serve indirect roles by facilitating tritrophic interactions. For example, volatile organic compounds (VOCs) such as methyl jasmonate or green-leaf volatiles are emitted in response to herbivory and act as airborne signals that attract parasitoids or predators of the herbivores. This mechanism enhances plant fitness by enlisting biological allies rather than relying solely on chemical warfare.

Plants also exhibit inducible defenses, whereby the biosynthesis of secondary metabolites is triggered upon herbivore attack. This adaptive response conserves metabolic resources while still providing timely protection. The jasmonic acid signaling pathway plays a central role in activating such inducible defenses, upregulating genes responsible for the biosynthesis of alkaloids, terpenes, and phenolic glycosides.

4. Roles in Plant-Pathogen Defense

Plant secondary metabolites are crucial for defense against pathogens like bacteria, fungi, viruses, and nematodes. These compounds act as both constitutive and inducible defenses, exhibiting antimicrobial, antifungal, and antiviral properties. Phytoalexins, synthesized in response to infection, inhibit pathogen growth—for example, medicarpin and pisatin in legumes disrupt fungal invasion. In crucifers, glucosinolates and their breakdown products (e.g., isothiocyanates) impair microbial enzymes and membranes through reactive sulfur species.

Phenolic compounds, activated via the phenylpropanoid pathway upon pathogen recognition, contribute to both chemical and structural defenses. Lignin precursors reinforce cell walls, while flavonoids like quercetin and kaempferol inhibit microbial metabolism and quorum sensing. Terpenoids, another key group, include saponins that disrupt microbial membranes and volatile compounds like pinene and limonene that deter pathogens and trigger immune signaling.

Salicylic acid (SA), a phenolic signal molecule, mediates systemic acquired resistance (SAR) by activating pathogenesis-related genes and preparing distant tissues for future attacks. Plants strategically accumulate and compartmentalize these compounds to target infected areas while protecting healthy tissue.

Advances in metabolomics and transcriptomics have deepened understanding of how plants modulate secondary metabolite production based on pathogen type and infection severity, enabling precise and efficient immune responses.

5. Involvement in Allelopathy and Plant-Plant Interactions

Beyond their roles in defending against herbivores and pathogens, plant secondary metabolites are essential mediators of allelopathy—a biological phenomenon where one plant releases biochemicals that influence the growth, survival, and reproduction of neighboring plants. These interactions can be antagonistic or facilitative and are particularly significant in shaping plant community structure, determining competitive dominance, and influencing ecological succession.

Allelopathic compounds are released through various mechanisms: volatilization from leaves, exudation from roots, leaching from fallen litter, or direct secretion from wounded tissues. Once in the soil or air, these compounds can inhibit seed germination, restrict root elongation, alter microbial communities, or interfere with hormone signaling in adjacent plants.

Phenolic acids such as ferulic, caffeic, and p-coumaric acids are well-documented allelochemicals. Released by the roots or decomposing residues of cereals and legumes, these compounds can disrupt cell membrane integrity, reduce chlorophyll biosynthesis, and cause oxidative stress in target species. In rice (*Oryza sativa*), the release of phenolic-rich root exudates is associated with the suppression of nearby weed species, offering a competitive advantage in dense agricultural systems.

Terpenoids also play significant allelopathic roles. For example, monoterpenes like camphor and cineole released by *Salvia* species can accumulate in soil and reduce germination of competing annuals. Similarly, *Eucalyptus* trees emit volatile terpenes that create a chemically inhospitable zone beneath their canopy, known as the "Eucalyptus effect," inhibiting undergrowth and promoting monoculture stands.

Alkaloids, though primarily associated with anti-herbivore roles, also show allelopathic activity in some cases. For instance, quinolizidine alkaloids from *Lupinus* species have been found to hinder root development in neighboring plants, especially in nutrient-poor soils, thereby reducing competition for limited resources.

Moreover, secondary metabolites influence not just plant-plant interactions directly, but also indirectly through microbial mediation. Allelochemicals can alter soil microbial communities, shifting the balance of beneficial mycorrhizal associations or suppressing nitrogen-fixing bacteria for neighboring plants, creating an asymmetric advantage for the allelopathic plant.

In agricultural contexts, allelopathy has been harnessed in crop rotation and intercropping strategies. Species such as rye (*Secale cereale*) and sorghum (*Sorghum bicolor*) release benzoxazinoids and sorgoleone, respectively, both potent inhibitors of weed growth. These properties make secondary metabolites valuable tools in developing eco-friendly weed management systems.

6. Ecological and Evolutionary Perspectives

The presence of secondary metabolites in plants is not a random occurrence but rather a product of intense evolutionary selection shaped by millions of years of biotic interactions. These compounds have diversified under co-

evolutionary pressures imposed by herbivores, pathogens, and competing plants, resulting in a vast chemical arsenal that reflects species-specific ecological niches.

From an ecological standpoint, secondary metabolites contribute to the fitness of individual plants by improving survival and reproductive success under hostile environmental conditions. For instance, plants growing in dense, species-rich habitats often invest more in allelochemicals and volatile defense compounds than those in isolated monocultures. Similarly, shade-tolerant understory plants may rely more heavily on anti-herbivore metabolites due to limited light availability for regrowth.

Evolutionarily, chemical diversity within plant populations confers resilience against pests that evolve counter-defenses. This concept, known as the "diversity-resistance hypothesis," suggests that genetic variability in secondary metabolite profiles can reduce the success of specialist herbivores or pathogens, slowing down evolutionary arms races. Plants also exhibit phenotypic plasticity in secondary metabolite production, responding dynamically to the presence or absence of biotic stress.

Studies using phylogenetics and metabolomics have revealed conserved biosynthetic pathways and gene families associated with secondary metabolism, such as terpene synthases, polyketide synthases, and cytochrome P450 monooxygenases. These gene clusters often show signatures of positive selection, indicating their adaptive value. Horizontal gene transfer and gene duplication events have also contributed to the metabolic innovation seen across plant lineages.

Thus, secondary metabolites are not merely biochemical by-products but represent ecological and evolutionary strategies that have shaped plant survival and diversification.

7. Agricultural and Biotechnological Applications

Understanding the defensive roles of plant secondary metabolites opens doors to numerous agricultural innovations. One prominent application is the development of natural pesticides and bioherbicides. Compounds like pyrethrins from *Chrysanthemum* and rotenone from *Derris* have been commercialized as botanical insecticides due to their efficacy and environmental safety compared to synthetic chemicals.

Crop varieties with enhanced secondary metabolite profiles are being bred or engineered to resist common pests and diseases. For instance, wild tomato species rich in acylsugars and alkaloids are being introgressed into commercial cultivars to improve resistance against whiteflies and aphids. Similarly, high-glucosinolate *Brassica* lines are effective in reducing soil-borne pathogens through biofumigation.

Biotechnological approaches also harness synthetic biology to produce plant secondary metabolites in microbial systems. Engineered yeast and *E. coli* strains are now capable of producing complex compounds like artemisinin (a sesquiterpene lactone used against malaria), making these compounds more accessible and sustainable.

Moreover, metabolomic screening and CRISPR-based editing of metabolic genes provide precision tools for manipulating plant chemistry, allowing targeted enhancement of specific defensive pathways without compromising growth or yield.

These applications illustrate the tremendous potential of secondary metabolites in creating sustainable, climate-resilient, and low-input agricultural systems.

8. Challenges and Future Directions

Despite their importance, the use of plant secondary metabolites in applied sciences faces several challenges. One of the primary issues is the context dependency of metabolite production. Environmental variables like temperature, light, and soil nutrients can significantly influence the quantity and composition of secondary metabolites, making it difficult to standardize their use in agriculture or pharmacology.

Additionally, many secondary metabolites show broad-spectrum bioactivity, which may unintentionally affect non-target organisms, including pollinators, soil microbes, or human consumers. Understanding and mitigating such side effects requires comprehensive ecological risk assessments and toxicological studies.

From a research perspective, the complexity of secondary metabolite biosynthetic pathways—often involving dozens of enzymes and regulatory elements—poses significant challenges for genetic engineering. Moreover, metabolic trade-offs can occur, where increased investment in defense reduces growth or reproductive output, especially under non-stress conditions.

Future research should focus on:

- Mapping the metabolic networks and regulatory circuits governing secondary metabolism using integrative omics.
- Developing tissue-specific or inducible promoters to fine-tune metabolite expression.
- Exploring synergistic interactions between different secondary metabolites and their role in microbiome shaping.
- Promoting interdisciplinary collaboration among plant scientists, ecologists, agronomists, and bioengineers.

9. Discussion

The role of plant secondary metabolites in mediating biotic interactions reflects the remarkable biochemical adaptability and evolutionary innovation of plants. As evidenced across numerous plant lineages, these metabolites provide a multi-layered defense system against herbivores, pathogens, and even competing plants. The diversity in chemical structures—ranging from simple phenolic acids to complex terpenoids—enables plants to combat a wide array of threats with both direct toxicity and indirect ecological strategies.

A consistent theme across plant-herbivore and plant-pathogen interactions is the inducibility and specificity of metabolite production. Unlike primary metabolism, secondary metabolite synthesis is tightly regulated in a context-dependent manner, often coordinated through hormonal signals like jasmonic acid and salicylic acid. This regulation allows plants to allocate resources efficiently, maintaining a balance between defense and growth.

The interplay between allelopathy and microbial mediation is particularly notable in competitive plant-plant dynamics. Allelochemicals not only directly inhibit competitors but also reshape rhizosphere communities, potentially altering nutrient availability and plant succession. These interactions underline the ecological intelligence embedded in plant secondary chemistry.

Furthermore, the evolutionary arms race between plants and their consumers is evident in the emergence of specialized metabolites and the corresponding adaptations in herbivores and pathogens. The diversification of biosynthetic pathways and gene families involved in secondary metabolism

supports the view that chemical innovation has been a key driver of plant speciation and resilience.

However, translating this rich chemical arsenal into agricultural applications requires deeper understanding and caution. While several secondary metabolites show promise as biopesticides or herbicidal agents, their ecological complexity and variability demand careful evaluation. Modern molecular tools and metabolomics offer unprecedented resolution for exploring these pathways, yet integration across disciplines remains essential to fully harness their potential.

10. Conclusion

Plant secondary metabolites are indispensable to survival and success in biotic environments. They act not only as defensive chemicals but as regulators, messengers, and competitors—integral to the ecological and evolutionary strategies of plants. Their functions span multiple biological levels, influencing interactions from cellular defenses to ecosystem dynamics.

As global agriculture faces increasing pressure from climate change, pest resistance, and the need for sustainability, leveraging the power of these natural compounds offers a promising path forward. By integrating traditional ecological knowledge with modern biotechnological tools, we can design next-generation crops that are both productive and ecologically intelligent. Continued exploration into the complex world of secondary metabolites is not only a scientific necessity but a step toward more resilient agroecosystems.

10. References

1. Wink M. Evolution of secondary metabolites from an ecological and molecular phylogenetic perspective. *Phytochemistry*. 2003;64(1):3-19.
2. Mithöfer A, Boland W. Plant defense against herbivores: chemical aspects. *Annu Rev Plant Biol*. 2012;63:431-450.
3. Piasecka A, Jedrzejczak-Rey N, Bednarek P. Secondary metabolites in plant innate immunity: conserved function of divergent chemicals. *New Phytol*. 2015;206(3):948-964.
4. Züst T, Agrawal AA. Trade-offs between plant growth and defense against insect herbivory: an emerging mechanistic synthesis. *Annu Rev Plant Biol*. 2017;68:513-534.
5. War AR, Paulraj MG, War MY, Ignacimuthu S. Role of plant volatiles in defence against herbivores. *Arthropod-Plant Interact*. 2011;5(3):179-191.
6. Macías FA, Molinillo JM, Varela RM, Galindo JC. Allelopathy—a natural alternative for weed control. *Pest Manag Sci*. 2007;63(4):327-348.
7. Gershenzon J, Dudareva N. The function of terpene natural products in the natural world. *Nat Chem Biol*. 2007;3(7):408-14.
8. Verpoorte R, van der Heijden R, Memelink J. Engineering the plant cell factory for secondary metabolite production. *Transgenic Res*. 2000;9(6):323-43.
9. Zhang H, Mallik AU, Zeng RS. Control of natural vegetation by allelopathy. *Allelopathy J*. 2010;25(1):57-70.
10. Erb M, Kliebenstein DJ. Plant secondary metabolites as defenses, regulators, and primary metabolites: the blurred functional trichotomy. *Plant Physiol*. 2020;184(1):39-52.