



ISSN: 2456-2912

NAAS Rating (2025): 4.61

VET 2025; 10(11): 228-230

© 2025 VET

www.veterinarypaper.com

Received: 15-09-2025

Accepted: 21-10-2025

Dr. R Mahaprabhu

Ph.D., Department of Veterinary
Pathology, Veterinary College and
Research Institute, Tirunelveli,
Tamil Nadu, India

Dr. E Madhesh

Ph.D., Department of Veterinary
Medicine, Veterinary College and
Research Institute, Tirunelveli,
Tamil Nadu, India

E Aishwarya Lakshmi

Research Scholar, Department of
Veterinary, Veterinary College and
Research Institute, Tirunelveli,
Tamil Nadu, India

J Antony Pearlsy Roy

Research Scholar, Department of
Veterinary, Veterinary College and
Research Institute, Tirunelveli,
Tamil Nadu, India

A case of lymphadenopathic theileriosis in a five-months pregnant jersey cross-bred: Clinical, cytological and therapeutic observations

R Mahaprabhu, E Madhesh, E Aishwarya Lakshmi and J Antony Pearlsy Roy

DOI: <https://www.doi.org/10.22271/veterinary.2025.v10.i11d.2732>

Abstract

A five-months pregnant Jersey cross-bred was presented to the Teaching Veterinary Clinical Complex, Veterinary College and Research Institute, Tirunelveli, with enlarged superficial lymph nodes, reduced feed intake, hind-limb weakness with tremors, and mucus-coated faeces. One week earlier, the cow had been treated by a field veterinarian for recumbency and weakness. Cytological examination of a lymph node aspirate revealed lymphocytes and monocytes containing characteristic Koch's blue bodies along with eosinophils. A diagnosis of the lymphoproliferative form of bovine tropical theileriosis caused by *Theileria annulata* was established. The animal was treated with oxytetracycline and supportive therapy, leading to complete recovery. This case highlights the diagnostic value of cytology for rapid identification of *T. annulata* infection and demonstrates the effectiveness of timely therapy in pregnant Jersey cross-bred cattle.

Keywords: Cross-bred Jersey, *Theileria annulata*, Koch's blue bodies, lymphadenopathy, cytology

Introduction

Tropical theileriosis of cattle, caused by *Theileria annulata*, is a tick-borne haemoprotozoan disease of major economic importance in tropical and subtropical regions (Gharbi *et al.*, 2011) [3]. The schizont stage of *T. annulata* invades bovine lymphocytes and macrophages, producing the characteristic "Koch's blue bodies" visible in cytological or histological smears (Gupta *et al.*, 2016) [1]. These schizonts induce lymphocytic proliferation, leading to lymph node enlargement, anaemia, emaciation, and weakness (OIE, 2019) [6].

Cross-bred and exotic breeds such as Jersey and Holstein-Friesian are highly susceptible and often exhibit severe clinical manifestations (Al-Hosary *et al.*, 2018) [2]. Early diagnosis and prompt administration of specific antiprotozoal therapy are crucial for a favourable prognosis (Gill *et al.*, 2015) [4].

This paper reports a case of lymphadenopathic theileriosis in a five-months pregnant Jersey cross-bred, emphasizing cytological diagnosis and successful therapeutic management.

Materials and Methods

Animal and History

A five-months pregnant Jersey cross-bred was presented to the Teaching Veterinary Clinical Complex, Tirunelveli, with a history of gradual enlargement of superficial lymph nodes (prescapular and prefemoral), decreased feed intake, hind-limb weakness with tremors, and passage of soft faeces coated with mucus.

Clinical Examination

On clinical examination, the animal appeared dull and depressed with poor body condition. The superficial lymph nodes were enlarged, firm, and movable. The hind limbs showed weakness and fine tremors. The mucous membranes were pale, and the faeces were soft and coated with mucus, without visible blood. Rectal temperature remained within normal limits.

Corresponding Author:

Dr. R Mahaprabhu

Ph.D., Department of Veterinary
Pathology, Veterinary College and
Research Institute, Tirunelveli,
Tamil Nadu, India

Cytological Examination

Fine-needle aspiration cytology (FNAC) was performed on the enlarged prescapular lymph node using a 22-gauge needle. Smears were air-dried, fixed in methanol, stained with Leishman stain, and examined microscopically. The smears revealed numerous large lymphocytes and occasional

lymphoblasts containing intracytoplasmic, blue-staining schizont inclusion bodies (Koch's blue bodies), (Figure 1) (Figure 3), along with 5-10% eosinophils and a few monocytes (Figure 2). These findings were diagnostic of lymphoproliferative theileriosis (Gupta *et al.*, 2016) ^[1].




Fig 1: Lymphnode cytology-Lymphoblasts containing intracytoplasmic, schizont inclusion bodies (Koch's blue bodies) along with eosinophils




Fig 2: Lymphnode cytology-cytoplasm of monocyte containing schizont inclusion




Fig 3: Lymphnode cytology-Lymphoblasts containing intracytoplasmic, Blue stained round schizont inclusion bodies (Koch's blue bodies)



Fig 4: Cow-Fully recovered in standing posture with normal prescapular lymph node

Therapy

The cow was treated as follows

- **Oxytetracycline:** @ 10 mg/kg intravenously once daily for 5 days
- **Supportive therapy**
 - Injection Chlorpheniramine maleate, 12.5 mL intramuscularly
 - Injection Meloxicam, 25 mL intramuscularly
 - Injection Tribivet, 20 mL intramuscularly
 - Bolus *Ecotas* (probiotic and rumen supplement) for 4 days

Follow-up

The animal was monitored daily for improvement in feed intake, lymph node size, and reduction in tremors. A re-examination was conducted on Day 5 post-treatment.

Results

On presentation, the prescapular lymph nodes measured approximately 8 cm in diameter. The animal exhibited marked anorexia and reluctance to walk. Faecal material was soft and covered with mucus.

Cytological examination confirmed *Theileria annulata* infection by the presence of Koch's blue bodies in

lymphocytes.

By Day 4, feed intake improved, and hind-limb tremors subsided. By Day 5, the lymph nodes had regressed significantly, and faeces returned to normal consistency. The animal was clinically normal (Figure 4), indicating complete recovery. No adverse drug reactions were observed, and pregnancy was maintained uneventfully.

Discussion

The clinical signs of lymphadenopathy, anorexia, weakness, and tremors in this case correspond with previous reports of bovine tropical theileriosis (Gharbi *et al.*, 2011; Gupta *et al.*, 2016) ^[3, 1]. The cytological detection of Koch's blue bodies within lymphocytes is pathognomonic for *Theileria annulata* infection (Al-Hosary *et al.*, 2018; OIE, 2019) ^[2, 6]. These findings reaffirm the diagnostic value of FNAC as a rapid, field-applicable test in endemic regions.

The use of oxytetracycline, along with supportive therapy, likely minimized secondary bacterial infections and improved systemic resilience. Cross-bred cattle, particularly Jerseys, are known to be more susceptible to theileriosis due to limited tick resistance and higher physiological stress during pregnancy (Durrani *et al.*, 2011) ^[5]. Therefore, theileriosis should be considered a differential diagnosis in pregnant

cross-bred heifers presenting with lymphadenopathy and weakness.

Conclusion

Cytological identification of Koch's blue bodies in lymphocytes provides a rapid and reliable diagnosis of *Theileria annulata* infection. Early treatment with oxytetracycline and supportive care led to complete recovery in this five-months pregnant Jersey cross-bred heifer. Early detection and timely intervention remain vital for the successful management of bovine tropical theileriosis in cross-bred cattle under field conditions.

Acknowledgements

The authors acknowledge the facilities and assistance provided by the Teaching Veterinary Clinical Complex, Veterinary College and Research Institute, Tirunelveli.

Conflict of Interest

Not available

Financial Support

Not available

Reference

1. Gupta A, Gupta K, Leishangthem G, Bal MS, Sood NK, Singh A. Molecular and pathological studies on natural cases of bovine theileriosis. J Parasit Dis. 2016;41(1):211-218.
2. Al-Hosary AAT, Nordengrahn A, Merza M. New approach to use blood smears for diagnosis of bovine theileriosis. Indian J Anim Res. 2018;54(11):1438-1440.
3. Gharbi M, Sassi L, Dorchies P, Darghouth MA. Diagnosis and control of bovine theileriosis in North Africa. Vet Parasitol. 2011;182(1):1-9.
4. Gill BS, Kaur P, Singh A, Sood NK. Therapeutic efficacy of buparvaquone against tropical theileriosis in crossbred cattle. Indian Vet J. 2015;92(3):72-75.
5. Durrani AZ, Kamal N, Mehmood N, Shakoori AR. Incidence of theileriosis in crossbred and exotic cattle. Pak Vet J. 2011;31(2):125-127.
6. OIE (World Organisation for Animal Health). Manual of Diagnostic Tests and Vaccines for Terrestrial Animals: Theileriosis. Paris: OIE; 2019.

How to Cite This Article

Mahaprabhu R, Madhesh E, Lakshmi EA, Roy JAP. A case of lymphadenopathic theileriosis in a five-months pregnant jersey cross-bred: Clinical, cytological and therapeutic observations. International Journal of Veterinary Sciences and Animal Husbandry. 2025;10(11):228-230.

Creative Commons (CC) License

This is an open-access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.