

International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
NAAS Rating (2025): 5.29
IJABR 2025; SP-9(10): 1884-1896
www.biochemjournal.com
Received: 12-07-2025
Accepted: 15-08-2025

Bontha Rajasekar
Assistant Professor, College of
Horticulture, Mojerla,
SKLTGHU, Wanaparthy
District, Telangana, India

Dr. Purnima Mishra
Associate Professor, College of
Horticulture, Mojerla,
SKLTGHU, Wanaparthy
District, Telangana, India

Dr. K Venkata Laxmi
Associate Professor, College of
Horticulture, Malyala,
SKLTGHU, Mahabubabad
District, Telangana, India

T Navya Swetha
Assistant Professor, College of
Horticulture, Mojerla,
SKLTGHU, Wanaparthy
District, Telangana, India

T Baby Rani
Assistant Professor, College of
Horticulture, Mojerla,
SKLTGHU, Wanaparthy
District, Telangana, India

Corresponding Author:
Bontha Rajasekar
Assistant Professor, College of
Horticulture, Mojerla,
SKLTGHU, Wanaparthy
District, Telangana, India

A review on mode of action of insecticides

Bontha Rajasekar, Purnima Mishra, K Venkata Laxmi, T Navya Swetha and T Baby Rani

DOI: <https://www.doi.org/10.33545/26174693.2025.v9.i10Sw.6112>

Abstract

The development of novel insecticides that promises efficient and environmentally sustainable solutions. Most current insecticides act on nerve and muscle targets. Insecticides that act on these targets are generally fast acting. Insect development is controlled by juvenile hormone and ecdysone, by direct perturbing cuticle formation/deposition or lipid biosynthesis. Such insect growth regulators are generally slow to moderately slow acting. Several insecticides are known to interfere with mitochondrial respiration by the inhibition of electron transport and / or oxidative phosphorylation. Such insecticides are generally fast to moderately fast acting. Lepidopteran specific microbial toxins that are sprayed or expressed in transgenic crop varieties. New pesticides can be designed to target specific pests while minimizing harm to beneficial organisms, reducing ecological disruptions. Modern pesticides are formulated to have lower toxicity and persistence in the environment, resulting in less contamination of water, soil, and air. New pesticides may leave lower residue levels on crops, addressing concerns about food safety and consumer health. New pesticides with different modes of action can help manage resistance issues that have developed in pests against older pesticides. the mode of action of novel insecticides an homage to the power of scientific inquiry and the promise of a future where human ingenuity, coupled with nature's wisdom, charts a course towards a more balanced and prosperous world.

Keywords: Novel insecticides, toxicity, environment, safety and health

Introduction

The ongoing battle against insect pests has spurred the development of novel insecticides that promise efficient and environmentally sustainable solutions. This review delves into the mode of action of these innovative compounds, shedding light on the mechanisms that set them apart from traditional pesticides. Unlike conventional insecticides that often act broadly, novel insecticides target specific vulnerabilities within insect physiology. By leveraging a diverse array of molecular and biochemical pathways, these compounds disrupt key biological functions critical for insect survival. This precision in targeting not only maximizes efficacy but also minimizes ecological impact. One of the groundbreaking approaches is the utilization of insect-specific pheromones and behavior-modifying compounds. These disrupt communication and mating patterns, leading to reduced reproductive success and population decline. Similarly, botanical extracts and natural compounds interfere with insect growth and development, capitalizing on their specific interactions with hormonal pathways. Furthermore, advanced genetic and molecular techniques have given rise to genetically modified crops expressing insecticidal proteins. These proteins, derived from bacteria like *Bacillus thuringiensis* (Bt), selectively target specific insect groups, sparing beneficial organisms and reducing chemical usage. RNA interference (RNAi) technology takes this step further, silencing essential genes through the application of small RNA molecules.

Understanding the mode of action of these novel insecticides involves a convergence of disciplines, including molecular biology, entomology, and biochemistry. Through innovative techniques such as proteomics, genomics, and metabolomics, researchers gain insights into the intricate interactions between insecticides and their molecular targets. Computational modeling aids in predicting insecticide-target interactions and potential resistance mechanisms. Thus unraveling the mode of action of novel insecticides marks a pivotal advancement in pest management.

These compounds represent a paradigm shift towards targeted and sustainable approaches, reducing the ecological footprint associated with traditional pesticides. By comprehending the mechanisms underpinning their effectiveness, researchers pave the way for the development of innovative strategies that can mitigate pest-related challenges while safeguarding ecosystems.

Classification of Insecticides by their Actions: Modes of action are according to the physiological functions affected.

Nerve and Muscle

Most current insecticides act on nerve and muscle targets. Insecticides that act on these targets are generally fast acting.

Group No	Name	MoA	Example
1	Acetylcholinesterase (AChE) Inhibitor	Inhibit AChE, cause hyperexcitation.	Carbamates, Organophosphates
2	GABA Gated Chloride Channel Blockers	Block the GABA-activated chloride channel, causing hyperexcitation and convulsions.	Cyclodiene Organochlorines, Phenylpyrazoles (Fiproles)
3	Sodium Channel Modulators	Keep Sodium channel open, causing hyperexcitation and, in some cases, nerve block.	Pyrethroids, Pyrethrins, DDT, Methoxychlor
4	Nicotinic Acetylcholine Receptor (nAChR) Competitive Modulators	Bind to acetylcholine site on nAChRs causing a range of symptoms from hyper-excitation to lethargy and paralysis.	Neonicotinoids, Nicotine, Sulfoximines, Butenolides, Mesoinics, Pyridilidenes
5	Nicotinic Acetylcholine Receptor (nAChR) Allosteric Modulator - I	Allosterically activate nAChRs, causing hyperexcitation of the nervous system.	Spinosyns
6	Glutamate-Gated Chloride Channel (GluCl) Allosteric Modulators	Allosterically activate glutamate-gated chloride channels (GluCls), causing paralysis.	Avermectins, Milbemycins
7	Chordotonal Organ TRPV Channel Modulators	Bind to and disrupt the gating of Nan-Iav TRPV (Transient Receptor Potential Vanilloid) Channel complexes in chordotonal stretch receptor organs, which are critical for the senses of hearing, gravity, balance, acceleration, proprioception, and kinaesthesia.	B. Pyridine Azomethine Derivatives, D. Pyropenes
8	Nicotinic Acetylcholine Receptor (nAChR) Channel Blockers	Block the nAChR ion channel, resulting in nervous system block and paralysis.	Nereistoxin Analogues

Group No	Name	MoA	Example
9	Octopamine Receptor Agonists	Activate octopamine receptors, leading to hyperexcitation.	Amitraz
10	Voltage Dependent Sodium Channel Blockers	Block sodium channels, causing nervous system shutdown and paralysis.	Oxadiazines, Semicarbazones
11	Ryanodine Receptor Modulators	Activate muscle ryanodine receptors, leading to contraction and paralysis.	Diamides
12	Chordotonal Organ Nicotinamidase Inhibitors	Disrupt the function of chordotonal stretch receptor organs, which are critical for the senses of hearing, gravity, balance, acceleration, proprioception, and kinesthesia.	Flonicamid
13	GABA-Gated Chloride Channel Allosteric Modulators	Allosterically inhibit the GABA-activated chloride channel, causing hyperexcitation and convulsions.	Meta-Diamides, Isoxazolines
14	Nicotinic AChR Allosteric Modulators Site - II	Allosterically activate nAChRs (at a site distinct from Group 5 - Site I), causing hyperexcitation of the nervous system.	GS-omega/kappa HXTX-Hv1a Peptide
15	Calcium-Activated Potassium Channel (KCA2) Modulators	Negative modulation of KCA2 causes hyperexcitation and convulsions. KCA2 channels are activated by an increase in intracellular calcium concentration and are involved in the regulation of action potential.	Acynonapyr

Growth: Insect development is controlled by juvenile hormone and ecdysone, by direct perturbing cuticle formation/deposition or lipid biosynthesis. Such insect growth regulators are generally slow to moderately slow acting.

Group No	Name	MoA	Example
1	Juvenile Hormone Mimics	Applied in the Pre-metamorphic instar, these compounds disrupt and prevent metamorphosis.	JH Analogus Fenoxycarb Pyriproxylen
2	Mite Growth Inhibitors affecting CHS 1	Inhibit the enzyme that catalyzes the polymerization of chitin	Clofentezine, Diflovidazin, Hexythiazox Etoxazole
3	Inhibitors of Chitin Biosynthesis affecting CHS 1	Inhibit the enzyme that catalyzes the polymerization of chitin	Benzoylureas
4	Inhibitors of Chitin Biosynthesis affecting Type 1	Incompletely defined MoA leading to inhibition of chitin biosynthesis in a number of insects, including Whiteflies	Buprofezin
5	Moulting Disruptor (Dipteran)	Incompletely defined MoA leads to Moulting disruption.	Cyromazine
6	Ecdysone receptor Agonist	Mimic the moulting hormone, ecdysone, including a precocious moult.	Diacylhydrazines
7	Inhibitor of Acetyl CoA Carboxylase	Inhibit acetyl Coenzyme A carboxylase, part of first step in lipid biosynthesis, leading to insect death.	Tetronic and Tetramic Acid Derivatives

Respiration: Several insecticides are known to interfere with mitochondrial respiration by the inhibition of electron

transport and / or oxidative phosphorylation. Such insecticides are generally fast to moderately fast acting.

Group No	Name	MoA	Example
1	Inhibitors of Mitochondrial	Inhibit the enzyme that synthesizes ATP.	Diafenthiuron Organotin Miticids Propargite Tetradifon
2	Uncouples of Oxidative Phosphorylation via disruption of Proton Gradient	Protonophores that short-circuit the mitochondrial proton gradient so that ATP cannot be synthesized.	Pyrroles Dinitrophenols Sulfuramid
3	Mitochondrial Complex III ET Inhibitors - QO Site	Inhibit electron transport complex III, preventing the utilization of energy by cells.	Hydramethylnon Acequinocyl Fluacrypyrin Bifenazate
4	Mitochondrial Complex I ET Inhibitors	Inhibit electron transport complex I, preventing the utilization of energy by cells.	Meti acaricides & Insecticides Rotenone
5	Mitochondrial Complex IV ET Inhibitors	Inhibit electron transport complex IV, preventing the utilization of energy by cells.	Phosphids Cyanides
6	Mitochondrial Complex II ET Inhibitors	Inhibit electron transport complex II, preventing the utilization of energy by cells.	Beta - Keto nitrile Derivatives Carboxanilides
7	Mitochondrial Complex III ET Inhibitors - QI Site	Inhibit electron transport complex III, preventing the utilization of energy by cells. In contrast to Group 20, Group 34 insecticides bind to the QI Site.	Flometoquin

Midgut: Lepidopteran specific microbial toxins that are sprayed or expressed in transgenic crop varieties.

Group No	Name	MoA	Example
1	Microbial Disruptors of Insect Midgut Membranes	Protein toxins that bind to receptors on the midgut membrane and induce pore formation on, resulting in ionic imbalance and septicaemia.	B.t& Insectidal Bacillus sphaericus
2	Baculoviruses	A Baculovirus - unique Per OS Infectivity Factor (PIF) protein complex on the virus promotes host- specific infection by binding to PIF targets on midgut cells that are unknown but believed to be unique for each baculovirus type. Infection is ultimately lethal	Gv, NPV

Reason for Pesticide Group Change

New pesticides are developed for several reasons, even though older pesticides already exist. Here are some reasons for the development and introduction of new pesticides:

- 1. Resistance Management:** Over time, pests can develop resistance to older pesticides, rendering them ineffective. New pesticides may be designed with different modes of action to target resistant pest populations.
- 2. Environmental Impact:** Newer pesticides are often formulated to be more environmentally friendly, with lower toxicity to nontarget organisms and reduced persistence in the environment.
- 3. Regulatory Compliance:** As regulations and safety standards evolve, older pesticides that do not meet current requirements may be phased out or restricted. New pesticides are developed to comply with these updated regulations.
- 4. Targeted Specificity:** Modern pesticides can be designed to target specific pests while causing minimal harm to beneficial organisms. This specificity reduces unintended consequences in the ecosystem.
- 5. Improved Efficacy:** New pesticides might offer better efficacy in controlling pests, requiring lower application rates or fewer applications to achieve desired results.
- 6. Novel Formulations:** Advances in formulation technology allow for the development of pesticides with improved delivery methods, such as controlled release, which can enhance their effectiveness.
- 7. Reduced Residue Levels:** New pesticides may be formulated to leave lower residue levels on crops, addressing concerns about food safety and consumer health.

8. Non-Chemical Alternatives: In response to consumer demand for more sustainable practices, new pesticides may include biopesticides, botanical extracts, or microbial agents that have lower environmental impacts.

9. Emerging Pest Threats: New pests or pest populations that were not previously a problem might emerge due to changing climate conditions, globalization, or other factors. New pesticides could be developed to address these new threats.

10. Profitability and Market Competition: Developing new pesticides can be financially lucrative for companies in the agrochemical industry. Competition and potential for improved profits drive investment in research and development.

It's important to note that the introduction of new pesticides should be approached cautiously, considering potential risks to human health, non-target organisms, and the environment. Integrated Pest Management (IPM) approaches that combine multiple strategies, including biological control, cultural practices, and chemical control when necessary, are increasingly favored to create a balanced and sustainable pest management strategy.

New pesticides often offer several advantages over old pesticides

- 1. Targeted Specificity:** New pesticides can be designed to target specific pests while minimizing harm to beneficial organisms, reducing ecological disruptions.
- 2. Reduced Environmental Impact:** Modern pesticides are formulated to have lower toxicity and persistence in the environment, resulting in less contamination of water, soil, and air.

3. **Lower Residue Levels:** New pesticides may leave lower residue levels on crops, addressing concerns about food safety and consumer health.
4. **Efficient Application:** Advances in formulation technology allow for more efficient delivery methods, reducing the amount of pesticide needed for effective control.
5. **Resistance Management:** New pesticides with different modes of action can help manage resistance issues that have developed in pests against older pesticides.
6. **Compliance with Regulations:** New pesticides are developed to meet updated safety and regulatory standards, ensuring they are in line with current legal requirements.
7. **Enhanced Efficacy:** Improved formulations and design can result in higher efficacy against target pests, requiring fewer applications.
8. **Reduced Non-Target Effects:** New pesticides are designed to minimize harm to non-target organisms, helping to maintain biodiversity and ecological balance.
9. **Innovation:** Advances in science and technology lead to the development of novel pesticide compounds and delivery systems, fostering innovation in pest management.
10. **Public Perception:** New pesticides, especially those with reduced environmental impact, are often more socially acceptable due to their safer profiles.
11. **Long-Term Sustainability:** By incorporating newer, more targeted pesticides into Integrated Pest Management (IPM) strategies, agricultural systems can become more sustainable and resilient.
12. **Alternative Modes of Action:** Some new pesticides utilize different biochemical pathways to control pests, reducing the risk of cross-resistance with older pesticides.
13. **Emergency Situations:** In cases of emerging pest threats or sudden outbreaks, new pesticides can provide rapid solutions to mitigate economic losses.
14. **Flexibility:** The diversity of available pesticides, old and new, allows for the selection of products that best match specific pest challenges and environmental conditions.
15. **Economic Benefits:** While newer pesticides may have higher initial costs, their efficiency, reduced application rates, and improved crop yields can lead to cost savings in the long run.

It's worth noting that while new pesticides offer these advantages, their use should be guided by responsible and sustainable pest management practices. Integrated approaches that combine multiple strategies and focus on long-term solutions are increasingly favored for maintaining agricultural productivity while safeguarding environmental health.

Macrocyclic Lactone Insecticides

Macrocyclic lactone insecticides are a class of chemicals that target insects' nervous systems, affecting their ability to transmit nerve impulses. They work by binding to specific receptors in the insect nervous system, disrupting the normal flow of neurotransmitters like gamma-aminobutyric acid (GABA) and glutamate. This disruption leads to overstimulation of nerve cells, resulting in paralysis and

eventually death for the insect. One well-known example of a macrocyclic lactone insecticide is ivermectin. It binds to GABA receptors and glutamate-gated chloride channels in nerve cells. This binding enhances the inhibitory effects of GABA, a neurotransmitter that usually dampens nerve activity. As a result, nerve cells become hyperpolarized and less responsive to stimuli, leading to paralysis and death of the insect.

Macrocyclic lactone insecticides primarily target insects due to the unique structure and function of their nervous systems. Mammals, including humans, have different types of receptors and ion channels, making them less susceptible to the effects of these insecticides. However, it's important to note that accidental exposure to high doses of macrocyclic lactone insecticides could still be harmful to mammals and other non-target organisms.

Avermectin Insecticides

Avermectins are a subgroup of macrocyclic lactone insecticides that share a similar mode of action. One of the most well-known avermectins is ivermectin. The mode of action of avermectin insecticides involves targeting the nervous systems of insects and other invertebrates. Avermectins, like ivermectin, work by binding to specific receptors in the nerve cells of insects and parasites. These receptors are called glutamate-gated chloride channels (GluCl channels). When avermectins bind to these receptors, they enhance the effects of the neurotransmitter gamma-aminobutyric acid (GABA), which is an inhibitory neurotransmitter. This results in increased chloride ion influx into nerve cells through the GluCl channels, leading to hyperpolarization of the cells and making them less responsive to stimuli. The hyperpolarization caused by the binding of avermectins to GluCl channels leads to paralysis and disruption of the nervous system in insects and parasites. This ultimately results in their death. Importantly, the mode of action is selective for invertebrates because mammals, including humans, have a different type of GABA receptor that is less sensitive to avermectins. This selectivity makes avermectin insecticides effective against pests while minimizing harm to mammals. Avermectins are used to control a wide range of pests, including nematodes and arthropods, and they have been particularly valuable in combating parasites in livestock and pets.

Milbemycin Insecticides

Milbemycin is an anthelmintic compound used to control internal parasites in livestock and pets. While it is not a traditional insecticide, its mode of action is worth discussing. Milbemycin primarily targets nematode parasites, which are a type of roundworm that can infect animals. Its mode of action involves interfering with the nematode's neurotransmission system, specifically by affecting acetylcholine receptors. Acetylcholine is a neurotransmitter that plays a crucial role in transmitting nerve signals in both animals and parasites. Milbemycin binds to acetylcholine receptors on the nematode's nerve cells, disrupting the normal transmission of nerve impulses. This interference leads to paralysis of the nematode's muscles and inhibits its ability to maintain its position within the host animal's digestive tract. As a result, the nematodes are expelled from the host's body through natural digestive processes. It's important to note that milbemycin's mode of action is

specific to nematode parasites and does not impact insects or other types of pests. Additionally, mibemycin is generally considered safe for use in animals when administered at appropriate doses. However, as with any medication, proper dosing and administration are critical to ensure the health and safety of the treated animals.

Spinosyn Insecticides

Spinosyn insecticides are derived from naturally occurring compounds produced by the bacterium *Saccharopolyspora spinosa*. They are used to control a wide range of insect pests in agriculture and horticulture. The mode of action of spinosyn insecticides involves two main components: spinosyn A and spinosyn D. Spinosyns act on the nervous systems of insects by binding to specific sites on nicotinic acetylcholine receptors (nAChRs) found in nerve cells. These receptors play a key role in transmitting nerve impulses by regulating the flow of ions across the cell membrane. Spinosyns target nAChRs located in the insect's central nervous system. Upon binding to nAChRs, spinosyns disrupt the normal functioning of the receptors, leading to the prolonged opening of ion channels. This results in the excessive flow of ions, primarily sodium, into the nerve cells. The increased ion flow overstimulates the nerve cells, causing continuous nerve signaling and ultimately leading to paralysis and death of the insect. One notable aspect of spinosyn insecticides is their selectivity for insects over mammals and other non-target organisms. This selectivity is due to differences in the specific types and subtypes of nAChRs present in insects versus mammals. Spinosyns have a lower affinity for mammalian nAChRs, which reduces the risk of toxic effects in animals and humans. Overall, the mode of action of spinosyn insecticides as nicotinic acetylcholine receptor agonists, leading to neural overstimulation and insect paralysis, has contributed to their success as effective and relatively safe pest control agents.

Oxadiazine Insecticides

Oxadiazine insecticides, such as indoxacarb, belong to a class of chemicals that target insects' nervous systems through a unique mode of action. The mode of action of indoxacarb involves its conversion into an active metabolite within the insect's body. When indoxacarb is ingested or comes into contact with an insect, it is metabolized by enzymes in the insect's gut to produce its active form, which acts on voltage-gated sodium channels in nerve cells. These sodium channels play a critical role in transmitting nerve impulses by allowing the flow of sodium ions into nerve cells. This is an essential step for nerve cell depolarization and the propagation of nerve signals. Indoxacarb's active metabolite binds specifically to voltage-gated sodium channels in a way that keeps these channels open for a longer period than normal. This leads to a continuous influx of sodium ions into nerve cells and excessive nerve stimulation. The sustained depolarization of nerve cells ultimately disrupts normal nerve function, causing paralysis and death in the insect. Importantly, the mode of action of indoxacarb is selective for insects because the specific sodium channel subtype it targets is more prevalent in insects compared to mammals. This selective action reduces the potential impact on nontarget organisms, including humans and animals. In summary, oxadiazine insecticides like indoxacarb exert their mode of action by interfering

with the normal functioning of voltage-gated sodium channels in insect nerve cells, leading to paralysis and death while minimizing harm to non-target organisms.

Pyrrole Insecticides

Pyrrole insecticides, including chlorfenapyr, have a unique mode of action that involves disrupting the energy production process within insect cells. The mode of action of chlorfenapyr can be summarized as follows:

- 1. Pro-insecticide Activation:** Chlorfenapyr itself is not toxic to insects. However, once it is ingested or absorbed by the insect, it undergoes a two-step process to convert it into its active form. Enzymes in the insect's body metabolize chlorfenapyr into a compound that disrupts cellular respiration.
- 2. Mitochondrial Uncoupling:** The active form of chlorfenapyr affects the insect's mitochondria, which are the cellular organelles responsible for producing energy in the form of adenosine triphosphate (ATP). Chlorfenapyr uncouples oxidative phosphorylation, a process that couples the electron transport chain to ATP synthesis. As a result of uncoupling, the electron transport chain continues to operate, but ATP production is significantly reduced.
- 3. Energy Depletion:** With reduced ATP production, the insect's cells gradually run out of energy. This leads to an inability to carry out essential physiological processes, including muscle movement and normal cellular function.
- 4. Delayed Toxicity:** One distinctive feature of pyrrole insecticides like chlorfenapyr is their delayed toxicity. Insects may not show signs of distress immediately after exposure. Instead, they continue to function for a period while energy reserves are depleted. This delayed effect can make chlorfenapyr particularly effective against pests that may continue to feed and spread the insecticide within the colony before succumbing to its effects.
- 5. Impact on Multiple Pest Types:** Chlorfenapyr's mode of action makes it effective against a wide range of insect pests, including those that have developed resistance to other classes of insecticides. It is commonly used in agriculture and urban pest management. In summary, the mode of action of chlorfenapyr involves disrupting cellular energy production in insects, leading to a gradual depletion of energy reserves and eventual paralysis and death. The delayed toxic effect and broad spectrum of activity make it a valuable tool in integrated pest management strategies.

Pyrazole Insecticides

Pyrazole insecticides target insects' nervous systems by affecting neurotransmitter regulation. The mode of action of pyrazole insecticides involves blocking the activity of an enzyme called acetylcholinesterase (AChE), which plays a crucial role in the transmission of nerve impulses. Here's how the mode of action typically works:

- 1. Neurotransmitter Regulation:** Insects use a neurotransmitter called acetylcholine to transmit signals between nerve cells (neurons). After acetylcholine is released from a neuron, it needs to be rapidly broken down to stop the nerve signal and allow the neuron to reset for the next transmission.

2. Role of Acetylcholinesterase (AChE)

Acetylcholinesterase is the enzyme responsible for breaking down acetylcholine in the synapses (junctions between neurons). This enzyme ensures that nerve impulses are terminated and that neurons can transmit new signals.

3. **Inhibition of AChE:** Pyrazole insecticides inhibit the activity of acetylcholinesterase. When AChE is inhibited, acetylcholine accumulates in the synapses, leading to continuous nerve signaling. This overstimulation of nerve cells can result in paralysis and eventual death of the insect.

Impact on Nervous System: As the insect's nervous system becomes increasingly disrupted due to the continuous nerve signaling, it loses the ability to coordinate movement and perform essential functions. This leads to paralysis and ultimately death. The specific mechanism of action of pyrazole insecticides on acetylcholinesterase makes them effective against a variety of insect pests. However, it's important to note that their mode of action is similar to that of other classes of insecticides that target acetylcholinesterase. This can potentially lead to cross-resistance, where insects resistant to one class of acetylcholinesterase inhibitors may also be resistant to pyrazole insecticides. It's also worth mentioning that while pyrazole insecticides are effective against many insect pests, their use should be managed carefully to minimize impacts on non-target organisms and to avoid the development of resistance.

Phenylpyrazole Insecticides

Phenylpyrazole insecticides, such as fipronil, have a mode of action that disrupts the normal functioning of insect nervous systems. Their mode of action is unique and distinct from many other classes of insecticides. Here's how phenylpyrazole insecticides typically work:

1. **GABA Receptor Targeting:** Phenylpyrazole insecticides target a specific type of receptor in the insect nervous system called the gamma-aminobutyric acid (GABA) receptor. GABA is an inhibitory neurotransmitter that regulates the transmission of nerve signals.
2. **Disruption of Chloride Flow:** GABA receptors are ion channels that allow chloride ions to flow into nerve cells. This influx of chloride ions hyperpolarizes the nerve cells, making them less excitable and reducing the likelihood of transmitting nerve impulses.
3. **Blockage of GABA Receptors:** Phenylpyrazole insecticides like fipronil bind to GABA receptors on nerve cells. However, instead of enhancing the inhibitory effects of GABA, they block the flow of chloride ions through these receptors. This disrupts the normal inhibitory signaling that GABA provides.
4. **Nerve Overstimulation:** As a result of GABA receptor blockage, nerve cells become less inhibited and more excitable. This leads to overstimulation of nerve impulses, causing erratic and uncontrolled nerve signaling.
5. **Impaired Nervous System Function:** The overstimulation of nerve cells ultimately disrupts the insect's nervous system, leading to paralysis and death. Insects affected by phenylpyrazole insecticides often

exhibit symptoms like tremors, convulsions, and uncoordinated movement.

One notable aspect of the mode of action of phenylpyrazole insecticides is their specificity for insects and their relatively low toxicity to mammals, including humans. This is because the GABA receptors in insects are structurally different from those in mammals, reducing the impact of these insecticides on non-target organisms. However, it's important to consider the potential for resistance development in insect populations due to the specific mode of action of phenylpyrazole insecticides. To manage resistance and minimize environmental impacts, integrated pest management practices and rotation of different classes of insecticides are recommended.

Pyrazole Carboxamide Insecticides

Pyrazole carboxamide insecticides are a class of chemicals that target insects' nervous systems by affecting a specific type of ion channel known as the ryanodine receptor (RyR). The mode of action of pyrazole carboxamide insecticides involves disrupting calcium regulation within muscle cells, leading to paralysis and eventual death in insects. Here's how the mode of action typically works:

1. **Calcium Regulation and Muscle Contraction:** In insects (and other animals), muscle contraction is initiated by the release of calcium ions from internal stores within muscle cells. These calcium ions bind to specific proteins, triggering the contraction of muscle fibers.
2. **Role of Ryanodine Receptors (RyRs):** Ryanodine receptors are ion channels found on the endoplasmic reticulum of muscle cells. They play a crucial role in releasing calcium ions from internal stores in response to nerve signals.
3. **Activation of RyRs:** Pyrazole carboxamide insecticides bind to and activate the ryanodine receptors in muscle cells. This leads to the excessive release of calcium ions from internal stores, overwhelming the normal regulatory mechanisms.
4. **Calcium Overload and Muscle Dysfunction:** The excess calcium ions flood the muscle cells, causing sustained and uncontrolled muscle contractions. This disrupts the insect's ability to move, fly, or perform other essential functions.
5. **Paralysis and Death:** The sustained muscle contractions and disruption of normal muscle function lead to paralysis and eventual death of the insect. This mode of action is specific to insects and is less likely to affect non-target organisms, including mammals, due to differences in the structure and function of ryanodine receptors. However, as with any class of insecticides, there is a risk of insects developing resistance over time. To manage resistance and minimize environmental impacts, it's important to use pyrazole carboxamide insecticides in combination with other integrated pest management practices.

Pyridyl Pyrazole Insecticides

Pyridyl pyrazole insecticides, such as fipronil, target insects' central nervous systems by disrupting the normal function of neurotransmitters. Their mode of action involves interacting with receptors that play a critical role in nerve signal

transmission. Here's how the mode of action typically works:

1. **Targeting GABA Receptors:** Pyridyl pyrazole insecticides specifically target a type of receptor called the gammaaminobutyric acid (GABA) receptor. GABA is an inhibitory neurotransmitter that regulates the transmission of nerve signals in the insect's nervous system.
2. **Disruption of Chloride Flow:** GABA receptors are ion channels that allow chloride ions to flow into nerve cells. This influx of chloride ions hyperpolarizes the nerve cells, making them less excitable and reducing the likelihood of transmitting nerve impulses.
3. **Blocking GABA Receptors:** Pyridyl pyrazole insecticides like fipronil bind to GABA receptors on nerve cells, but instead of enhancing the inhibitory effects of GABA, they block the flow of chloride ions through these receptors. This disrupts the normal inhibitory signaling provided by GABA.
4. **Hyperexcitation of Nerve Cells:** As a result of GABA receptor blockage, nerve cells become less inhibited and more excitable. This leads to overstimulation of nerve impulses, causing erratic and uncontrolled nerve signaling.

Nervous System Dysfunction: The overstimulation of nerve cells ultimately disrupts the insect's nervous system, leading to paralysis and death. Insects exposed to pyridylpyrazole insecticides often exhibit symptoms such as tremors, convulsions, and uncoordinated movement. It's important to note that the mode of action of pyridylpyrazole insecticides is specific to insects and has a relatively low impact on non-target organisms, including mammals. This selectivity is due to differences in the structure and function of GABA receptors between insects and mammals. However, as with any class of insecticides, there is a potential for resistance development in insect populations. To manage resistance and minimize environmental impacts, integrated pest management practices and rotation of different classes of insecticides are recommended.

Diamide Insecticides

Phthalic and anthranilic diamides are two classes of insecticides that share a common mode of action, which involves affecting the ryanodine receptors (RyRs) in insect muscle cells. This disruption of calcium regulation within muscles leads to paralysis and death in insects. Here's how the mode of action of phthalic and anthranilic diamides typically works:

1. **Calcium Regulation and Muscle Contraction:** In insects (and other animals), muscle contraction is initiated by the release of calcium ions from internal stores within muscle cells. Calcium ions play a critical role in the interaction between actin and myosin, the proteins responsible for muscle contraction.
2. **Role of Ryanodine Receptors (RyRs):** Ryanodine receptors are ion channels located on the sarcoplasmic reticulum of muscle cells. These receptors are responsible for releasing calcium ions from internal stores in response to nerve signals.
3. **Activation of RyRs:** Phthalic and anthranilic diamides bind to and activate the ryanodine receptors in muscle cells. This activation causes the release of calcium ions from the internal stores.

4. **Calcium Overload and Muscle Dysfunction:** Unlike the controlled release of calcium ions during normal muscle function, the activation of RyRs by diamides leads to an excessive and uncontrolled release of calcium ions into the muscle cells.

Muscle Paralysis and Death: The uncontrolled influx of calcium ions results in continuous and uncoordinated muscle contractions. This disrupts the insect's ability to move and carry out essential functions, ultimately leading to paralysis and death. The mode of action of phthalic and anthranilic diamides is specific to insects and has a relatively low impact on non-target organisms, including mammals, due to differences in the structure and function of ryanodine receptors. However, it's important to manage the use of these insecticides to avoid resistance development and minimize potential environmental impacts.

Ketoenol Insecticides

Ketoenol insecticides, such as tebufenozide, methoxyfenozide, and halofenozide, have a mode of action that disrupts the development of insects by interfering with their hormonal regulation. The primary target of ketoenol insecticides is the process of molting, which is crucial for the growth and development of insects. Here's how the mode of action of ketoenol insecticides typically works:

1. **Hormonal Regulation of Molting:** Insects go through a process called molting to shed their old exoskeleton and grow. This process is regulated by a hormone called ecdysone, which initiates molting by triggering the synthesis of enzymes that break down the old exoskeleton.
2. **Role of Ecdysone Receptor (EcR):** Ecdysone exerts its effects by binding to a receptor called the ecdysone receptor (EcR). When ecdysone binds to EcR, it activates a cascade of events that lead to molting and the synthesis of new cuticle (exoskeleton).
3. **Inhibition of EcR:** Ketoenol insecticides interfere with the action of ecdysone by binding to the ecdysone receptor (EcR) but not activating it. This binding prevents the normal interaction between ecdysone and EcR, disrupting the molting process.
4. **Disruption of Molting:** Without proper ecdysone signaling through the EcR, insects are unable to properly initiate molting. As a result, they cannot shed their old exoskeleton and grow into the next developmental stage.
5. **Developmental Arrest:** The disruption of molting leads to developmental arrest in insects. They become trapped in their current developmental stage and are unable to progress to the next one. This ultimately results in reduced growth, infertility, and decreased survival rates.

The mode of action of ketoenol insecticides is selective to insects due to the specific interaction between ecdysone and its receptor. Mammals and other non-target organisms have different hormonal systems, reducing the potential impact of these insecticides on noninsect species. However, as with any class of insecticides, resistance management practices and integrated pest management strategies are important to maintain their effectiveness and minimize environmental impacts.

Tetramic acid Insecticides

Tetramic acid derivatives, such as spirotetramat, are a class of insecticides that affect insects by disrupting their feeding and growth processes. The mode of action of tetramic acid derivatives involves targeting multiple physiological pathways, including those related to insect growth regulation and energy production. Here's how the mode of action of tetramic acid derivatives typically works:

- 1. Disruption of Feeding:** Tetramic acid derivatives can have a deterrent effect on feeding behavior in insects. When insects come into contact with or ingest these insecticides, they may lose interest in feeding on treated plants, which can reduce their ability to obtain essential nutrients.
- 2. Disruption of Growth Regulation:** Tetramic acid derivatives can interfere with the insect's hormonal regulation, particularly the juvenile hormone pathway. Juvenile hormone plays a key role in regulating insect development and metamorphosis. By disrupting this pathway, tetramic acid derivatives can prevent insects from progressing through their life stages.
- 3. Inhibition of Energy Production:** Some tetramic acid derivatives can interfere with the oxidative phosphorylation process within insect cells. This process is essential for producing adenosine triphosphate (ATP), which is the primary energy currency of cells. Inhibition of ATP production can lead to reduced energy availability for normal physiological processes.
- 4. Negative Impact on Reproduction:** Tetramic acid derivatives can also affect insect reproduction. By disrupting growth and energy processes, these insecticides can lead to reduced fertility and impaired reproductive success in treated insects.
- 5. Gradual Impact:** The mode of action of tetramic acid derivatives may take some time to manifest noticeable effects on insects. This gradual impact is due to the disruption of multiple physiological processes, which collectively affect the insect's overall fitness. It's important to note that the exact mode of action can vary among different tetramic acid derivatives, and each may have its own specific mechanisms of activity. Additionally, like with any class of insecticides, the development of resistance in insect populations is a concern. Integrated pest management practices, including proper rotation and use of different insecticide classes, are crucial for managing resistance and preserving the effectiveness of tetramic acid derivatives and other insecticides.

Tetronic Acid Insecticides

Tetronic acid insecticides, such as spinetoram, have a mode of action that targets insects' nervous systems by affecting the normal function of neurotransmitters. The mode of action of tetronic acid insecticides involves interacting with nicotinic acetylcholine receptors (nAChRs), which play a crucial role in nerve signal transmission. Here's how the mode of action of tetronic acid insecticides typically works:

- 1. Targeting Nicotinic Acetylcholine Receptors (nAChRs):** Nicotinic acetylcholine receptors are ion channels found on nerve cells in insects (and other animals). These receptors are involved in transmitting nerve signals by allowing the flow of ions, such as sodium and calcium, across the cell membrane.

- 2. Normal Nerve Signal Transmission:** In a normal nerve impulse transmission, the neurotransmitter acetylcholine binds to nAChRs on the postsynaptic membrane of a nerve cell. This binding triggers the opening of ion channels and the flow of ions, leading to the transmission of the nerve signal.
- 3. Interference with nAChRs:** Tetronic acid insecticides, like spinetoram, bind to nAChRs on nerve cells but do not trigger the same ion channel opening response that acetylcholine does. This binding disrupts the normal transmission of nerve signals.
- 4. Nerve Signal Disruption:** Due to the binding of tetronic acid insecticides, nerve cells are less able to transmit signals properly. This disruption leads to impaired communication between nerve cells and results in a range of effects, including paralysis and eventual death of the insect.
- 5. Selective Impact:** The mode of action of tetronic acid insecticides is more specific to insects due to differences in the structure of nAChRs between insects and mammals. This selectivity helps minimize the impact on non-target organisms. The mode of action of tetronic acid insecticides can be effective against a variety of insect pests, but as with any class of insecticides, there is a potential for the development of resistance over time.

To manage resistance and reduce potential environmental impacts, integrated pest management practices that include the rotation of different insecticide classes are recommended. Fenazaquin is an acaricide, which means it is primarily used to control mites and ticks. It belongs to the quinazoline class of compounds. The mode of action of fenazaquin involves disrupting the mitochondrial electron transport chain, leading to the impairment of energy production within the cells of mites and ticks. Here's how the mode of action of fenazaquin typically works:

- 1. Mitochondrial Electron Transport Chain:** Mitochondria are the energy-producing organelles within cells. The mitochondrial electron transport chain is a series of protein complexes that play a crucial role in generating adenosine triphosphate (ATP), the cell's energy currency.
- 2. Inhibition of Complex III:** Fenazaquin interferes with the function of complex III (cytochrome bc₁ complex) of the mitochondrial electron transport chain. Complex III is responsible for transferring electrons between molecules in the chain.
- 3. Disruption of Electron Transport:** Fenazaquin binds to the Q_o site of complex III, which inhibits the transfer of electrons. As a result, the normal flow of electrons through the chain is disrupted.
- 4. Production of Reactive Oxygen Species (ROS):** The disruption of the electron transport chain by fenazaquin can lead to the accumulation of electrons and a buildup of certain molecules. This can increase the production of reactive oxygen species (ROS), which are harmful to cells.
- 5. Mitochondrial Dysfunction:** The disruption of the electron transport chain and the increase in ROS production ultimately lead to mitochondrial dysfunction. Mitochondria are less efficient at producing ATP, which is essential for various cellular

processes, including muscle function and energy generation.

Paralysis and Death: The impairment of cellular energy production and overall cellular dysfunction result in paralysis and death of the mites or ticks that have been exposed to fenazaquin. It's important to note that the mode of action of fenazaquin is specific to mites and ticks due to differences in their physiology compared to other organisms. While fenazaquin has low toxicity to mammals, including humans, it's still essential to use it properly and follow recommended guidelines to minimize potential risks and to avoid the development of resistance in target populations.

Sulfoximine Insecticides

Sulfoximine insecticides, like sulfoxaflor, have a mode of action that affects the nervous systems of insects by targeting a specific type of receptor called the nicotinic acetylcholine receptor (nAChR). This disruption in nerve signal transmission leads to paralysis and eventual death in the targeted insects. Here's how the mode of action of sulfoximine insecticides typically works, using sulfoxaflor as an example:

1. **Targeting nAChRs:** Nicotinic acetylcholine receptors (nAChRs) are ion channels found on nerve cells in insects (and other animals). These receptors are involved in transmitting nerve signals by allowing the flow of ions, such as sodium and calcium, across the cell membrane.
2. **Normal Nerve Signal Transmission:** In a normal nerve impulse transmission, the neurotransmitter acetylcholine binds to nAChRs on the postsynaptic membrane of a nerve cell. This binding triggers the opening of ion channels and the flow of ions, leading to the transmission of the nerve signal.
3. **Disruption of nAChRs:** Sulfoxaflor and other sulfoximine insecticides bind specifically to AChRs on nerve cells, causing these receptors to become overstimulated.
4. **Hyperexcitation of Nerve Cells:** The binding of sulfoxaflor leads to excessive and prolonged opening of ion channels in nAChRs. This results in hyperexcitation of nerve cells, causing continuous and uncontrolled nerve signaling.
5. **Nerve Signal Disruption:** The hyperexcitation of nerve cells disrupts the normal transmission of nerve signals. This can lead to a range of effects, including paralysis and eventual death of the insect.

Selective Impact: The mode of action of sulfoximine insecticides like sulfoxaflor is relatively specific to insects due to differences in the structure of nAChRs between insects and mammals. This selectivity helps minimize the impact on non-target organisms. As with any class of insecticides, there is a potential for the development of resistance in insect populations. To manage resistance and minimize potential environmental impacts, integrated pest management practices that include the rotation of different insecticide classes are recommended when using sulfoximine insecticides.

Oxime Carbamate Insecticide

Thiodicarb is an oxime carbamate insecticide with a mode of action that affects insects' nervous systems by inhibiting the activity of the enzyme acetylcholinesterase (AChE). This disruption in nerve signal transmission leads to paralysis and eventual death in the targeted insects. Here's how the mode of action of thiodicarb typically works:

1. Role of Acetylcholinesterase (AChE)

Acetylcholinesterase is an enzyme that plays a critical role in the transmission of nerve signals. After the neurotransmitter acetylcholine is released from a nerve cell, AChE breaks down acetylcholine in the synapses (junctions between nerve cells). This termination of the nerve signal allows the nerve cell to reset for the next transmission.

2. **Normal Nerve Signal Termination:** In a normal nerve impulse transmission, acetylcholine binds to receptors on the postsynaptic membrane of a nerve cell. This triggers the transmission of the nerve signal. After the signal is transmitted, AChE rapidly breaks down acetylcholine, terminating the signal.
3. **Inhibition of AChE:** Thiodicarb inhibits the activity of AChE. When AChE is inhibited, acetylcholine accumulates in the synapses, prolonging nerve signaling.
4. **Nerve Signal Disruption:** Due to the inhibition of AChE, nerve signals are not properly terminated. This results in continuous and uncontrolled nerve signaling, which can lead to paralysis and other effects in insects.
5. **Impact on Nervous System:** The prolonged nerve signaling and disrupted nerve transmission ultimately lead to paralysis and death of the insect. It's important to note that the mode of action of thiodicarb is specific to insects due to the differences in the structure of AChE between insects and mammals. However, similar to other classes of insecticides, there is a potential for the development of resistance in insect populations. Integrated pest management practices that include the rotation of different insecticide classes are recommended to manage resistance and minimize potential environmental impacts.

Thiourea Insecticides

Diafenthiuron is a thiourea insecticide that affects insects by disrupting their nervous systems and causing paralysis. The mode of action of diafenthiuron involves interfering with the proper functioning of mitochondria, which are the energy-producing organelles within cells. Here's how the mode of action of diafenthiuron typically works:

- **Mitochondrial Uncoupling:** Diafenthiuron interferes with the process of oxidative phosphorylation that occurs in the mitochondria. This process is responsible for generating adenosine triphosphate (ATP), which is the cell's primary energy source. Diafenthiuron acts as a mitochondrial uncoupler, which means it disrupts the normal coupling of electron transport and ATP synthesis.
- **Proton Leakage:** Diafenthiuron allows protons (positively charged hydrogen ions) to leak across the mitochondrial inner membrane. Normally, the flow of protons is tightly regulated and is coupled with ATP synthesis. However, the uncoupling action of diafenthiuron disrupts this regulation, leading to the loss of the proton gradient used to generate ATP.

- **Reduced ATP Production:** As protons leak back into the mitochondrial matrix without contributing to ATP synthesis, the production of ATP is significantly reduced. ATP is essential for many cellular processes, including muscle contraction and nerve signal transmission.
- **Energy Depletion:** The reduced production of ATP impairs the insect's ability to carry out essential physiological functions, such as muscle movement and normal nerve signaling.

Paralysis and Death: The disruption of ATP production and overall energy depletion ultimately leads to paralysis and death in insects. It's important to note that the mode of action of diafenthiuron is specific to insects due to differences in the structure and function of mitochondria between insects and other organisms. However, as with any class of insecticides, the potential for the development of resistance exists. Integrated pest management practices, including proper rotation and use of different insecticide classes, are essential for managing resistance and minimizing potential environmental impacts.

Butenolide Insecticide

Flupyradifurone is a butenolide insecticide that has a mode of action affecting the nervous systems of insects. It targets a specific type of receptor known as the nicotinic acetylcholine receptor (nAChR), disrupting nerve signal transmission and leading to paralysis and death in the targeted insects. Here's how the mode of action of flupyradifurone typically works:

1. **Targeting nAChRs:** Nicotinic acetylcholine receptors (nAChRs) are ion channels found on nerve cells in insects (and other animals). They play a crucial role in transmitting nerve signals by allowing the flow of ions, such as sodium and calcium, across the cell membrane.
2. **Normal Nerve Signal Transmission:** In a normal nerve impulse transmission, the neurotransmitter acetylcholine binds to nAChRs on the postsynaptic membrane of a nerve cell. This binding triggers the opening of ion channels and the flow of ions, leading to the transmission of the nerve signal.
3. **Disruption of nAChR:** Flupyradifurone and other butenolide insecticides bind specifically to nAChRs on nerve cells, causing these receptors to become overstimulated.
4. **Hyperexcitation of Nerve Cells:** The binding of flupyradifurone leads to excessive and prolonged opening of ion channels in nAChRs. This results in hyperexcitation of nerve cells, causing continuous and uncontrolled nerve signaling.
5. **Nerve Signal Disruption:** The hyperexcitation of nerve cells disrupts the normal transmission of nerve signals. This can lead to paralysis and eventual death of the insect.
6. **Selective Impact:** The mode of action of flupyradifurone is relatively specific to insects due to differences in the structure of nAChRs between insects and mammals. This selectivity helps minimize the impact on non-target organisms. As with any class of insecticides, there is a potential for the development of resistance in insect populations. Integrated pest management practices that include the rotation of different insecticide classes are recommended to

manage resistance and minimize potential environmental impacts when using flupyradifurone and other butenolide insecticides.

Mode of Action of *Bacillus thuringiensis*

Bacillus thuringiensis (Bt) is a type of bacteria commonly used as a biological pesticide. Its mode of action involves producing proteins called crystal toxins or Cry toxins. When insects consume plants treated with Bt or Bt-based insecticides, these toxins are ingested. Inside the insect's gut, the toxins bind to specific receptors on the gut lining. This binding disrupts the integrity of the gut membrane, causing it to break down. As a result, the insect's gut contents, including bacteria and toxins, leak into its body, leading to a condition called septicemia. This systemic infection overwhelms the insect's immune system and causes it to stop feeding. Additionally, the Cry toxins can disrupt ion balance within the gut cells, leading to paralysis and further gut disruption. Ultimately, the combination of gut breakdown, systemic infection, and paralysis leads to the death of the insect. Importantly, Bt toxins are highly specific to certain groups of insects and are generally considered safe for humans, animals, and most non-target organisms. This specificity is due to the specific receptors present on the gut lining of susceptible insects. Bt-based insecticides have been widely used in agriculture to control pest populations, offering an environmentally friendly alternative to chemical pesticides. However, the development of resistance in some insect populations over time has highlighted the importance of using Bt in integrated pest management strategies to maintain its effectiveness.

Mode of Action Nuclear Polyhedrosis Virus

The mode of action of NPV, or Nuclear Polyhedrosis Virus, involves several steps.

1. **Attachment and Entry:** NPV infects insects, particularly caterpillar larvae. The virus particles attach to the insect's cuticle (outer surface) and are ingested when the insect feeds on virus-contaminated plant material. The alkaline environment of the insect's midgut dissolves the protective protein coat of the virus, releasing the viral particles.
2. **Primary Infection and Replication:** Once inside the insect's midgut cells, the virus starts replicating. The viral genome takes control of the cellular machinery to produce more virus particles. This process eventually causes the infected cells to lyse (burst), releasing a large number of new virus particles into the insect's body.
3. **Secondary Infection and Spread:** The newly produced virus particles infect neighboring cells within the insect's body, leading to the further replication of the virus. The virus spreads throughout the insect's hemolymph (insect blood), affecting various tissues and organs.
4. **Polyhedra Formation:** As the infection progresses, some cells near the end of the infection cycle produce large crystalline structures called polyhedra. These polyhedra contain thousands of virus particles each and serve as a protective casing for the virus, helping to preserve it in the environment.
5. **Lethal Effects:** The rapid replication and spread of NPV within the insect's body eventually cause severe damage to the insect's tissues and disrupt its physiological processes. This leads to the characteristic

symptoms of NPV infection, such as lethargy, loss of appetite, and eventually death.

Transmission: After the insect dies, the polyhedra are released into the environment along with the insect's body fluids. These polyhedra can persist for some time, protecting the virus from environmental factors. When other insects come into contact with the infected remains, they can ingest the polyhedra and become infected themselves, continuing the cycle. Overall, NPV is an example of a naturally occurring biological control agent used in pest management. It's effective in reducing pest populations because it specifically targets certain insect species without harming non-target organisms.

Mode of action of Granulovirus

Granulovirus is a type of virus that infects insects, particularly various species of Lepidoptera (moths and butterflies). Its mode of action involves several steps:

- 1. Attachment and Entry:** Granulovirus particles attach to the insect's cuticle (outer protective layer) through specific interactions. Once attached, the virus enters the insect host through natural openings or minor abrasions on the cuticle.
- 2. Primary Replication:** After entering the host, the virus particles release their genetic material, which typically consists of double-stranded DNA. This genetic material starts replicating within the insect cells, utilizing the host's cellular machinery to create more virus particles.
- 3. Expression of Viral Genes:** Granuloviruses produce various proteins encoded by their genes. Some of these proteins manipulate the host's cellular processes, effectively shutting down its normal functions to favor virus replication. Others are involved in the formation of specialized structures called occlusion bodies.
- 4. Occlusion Body Formation:** One of the distinguishing features of granuloviruses is the production of occlusion bodies. These are dense protein structures that encapsulate numerous virus particles. They provide protection for the virus against environmental factors, ensuring its survival until it can infect another host.
- 5. Cell Lysis and Spreading:** As the virus replication progresses, infected cells start to lyse (burst open), releasing a large number of occlusion bodies into the insect's body cavity. These occlusion bodies contain viral particles and are the source of infection for other insects. When an uninfected insect comes into contact with these occlusion bodies, the cycle begins anew.
- 6. Secondary Replication and Spread:** Once ingested or otherwise entered into an uninfected insect, the occlusion bodies dissolve, releasing the virus particles. These particles infect the midgut cells of the new host insect. The replication process starts again, creating more virus particles and ultimately leading to the formation of new occlusion bodies.
- 7. Death and Transmission:** The host insect eventually succumbs to the infection, usually due to the disruption of vital functions caused by the virus. The dead insect's body contains a high concentration of occlusion bodies. This provides a potent source of infection for other insects that might consume the remains or come into contact with them. Overall, granuloviruses use a combination of strategies to infect, replicate within, and spread to new host insects. The production of occlusion

bodies is a key element in their lifecycle, ensuring their survival in the environment and aiding in the transmission of infection to other susceptible insects.

Juvenile Hormone Mimics

Juvenile hormone mimics are chemicals that imitate the effects of juvenile hormone in insects. Juvenile hormones are natural compounds that play a crucial role in regulating insect development and metamorphosis. Juvenile hormone mimics have been developed as insecticides to disrupt the normal growth and development of insects. Their mode of action involves several steps:

- 1. Binding to Juvenile Hormone Receptors:** Juvenile hormone mimics are designed to bind to the same receptors as the natural juvenile hormones. These receptors are found on the surface of insect cells, particularly those involved in regulating growth and metamorphosis.
- 2. Disruption of Hormonal Signaling:** When the juvenile hormone mimic binds to its receptor, it triggers a cascade of signaling events within the insect's cells. This disrupts the normal hormonal signaling pathways that control various aspects of development, including metamorphosis, reproduction, and behavior.
- 3. Prolonged Larval State:** Insects typically undergo metamorphosis from larva to pupa and then to the adult stage. Juvenile hormone mimics interfere with this process by preventing the transition from larva to pupa. This leads to the insect remaining in its larval state for an extended period, disrupting its development and preventing it from maturing into a reproductive adult.
- 4. Inhibition of Molting:** Juvenile hormone mimics can also interfere with the process of molting, which is the shedding of the old exoskeleton to allow for growth. By disrupting the proper timing and regulation of molting, these mimics cause abnormalities in the insect's growth and development.
- 5. Reduced Fertility and Reproduction:** The disruption of hormonal signaling caused by juvenile hormone mimics often leads to reduced fertility and reproductive capacity in insects. This is particularly important in agricultural and pest control contexts, as it can limit the insect population's ability to reproduce and grow.

Insecticidal Effects: Over time, the prolonged exposure to juvenile hormone mimics and the resulting disruption of growth and development ultimately lead to the death of the insect. Without the ability to complete metamorphosis, insects become more vulnerable to environmental stresses and predators, further contributing to their decline. In summary, the mode of action of juvenile hormone mimics involves interfering with the normal hormonal regulation of insect development. By disrupting growth, development, and reproduction, these mimics effectively act as insecticides, providing a targeted way to control insect populations in agriculture and other settings.

Chitin Biosynthesis Inhibitors

Chitin biosynthesis inhibitors (CBIs) are a class of insecticides that disrupt the formation of chitin, a key component of insects' exoskeletons. The mode of action of CBIs involves targeting the synthesis of chitin and subsequently interfering with insect growth and development. Here's how they work:

1. **Chitin Synthesis:** Chitin is a tough, structural polysaccharide that forms the major component of insects' exoskeletons, as well as other arthropods. Insects produce chitin through a process called chitin biosynthesis. This involves the enzymatic conversion of glucose, a simple sugar, into chitin chains that are then assembled into the exoskeleton.
2. **CBIs Binding:** Chitin biosynthesis inhibitors are designed to bind to specific enzymes involved in the chitin biosynthesis pathway. These enzymes include chitin synthase, which is responsible for polymerizing glucose into chitin chains.
3. **Enzyme Inhibition:** Once CBIs bind to chitin synthase or other related enzymes, they inhibit the enzymes' activity. This prevents the proper polymerization of glucose into chitin chains, disrupting the formation of a functional exoskeleton during molting and growth.
4. **Molting and Growth Disruption:** Insects undergo molting, a process where they shed their old exoskeleton to accommodate growth. Without a properly formed exoskeleton, insects cannot undergo successful molting. CBIs effectively block this process, leading to deformities, incomplete molting, and growth abnormalities.
5. **Exoskeleton Integrity:** The exoskeleton not only provides structural support but also helps insects retain body moisture and protects them from environmental factors. When the chitin biosynthesis is inhibited, the integrity of the exoskeleton is compromised, leaving insects vulnerable to desiccation, physical damage, and predation.
6. **Delayed Development and Death:** Insect larvae treated with CBIs typically experience prolonged larval stages due to the inability to molt and progress through their development. This delayed development can lead to reduced feeding, decreased mobility, and increased susceptibility to stressors. Eventually, the insects may die due to the disruption of their growth and their vulnerability to external threats.
7. **Selective Impact:** One advantage of CBIs is their relatively specific mode of action, targeting insects' chitin biosynthesis pathways without affecting vertebrates or other non-target organisms that lack chitin in their bodies. Overall, the mode of action of chitin biosynthesis inhibitors results in growth disruption, molting abnormalities, and eventually, the death of insects. CBIs are widely used in agricultural pest management to control various insect pests that can damage crops while posing a reduced risk to non-target species.

Ecdysone Agonists

Ecdysone agonists are a class of insecticides that mimic the action of ecdysone, a natural hormone responsible for regulating the molting and metamorphosis processes in insects. The mode of action of ecdysone agonists involves manipulating the insect's hormonal system to disrupt its development and ultimately lead to its death. Here's how they work:

- **Ecdysone Receptor Binding:** Ecdysone agonists are designed to bind to ecdysone receptors on the surface of insect cells. Ecdysone receptors are proteins that normally bind to the insect's natural ecdysone hormone, initiating a series of biological responses related to molting and metamorphosis.

- **Hormonal Signaling:** When an ecdysone agonist binds to the ecdysone receptor, it triggers a signaling cascade within the insect's cells. This cascade mimics the natural signaling pathway initiated by ecdysone, which is crucial for controlling the timing and regulation of molting and metamorphosis.
- **Molting Disruption:** Ecdysone agonists can disrupt the precise timing and coordination of molting in insects. Instead of progressing through the usual molting stages, insects treated with these agonists may experience abnormal and incomplete molting, leading to deformities and growth inhibition.
- **Metamorphosis Abnormalities:** Insects go through a complex process of metamorphosis, transitioning from larva to pupa and then to the adult stage. Ecdysone agonists can interfere with this process, causing insects to remain stuck in earlier developmental stages or preventing them from reaching the adult stage altogether.
- **Inhibition of Reproduction:** By disrupting the natural hormonal processes, ecdysone agonists often lead to reduced fertility and reproductive capacity in insects. This is a significant advantage in pest management, as it limits the ability of insects to reproduce and maintain their populations.
- **Impaired Adult Formation:** Even if insects treated with ecdysone agonists manage to reach the adult stage, they may exhibit abnormalities in adult morphology and behavior. This further reduces their ability to survive, reproduce, and contribute to the pest population.

Selective Impact: Ecdysone agonists can have a relatively selective impact on insects due to the specific targeting of ecdysone receptors. This reduces the risk of harming non-target organisms that do not have the same hormonal pathways. In summary, the mode of action of ecdysone agonists involves manipulating the hormonal regulation of insect development, leading to disrupted molting, abnormal metamorphosis, and reduced reproductive capacity. These effects collectively contribute to the control of insect populations and are utilized in integrated pest management strategies.

Conclusion

In the grand tapestry of pest management, the mode of action of novel insecticides emerges as a thread that weaves together innovation, precision, and ecological stewardship. As we cast our gaze back over this exploration, we glimpse a future where the battle against insect pests is no longer waged with broad strokes, but rather with the finesse of a surgeon's scalpel. The conventional era of indiscriminate chemical warfare is yielding to a new era of sophistication one where science listens to nature's whispers and adapts its solutions accordingly. The mode of action of novel Insecticides is not merely a scientific curiosity; it is a beacon guiding us towards a landscape where agricultural bounty thrives without leaving ecological wreckage in its wake. It is a testament to the relentless pursuit of knowledge, where molecular insights become the compass that directs us away from the pitfalls of resistance and towards the shores of sustainable pest control. The narrative of innovation does not end here; it is a living narrative, one that continues to unfold in laboratories, fields, and research institutions around the world. As we bid adieu to this exploration, let us

carry forward the lessons learned from the mode of action of novel insecticides an homage to the power of scientific inquiry and the promise of a future where human ingenuity, coupled with nature's wisdom, charts a course towards a more balanced and prosperous world.

References

1. Biondi A, Mommaerts V, Smagghe G, Viñuela E, Zappalà L, Desneux N. The nontarget impact of spinosyns on beneficial arthropods. *Pest Management Science*. 2012;68(12):1523-1536.
2. Bret BL, Larson LL, Schoonover JR, Sparks TC, Thompson GD. Biological properties of spinosad. *Down to Earth*. 1997;52(1):6-13.
3. Cloyd RA, Bethke JA. Impact of neonicotinoid insecticides on natural enemies in greenhouse and interiorscape environments. *Pest Management Science*. 2010;66(2):136-143.
4. Cordova D, Benner EA, Sacher MD, Rauh JJ, Sopa JS, Lahm GP, *et al*. Anthranilic diamides: A new class of insecticides with a novel mode of action, ryanodine receptor activation. *Pesticide Biochemistry and Physiology*. 2006;84(3):196-214.
5. Eger JE, Lindenburg LB. Utility of spinosad for insect control in vegetables. *Proceedings of the Florida State Horticultural Society*. 1998;111:55-57.
6. Ghanim M, Ishaaya I. Insecticides with novel modes of action: Mechanism and resistance management. In: *Tolerance to Environmental Contaminants*. K10919.indb. 2016. p. 385-407.
7. Hardy MC. Resistance is not futile: It shapes insecticide discovery. *Insects*. 2014;5(1):227-242.
8. Hemingway J, Hawkes NJ, McCarroll L, Ranson H. The molecular basis of insecticide resistance in mosquitoes. *Insect Biochemistry and Molecular Biology*. 2004;34(7):653-665.
9. Insecticide Resistance Action Committee (IRAC). Resistance management for sustainable agriculture and improved public health. 2007. Available from: <https://irac-online.org/>