

International Journal of Statistics and Applied Mathematics

ISSN: 2456-1452

NAAS Rating (2025): 4.49

Maths 2025; 10(12): 173-176

© 2025 Stats & Maths

<https://www.mathsjournal.com>

Received: 05-11-2025

Accepted: 09-12-2025

Mamadou Diallo

Department of Agricultural
Economics, University of Dakar,
Dakar, Senegal

Fatoumata Sarr

Department of Agricultural
Economics, University of Dakar,
Dakar, Senegal

Statistical analysis of papaya yield variability in the Niayes Area: A typological approach

Mamadou Diallo and Fatoumata Sarr

DOI: <https://www.doi.org/10.22271/math.2025.v10.i12b.2222>

Abstract

Papaya (*Carica papaya*) cultivation plays a significant role in the agricultural economy of the Niayes area in Senegal. Despite its economic importance, there remains considerable variability in papaya yield due to various factors including environmental conditions, farm management practices, and socio-economic influences. This research aims to analyze the yield variability of papaya farms in the region using a statistical typological approach. By applying cluster analysis and regression models, this research identifies distinct typologies of papaya farms based on factors such as farm size, irrigation techniques, and fertilizer usage. Furthermore, the research explores the relationship between farm typologies and yield variability, aiming to provide insights into the underlying factors contributing to production differences. The data used in this research was collected from a sample of papaya farms across the Niayes area. The results indicate that farm management practices, particularly irrigation and fertilizer usage, significantly influence yield outcomes, while environmental factors such as soil type and rainfall patterns also play a critical role. By developing typologies of farms, this research offers a more targeted approach to improving papaya production, with recommendations for optimizing farm practices based on the identified typologies. These findings have practical implications for agricultural policy-makers and farmers in the region, offering insights into how farm management improvements can be tailored to specific farm types to reduce yield variability and increase overall production efficiency.

Keywords: Papaya yield variability, Niayes area, typological approach, statistical analysis, farm management, irrigation, fertilizer, farm typologies

Introduction

Papaya cultivation is a vital agricultural activity in the Niayes area of Senegal, contributing significantly to the local economy. The region's farmers face considerable challenges in achieving consistent yields, which are influenced by a range of factors including environmental conditions, farming practices, and socio-economic factors. Despite the growing importance of papaya as a cash crop, yield variability continues to be a challenge, hindering farmers' ability to maximize production and income. Understanding the causes of this variability is crucial for developing strategies to improve productivity.

The purpose of this research is to statistically analyze papaya yield variability in the Niayes area using a typological approach. This approach allows for the identification of distinct farm types based on factors such as farm size, irrigation techniques, and fertilizer usage. Previous studies have shown that farm management practices are closely linked to agricultural output, with practices such as irrigation and fertilization having significant impacts on crop yields [3]. However, there is limited research specifically addressing papaya yield variability in the Niayes area, and even less that uses a typological framework to categorize farms based on production characteristics.

The research objectives of this research are threefold.

First, it aims to identify typologies of papaya farms in the Niayes area based on key agricultural factors.

Second, the research seeks to analyze the relationship between these typologies and yield variability.

Corresponding Author:

Mamadou Diallo

Department of Agricultural
Economics, University of Dakar,
Dakar, Senegal

Finally, the research aims to provide recommendations for improving papaya production efficiency by tailoring interventions to specific farm types. The hypothesis of this research is that farm management practices, particularly irrigation and fertilizer use, have a direct impact on papaya yield, and that these impacts vary across different farm typologies.

The typological analysis is expected to provide valuable insights for farmers, agricultural practitioners, and policy-makers, facilitating more targeted interventions to enhance papaya yield stability and reduce the impacts of yield variability in the region ^[4]. The findings will contribute to the body of knowledge on papaya cultivation and farm management in West Africa, specifically in areas where environmental and socio-economic factors drive production outcomes.

Materials and Methods

Materials

The research was conducted in the Niayes area of Senegal, an important region for papaya cultivation, characterized by a semi-arid climate with distinct rainy and dry seasons. Fifty papaya farms were selected for the study for this research, representing a range of farm sizes, management practices, and environmental conditions. The farms were stratified based on farm size, irrigation methods, and fertilizer usage. The selected farms varied in size, ranging from smallholder farms of less than 1 hectare to larger commercial operations exceeding 5 hectares. Data collection involved a comprehensive survey, which was conducted between January and April 2023, during the dry season. The survey focused on farm characteristics such as the use of irrigation systems (drip, sprinkler, or flood irrigation), the type and number of fertilizers used (organic vs. inorganic), and the farm management practices employed. Soil samples were also collected from each farm to assess soil fertility, pH, and other relevant soil parameters that may influence papaya yield. Weather data, including rainfall and temperature, were obtained from the local meteorological station for the duration of the research period. The farms were also categorized based on socio-economic factors, such as access to markets and extension services, to understand their influence on farm productivity ^{[1][2]}.

Methods

To assess the variability in papaya yield, the research applied

a typological approach using cluster analysis to group farms based on common characteristics such as size, irrigation method, fertilizer use, and socio-economic factors. Descriptive statistics were calculated for each typology, including mean yield per farm, farm size, and inputs used. The data were then subjected to regression analysis to explore the relationships between farm typologies and yield variability. A multiple regression model was employed to examine the influence of farm management practices on yield, controlling for environmental variables such as soil type and weather conditions. The yield data were collected over a full production cycle, from planting to harvesting, and measured in terms of both quantity and fruit quality. Statistical analysis was performed using R software (version 4.0.3) to conduct both cluster and regression analyses. To ensure the reliability and validity of the findings, data were cross-validated with secondary sources, including government agricultural reports and previous studies in the region ^{[3][4][5]}. The findings from this analysis were used to develop practical recommendations for improving papaya production, tailored to the specific needs of different farm typologies in the Niayes area ^[6].

Results

Farm Typologies and Yield Variability

The farm typologies were categorized based on key factors such as farm size, irrigation techniques, fertilizer use, and socio-economic status. Three distinct typologies were identified through cluster analysis: small-scale farms (Typology 1), medium-scale farms (Typology 2), and large-scale farms (Typology 3). Each typology exhibited significant differences in both papaya yield and management practices.

The average papaya yield for

Typology 1: Small-scale farms was 15 tons per hectare, while Typology 2 (medium-scale farms) showed a higher yield of 25 tons per hectare.

Typology 3 (large-scale farms) had the highest yield at 35 tons per hectare.

A one-way ANOVA was conducted to determine whether the differences in yield between farm typologies were statistically significant. The results revealed that the mean yield for large-scale farms was significantly higher than both small- and medium-scale farms ($p < 0.05$). This is consistent with the findings of Faye *et al.* ^[4], who observed similar trends in horticultural crop yields in the Niayes area.

Table 1: Papaya Yield by Farm Typology

Typology	Average Yield (tons/hectare)	Farm Size (hectares)	Irrigation Type	Fertilizer Use
Typology 1 (Small)	15	<1	Drip	Organic/Inorganic
Typology 2 (Medium)	25	1-3	Sprinkler	Inorganic
Typology 3 (Large)	35	>5	Flood	Inorganic




Fig 1: Yield distribution across farm typologies

Regression Analysis of Farm Management Factors on Yield

To further understand the relationship between farm management practices and yield, multiple regression analysis was conducted. The regression model examined the impact of irrigation type, fertilizer use, and farm size on papaya yield. The model's R-squared value was 0.75, indicating that 75% of

the variation in yield could be explained by these factors. The results showed that fertilizer use ($p = 0.02$) and irrigation type ($p = 0.01$) were statistically significant predictors of yield, while farm size had a smaller but still significant effect ($p = 0.05$). These findings align with the work of Gueye *et al.* [6], who found that improved irrigation and fertilizer application significantly increased crop yields in the Niayes region.




Fig 2: Regression of Papaya Yield on Farm Management Factors

Interpretation of Results

The findings from the cluster analysis and regression models indicate that papaya yield in the Niayes area is influenced by farm typology and farm management practices. Larger farms with more access to irrigation and fertilizers tend to have higher yields, consistent with previous studies in the region [5][6]. This suggests that improvements in irrigation infrastructure and fertilizer use could lead to significant yield improvements for smaller and medium-scale farms.

The regression analysis further emphasizes the importance of fertilizer use and irrigation in optimizing papaya production. These results are in line with previous research, which highlights the role of proper farm management in increasing crop yields, especially in semi-arid regions like the Niayes area [3][4]. However, the impact of farm size is less pronounced, suggesting that other factors, such as access to markets and technical knowledge, may also play a role in determining yield outcomes.

Overall, these results provide a comprehensive understanding of the factors driving papaya yield variability in the Niayes area. They highlight the need for targeted interventions to optimize farm practices, particularly for small-scale farmers who may not have the same resources as larger farms.

Discussion

The findings of this research highlight the significant role that farm management practices and farm typology play in influencing papaya yield variability in the Niayes area of Senegal. The typological classification of farms revealed three distinct categories, with large-scale farms consistently outperforming smaller ones in terms of yield. This trend aligns with previous research, where larger farms with better access to resources such as irrigation and fertilizers were found to have higher crop yields [1][4]. Our results suggest that farm size, while not the only determinant of yield, serves as a strong indicator of access to necessary agricultural inputs that can substantially increase production.

The regression analysis further supported the notion that farm management practices, particularly fertilizer use and irrigation, significantly affect papaya yield. This finding corroborates the work of Faye *et al.* [4] and Ndiaye *et al.* [5],

who demonstrated that appropriate nutrient management and irrigation methods lead to improved productivity. Specifically, the results showed that fertilizer use had a strong positive impact on papaya yield, with a statistically significant relationship ($p = 0.02$). The importance of fertilizers, especially in semi-arid areas like the Niayes region, is crucial, as these areas often experience soil nutrient deficiencies that limit crop growth. This highlights the need for targeted interventions that promote the use of optimized fertilization techniques, which could improve yields, especially for small-scale farmers who may lack access to high-quality fertilizers. Moreover, the regression model revealed that irrigation type (drip, sprinkler, or flood) also significantly influenced papaya yield ($p = 0.01$). This finding emphasizes the critical role of water management in agricultural production. The advantages of drip irrigation, in particular, have been well-documented, as it provides efficient water usage, reduces water waste, and can improve crop yield by ensuring plants receive consistent moisture [6]. Given the water scarcity in the Niayes area, enhancing irrigation infrastructure, especially for smaller farms, could greatly reduce yield variability and improve the sustainability of papaya farming in the region.

The research's findings also underscore the potential for farm typology-based interventions to improve agricultural outcomes. By categorizing farms based on their management practices and resources, policy-makers and agricultural support programs can tailor their strategies to the specific needs of each typology. For instance, smaller farms may benefit more from access to affordable irrigation systems and subsidized fertilizers, while larger farms may require advanced technologies for pest management and production optimization. This targeted approach could reduce yield disparities across farms and promote more equitable agricultural development in the region.

Conclusion

This research provides a comprehensive analysis of the factors influencing papaya yield variability in the Niayes area of Senegal. The results indicate that farm typology, characterized by factors such as farm size, irrigation methods, and fertilizer use, plays a crucial role in determining yield

outcomes. Large-scale farms were found to have significantly higher yields compared to smaller farms, primarily due to better access to resources such as irrigation and fertilizers. The regression analysis further confirms the significant influence of farm management practices, particularly fertilizer use and irrigation techniques, on yield variability. The findings underscore the critical role of water management and optimized fertilization in improving papaya production in the region.

The research's results suggest that improving farm practices, especially for small-scale farmers, could significantly reduce yield variability and enhance overall productivity. Practical recommendations based on the findings include increasing access to efficient irrigation systems such as drip irrigation, which can ensure a more consistent water supply and minimize water wastage. For small- and medium-scale farms, targeted subsidies or financial support for purchasing quality fertilizers and irrigation equipment could help improve productivity. Additionally, training programs should be implemented to educate farmers on the best practices for fertilizer application and irrigation management, focusing on cost-effective and environmentally sustainable methods. Further investment in agricultural infrastructure, including improved soil testing facilities and weather forecasting services, would enable farmers to make better-informed decisions regarding crop management. Extension services should be strengthened to provide tailored advice to farmers, particularly in areas with limited access to resources, ensuring that interventions are based on farm typology and specific needs. Finally, policymakers should prioritize the establishment of farmer cooperatives to improve market access, reduce input costs, and foster the exchange of knowledge and best practices. By implementing these recommendations, agricultural productivity in the Niayes region can be optimized, reducing the gap between farm typologies and increasing the overall sustainability of papaya production. These efforts could pave the way for a more resilient agricultural sector that is better equipped to withstand environmental challenges and ensure food security for the growing population in the region.

References

1. Gueye M, Thiam A, Diawara A. Farm typology and its relation to papaya yield variability in Senegal. *Int J Agric Econ*. 2023;47(3):105-115.
2. Sow B, Ndiaye MA, Diouf R. Impact of soil fertility management on crop yield in the Niayes area of Senegal. *J Agric Sci*. 2022;11(2):78-89.
3. Ndiaye A, Tine J, Fall D. Effects of climate change on papaya production in Senegal. *J Environ Sci*. 2020;8(1):33-47.
4. Faye M, Diagne A. Irrigation practices and their effect on the productivity of horticultural crops in the Niayes area of Senegal. *Agric Water Manag*. 2021;240:106245.
5. Ndiaye S, Laye H, Mbacké A. Fertilizer application and its impact on papaya yield in Senegal. *African J Agric Res*. 2020;16(5):159-168.
6. Cissé AM, Dieye C, Diedhiou PM, Mbacké B, Diedhiou I. Typology of papaya-producing farms in the Niayes area (Senegal). *Int J Hortic Food Sci*. 2024;6(2):45-52. DOI:10.33545/26631067.2024.v6.i2a.228.
7. Faye B, Lo B. The role of irrigation in improving fruit yields in semi-arid areas of Senegal. *J Agric Res*. 2021;10(1):15-23.
8. Diouf M, Ba FA, Diaw M. The socio-economic factors affecting papaya production in Senegal. *J Agric Econ Dev*. 2021;34(4):85-99.
9. Kane A, Diallo D, Toure M. Typological approaches to agricultural productivity in West Africa. *Agric Syst*. 2020;132:87-97.
10. Mbengue M, Khouma A, Ndione M. Modeling the impact of fertilizer use on papaya production in the Niayes area. *Agric Syst*. 2022;43(2):123-135.
11. Fall D, Keita K, Mbaye A. Papaya production and environmental factors in the Senegalese Niayes region. *J Clim Agric*. 2023;11(3):234-245.
12. Mendy S, Sidibe S, Niane K. Analysis of irrigation methods and their effect on fruit yield in Senegal's Niayes area. *Irrig Sci*. 2019;37(4):357-365.
13. Diatta T, Thiam L, Ndour M. Land use and farm practices influencing papaya productivity in the Niayes area of Senegal. *Land Use Policy*. 2022;99:105244.
14. Sarr H, Sow P, Cissé A. Sustainable agriculture in the Niayes area: Managing papaya yields under varying climatic conditions. *Sustainability*. 2021;13(1):119-128.