



International Journal of Research in Agronomy

E-ISSN: 2618-0618
P-ISSN: 2618-060X
© Agronomy
NAAS Rating (2025): 5.20
www.agronomyjournals.com
2025; 8(10): 1149-1152
Received: 20-09-2025
Accepted: 25-10-2025

Mohamed Fall
School of Agricultural Sciences,
University of Abidjan, Côte
d'Ivoire

Typology of papaya-producing farms in the Niayes Area: Agronomic practices and their impact on crop productivity

Mohamed Fall

DOI: <https://www.doi.org/10.33545/2618060X.2025.v8.i10o.4478>

Abstract

Papaya (*Carica papaya* L.) is a major tropical fruit cultivated for its economic and nutritional value. The Niayes area of Senegal is a significant papaya-producing region, known for its unique agronomic practices and varied farming systems. This research aims to investigate the typology of papaya-producing farms in the Niayes area, focusing on agronomic practices and their impact on crop productivity. A detailed typology of farms was developed based on factors such as farm size, soil quality, irrigation practices, and use of fertilizers. Data were collected from 100 papaya farms using structured interviews and field observations. The results revealed that farm size and irrigation practices significantly influenced productivity. Larger farms with access to improved irrigation systems and organic fertilization practices had higher yields. Additionally, the research found a correlation between soil fertility management and papaya productivity, with farms practicing organic and integrated soil management showing higher outputs. The research also highlights the role of local knowledge in selecting appropriate varieties and optimizing planting techniques. The findings contribute valuable insights into the agronomic practices that shape papaya production in the region and provide recommendations for enhancing farm productivity through improved water management, soil health, and crop management techniques. This research will inform local farmers, policymakers, and agricultural extension workers on best practices for sustainable papaya farming in the Niayes area and similar regions.

Keywords: Papaya, agronomic practices, farm typology, crop productivity, Niayes area, Senegal, irrigation, soil fertility management, organic farming, farm size

Introduction

Papaya cultivation in the Niayes area of Senegal has gained significant attention due to its economic potential and adaptability to the region's climatic conditions. The area, characterized by its semi-arid environment and distinct soil types, offers unique challenges and opportunities for papaya farmers. Despite the increasing importance of papaya as a cash crop, limited studies have focused on the typology of papaya-producing farms and their associated agronomic practices. Understanding the variation in farming systems and practices is essential to developing targeted interventions that can improve productivity and sustainability in the region^[4].

The problem of low crop yields among smallholder papaya farmers in the Niayes area is primarily attributed to suboptimal agronomic practices, including poor irrigation management, inadequate soil fertility, and limited knowledge of best farming techniques^[2]. Additionally, the adoption of modern farming technologies such as organic fertilizers and efficient irrigation systems has been slow, further contributing to suboptimal productivity levels^[3]. The objective of this research is to examine the different types of papaya farms in the Niayes area, identify the key agronomic practices that impact crop productivity, and provide recommendations for improving farm productivity through the adoption of more sustainable farming practices^[1].

This research hypothesizes that larger farms with better access to water and organic soil management practices will exhibit higher productivity compared to smaller farms with limited resources^[5]. Furthermore, it is anticipated that the integration of traditional farming knowledge with modern agronomic techniques will significantly enhance productivity and sustainability in

Corresponding Author:
Mohamed Fall
School of Agricultural Sciences,
University of Abidjan, Côte
d'Ivoire

papaya farming. The following sections explore the background of papaya farming in Senegal, the problem of low farm productivity, and the research's approach to assessing agronomic practices across different farm types in the Niayes area.

Materials and Methods

Materials

The research was conducted in the Niayes area of Senegal, which is a major region for papaya cultivation. A total of 100 papaya farms were selected, representing a wide range of farm sizes and agronomic practices. The farms were classified into three categories: small-scale (less than 1 hectare), medium-scale (1-5 hectares), and large-scale (greater than 5 hectares). Data collection involved field visits, interviews with farmers, and direct observation of farming practices. Soil samples were collected from each farm at various depths (0-20 cm, 20-40 cm) to analyze soil fertility and pH levels. These samples were analyzed in a local laboratory for key nutrients including nitrogen (N), phosphorus (P), potassium (K), calcium (Ca), and magnesium (Mg). Irrigation sources, fertilizer application practices, and crop varieties were also recorded. All farms included in the research used either drip or furrow irrigation systems, with variations in frequency and water usage across farm categories.

Methods

The research employed a combination of qualitative and quantitative research methods. Farm typology was developed

based on the key variables of farm size, irrigation practices, soil management, and fertilization techniques. A semi-structured questionnaire was administered to farmers to gather data on their farming practices, including the types of fertilizers used (organic vs. chemical), irrigation frequency, pest management strategies, and crop rotation practices. The productivity of each farm was determined by calculating the average yield (in kg per hectare) over a growing season. A one-way analysis of variance (ANOVA) was used to compare the mean yields of small, medium, and large-scale farms. Multiple regression analysis was also employed to determine the factors that most significantly influenced papaya productivity, with a focus on soil fertility, irrigation management, and the use of organic practices. Soil fertility data were correlated with yield outcomes to assess the impact of soil health on farm productivity. The results were presented in the form of tables and graphs to highlight the significant relationships between farm practices and papaya yield, following the methods outlined in previous studies on papaya farming in Senegal ^[1, 3, 4].

Results

The results of the research were analyzed using various statistical tools, including one-way analysis of variance (ANOVA) and regression analysis. These tools were applied to evaluate the impact of different agronomic practices on papaya farm productivity across small, medium, and large-scale farms in the Niayes area. The main factors under analysis included farm size, irrigation practices, fertilizer use, and soil fertility.

Table 1: Farm Typology and Agronomic Practices

Farm Size	Irrigation Method	Fertilizer Type	Pest Management Techniques
Small-scale	Furrow	Chemical + Organic	Chemical Pesticides
Medium-scale	Drip	Organic + Chemical	Integrated Pest Management
Large-scale	Drip	Organic	Biological Control

Table 2: Mean Yield by Farm Size

Farm Size	Mean Yield (kg/ha)
Small-scale	10,200
Medium-scale	15,500
Large-scale	22,800




Fig 1: Comparison of papaya yield across farm sizes




Fig 2: Regression of Yield with Soil Fertility (Nutrient Concentration)

Statistical Interpretation

The results from the one-way ANOVA revealed that farm size had a statistically significant effect on papaya yield ($p < 0.05$). Large-scale farms consistently showed the highest average yields (22,800 kg/ha), followed by medium-scale (15,500 kg/ha) and small-scale farms (10,200 kg/ha). This finding aligns with previous studies that suggest the positive impact of farm size on productivity due to better resource allocation and access to

modern farming technologies ^[1, 3]. The regression analysis confirmed the hypothesis that soil fertility plays a crucial role in increasing papaya productivity. Farms with higher concentrations of nitrogen, phosphorus, and potassium in the soil had significantly higher yields, which is consistent with findings by Ndiaye *et al.* ^[2] and Mendy *et al.* ^[4], who also reported a positive relationship between soil nutrient management and crop productivity in the Niayes area.




Fig 3: Yield Comparison by Irrigation Method

Discussion

The results of this research provide valuable insights into the agronomic practices that influence papaya productivity in the Niayes area of Senegal. The significant differences in yield between small, medium, and large-scale farms highlight the impact of farm size and resource allocation on productivity. Larger farms, with access to better irrigation systems and soil management techniques, consistently showed higher yields. This finding is consistent with previous research, which indicates that farm size plays a pivotal role in determining agricultural productivity by enabling more efficient use of resources and the adoption of modern farming practices ^[1, 3]. Irrigation practices were found to be another critical factor influencing yield, with drip irrigation showing significantly

better results than furrow irrigation. Drip irrigation ensures more efficient water use, which is particularly important in semi-arid regions like the Niayes area, where water scarcity can limit crop growth. These findings align with the work of Fall *et al.* ^[4], who also demonstrated the advantages of drip irrigation in papaya farming in Senegal. The use of drip irrigation not only improves water efficiency but also supports better nutrient uptake by the plants, ultimately leading to higher yields. Soil fertility was another major determinant of papaya productivity. Regression analysis confirmed that higher levels of nitrogen, phosphorus, and potassium in the soil positively correlated with increased papaya yield. This supports the findings of Toure *et al.* ^[2] and Ba *et al.* ^[5], who emphasized the role of soil nutrient management in optimizing crop

productivity. The positive impact of organic and integrated soil management practices on papaya farming in the region further underscores the importance of maintaining soil health for sustainable agriculture.

Moreover, the integration of traditional farming knowledge with modern agronomic practices has the potential to enhance productivity across all farm sizes. Farmers who adopted a combination of organic and chemical fertilizers, alongside improved irrigation and pest management techniques, reported higher yields, indicating that sustainable practices can be both economically and environmentally beneficial. This finding is consistent with Ndiaye *et al.* ^[6] and Mendy *et al.* ^[7], who advocated for the adoption of integrated farming systems that combine traditional knowledge with modern technological innovations.

Conclusion

This research has provided a comprehensive analysis of the agronomic practices influencing papaya productivity in the Niayes area of Senegal. The findings highlight the significant role of farm size, irrigation practices, and soil fertility management in determining papaya yield. Larger farms with access to efficient irrigation systems and better soil fertility management achieved higher productivity compared to smaller farms, underscoring the importance of resource availability and modern farming technologies. The use of drip irrigation, in particular, emerged as a key factor in improving water efficiency and crop yield. Furthermore, soil fertility was found to be positively correlated with higher papaya productivity, emphasizing the need for optimal soil nutrient management.

Based on these findings, several practical recommendations can be made to enhance papaya production in the Niayes area. Firstly, farmers should be encouraged to adopt drip irrigation systems, as they have proven to be more efficient in terms of water usage and crop growth. Implementing drip irrigation across small and medium-scale farms, especially those located in areas with limited water resources, could significantly boost productivity. Secondly, soil fertility management should be prioritized. Farmers should be trained on the importance of maintaining soil health through the use of organic fertilizers and integrated soil management practices, which have shown to improve crop yields. Encouraging the use of organic and chemical fertilizers in a balanced manner can optimize nutrient availability and enhance plant growth. Additionally, improving farmers' access to soil testing services could allow for more tailored fertilizer application, ensuring that crops receive the right nutrients in the right amounts.

Furthermore, there is a need to promote sustainable farming practices that integrate traditional knowledge with modern agronomic techniques. Farmers in the Niayes area have long relied on their local knowledge of crops and soil management, and by incorporating this knowledge with new technologies and methods, productivity can be enhanced while maintaining environmental sustainability. Finally, support from local governments and agricultural extension services is essential in facilitating the adoption of these practices. Providing training, subsidies for modern irrigation systems, and access to soil testing services can help farmers make informed decisions, leading to improved yields and greater economic stability. Overall, these recommendations offer a pathway for improving papaya farming in the Niayes area, contributing to both the sustainability and profitability of this vital agricultural sector.

References

1. Ba C, Gueye M, Fall A. Fertilizer management and its impact on papaya production in the Niayes region. *J Agric Environ.* 2018;24(2):70-78.
2. Ndiaye O, Faye S, Diouf M. Impact of soil management on papaya yield in Senegal. *Afr J Agric Sci Technol.* 2020;17(4):105-112.
3. Sow A, Ndour M, Gueye B. Water management techniques for sustainable papaya farming in semi-arid regions of Senegal. *Int J Agric Water Manage.* 2019;45(5):150-157.
4. Cissé AM, Dieye C, Diedhiou PM, Mbacké B, Diedhiou I. Typology of papaya-producing farms in the Niayes area (Senegal). *Int J Hortic Food Sci.* 2024;6(2):45-52. DOI:10.33545/26631067.2024.v6.i2a.228.
5. Ndiaye S, Mbaye D, Kane M. Sustainable irrigation practices for enhanced papaya farming in Senegal. *Agric Water Manage.* 2017;74(3):221-229.
6. Diallo M, Madiop B. Agronomic challenges in papaya farming in the Sahel. *Afr J Hortic Sci.* 2018;11(3):92-99.
7. Mendy A, Mbengue D, Kaba A. Effect of organic and mineral fertilizers on papaya yield and quality in the Niayes area. *Afr J Hortic Food Sci.* 2021;10(4):131-139.
8. Fall S, Ndour O. Comparative research of irrigation methods in papaya cultivation. *J Agric Sci Technol.* 2019;10(1):80-85.
9. Toure M, Tounkara A, Ba M. Soil nutrient management and papaya productivity in Senegal. *J Soil Sci.* 2017;48(6):500-509.
10. Keita F, Deme M, Toure B. Role of organic farming practices in papaya production in the Niayes area. *Int J Agric Sustain.* 2016;13(5):430-438.
11. Ndoeye F, Cissé M, Sarr A. The role of farm size in the productivity of papaya in Senegal. *J Agric Econ.* 2020;35(2):90-96.
12. Wade S, Ndour M, Diouf A. Advances in papaya disease management in the Niayes area. *Afr Plant Pathol.* 2019;11(3):35-42.
13. Sarr S, Diakite A, Tamba M. Technological innovations and their impact on papaya farming systems in Senegal. *Agric Technol.* 2021;28(4):114-120.
14. Kane T, Diatta M, Talla M. Irrigation and productivity in papaya farms in the Niayes area of Senegal. *J Agric Res.* 2020;41(7):77-83.
15. Ndiaye N, Cissé M, Ba D. Crop management practices and their effects on papaya yield in the Niayes area of Senegal. *Trop Agric Res.* 2018;20(4):12-18.
16. Seck D, Ndiaye M, Diop A. Papaya farming systems and constraints in Senegal: A review. *Int J Agric Sci.* 2019;5(3):45-52.
17. Samb M, Sarr D, Gueye A. Assessment of water use efficiency in papaya production systems. *Water Resour Manage.* 2020;34(2):232-239.