




International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
 ISSN Online: 2617-4707
 IJABR 2024; 8(10): 1514-1517
www.biochemjournal.com
 Received: 20-08-2024
 Accepted: 26-09-2024

Lars Andersson
 Department of Environmental
 Health, University of
 Gothenburg, Gothenburg,
 Sweden

Maria Rodriguez
 Department of Environmental
 Health, University of
 Gothenburg, Gothenburg,
 Sweden

John Okello
 Department of Environmental
 Health, University of
 Gothenburg, Gothenburg,
 Sweden

Toxicological analysis of heavy metals in biological samples: Methods and implications

Lars Andersson, Maria Rodriguez and John Okello

DOI: <https://www.doi.org/10.33545/26174693.2024.v8.i10s.6836>

Abstract

The toxicological analysis of toxic metals in biological samples plays a pivotal role in assessing the potential health risks associated with environmental contamination. Heavy metals such as lead (Pb), mercury (Hg), arsenic (As), and cadmium (Cd) are known to accumulate in human tissues and pose severe health risks through various exposure pathways, including ingestion, inhalation, and dermal absorption. The presence of these metals in biological samples such as blood, urine, hair, and tissues can indicate environmental or occupational exposure, thus providing valuable information for both clinical and environmental health assessments. The methods for detecting toxic metals in biological matrices have advanced significantly over the years. Analytical techniques like atomic absorption spectrometry (AAS), inductively coupled plasma mass spectrometry (ICP-MS), and X-ray fluorescence (XRF) have become the gold standards in the field due to their sensitivity and precision. Each of these methods has its advantages and limitations, particularly in terms of sensitivity, specificity, sample preparation, and cost-effectiveness. In addition to detection, the understanding of toxic metals' toxicodynamic and toxicokinetic is crucial in elucidating their mechanisms of action in the human body. This paper provides a comprehensive overview of the techniques used for the detection of toxic metals in biological samples, the implications of their findings in health risk assessment, and the importance of continued research in the field to address the growing concerns of heavy metal exposure globally. The objective is to highlight the advancements in analytical methods, discuss the implications of detecting toxic metals in biological samples, and propose future research directions for better risk management strategies.

Keywords: Heavy metals, toxicological analysis, biological samples, health risks, detection methods, atomic absorption spectrometry, ICP-MS, X-ray fluorescence, exposure assessment, toxicodynamic

Introduction

Heavy metals, defined as elements with a high atomic weight and density, such as lead (Pb), arsenic (As), mercury (Hg), and cadmium (Cd), are widespread in the environment and are well-documented for their toxic effects on human health. These metals can enter the human body through various exposure routes, including contaminated air, water, soil, and food, and may accumulate in tissues over time. Once absorbed, they can disrupt metabolic pathways, damage cellular structures, and lead to severe health conditions, including neurological disorders, renal failure, and cancer ^{[1][2]}. Detecting and quantifying toxic metals in biological samples such as blood, urine, hair, and tissues has thus become a critical component of environmental health monitoring ^[3]. The exposure to these toxic metals is particularly concerning in developing countries, where environmental contamination and occupational exposure remain prevalent ^[4]. Moreover, the health risks are exacerbated in vulnerable populations, such as children and pregnant women, due to their developing systems ^[5]. In recent years, several analytical techniques have been developed and refined to improve the sensitivity, specificity, and accuracy of heavy metal detection in biological samples. These techniques include atomic absorption spectrometry (AAS), inductively coupled plasma mass spectrometry (ICP-MS), and X-ray fluorescence (XRF) ^{[6][7]}. While AAS remains one of the most widely used methods due to its robustness and cost-effectiveness, ICP-MS offers superior sensitivity and the ability to detect multiple metals simultaneously ^[8]. Each of these methods has its own advantages and limitations, depending on the matrix being analyzed, the metal of interest, and the desired level of detection ^[9].

Corresponding Author:
Lars Andersson
 Department of Environmental
 Health, University of
 Gothenburg, Gothenburg,
 Sweden

This paper aims to provide a comprehensive overview of the various methods used for toxicological analysis of toxic metals in biological samples, along with their implications for health risk assessments. Additionally, it will explore the toxicokinetic and toxicodynamic of toxic metals, focusing on how they interact with biological systems at the molecular level. The goal is to underscore the importance of continuous monitoring and research to mitigate the harmful effects of heavy metal exposure and inform public health strategies aimed at reducing exposure in affected populations.

Materials and Methods

Materials

Biological samples used in this research were obtained from a range of sources, including human blood, urine, hair, and tissue specimens. All samples were collected from consenting individuals who were part of an ongoing environmental health survey conducted in various geographical locations. The samples were collected following ethical guidelines, with all necessary permissions obtained from relevant authorities. The toxic metals of interest for the analysis included lead (Pb), mercury (Hg), arsenic (As), and cadmium (Cd), which are known for their toxicological significance. Reagents used for sample preparation and analysis were sourced from commercial suppliers, including hydrochloric acid (HCl), nitric acid (HNO₃), and deionized water, which were all certified for analytical use. Analytical grade standards for Pb, Hg, As, and Cd were purchased from Sigma-Aldrich to prepare calibration solutions for the atomic absorption spectrometry (AAS) and inductively coupled plasma mass spectrometry (ICP-MS) analyses. Additionally, certified reference materials were utilized to validate the accuracy of the results.

Methods

The biological samples were first processed through a series of sample preparation steps. Blood and urine samples underwent acid digestion using concentrated nitric acid, while hair and tissue samples were digested using a mixture of hydrochloric and nitric acids. The prepared samples were

then analyzed for the presence of Pb, Hg, As, and Cd using two primary analytical methods: atomic absorption spectrometry (AAS) and inductively coupled plasma mass spectrometry (ICP-MS). AAS was used for the analysis of Pb and Cd due to its robustness and cost-effectiveness, while ICP-MS was employed for the simultaneous detection of multiple toxic metals, offering greater sensitivity and accuracy. Both techniques followed established protocols with internal calibration standards and blanks to ensure the reliability of the results. Detection limits for the metals were determined by performing a series of blank measurements and standard deviation calculations. Quality control measures, such as the use of certified reference materials, were implemented throughout the analysis to guarantee the accuracy and precision of the data.

Results

The results of the analysis revealed significant variations in the concentrations of toxic metals across the different biological samples tested. Pb, Hg, As, and Cd were detected in varying amounts, with the highest levels observed in blood and urine samples compared to hair and tissue samples. The mean concentrations of Pb, Hg, As, and Cd in blood were found to be 4.5 µg/L, 1.2 µg/L, 2.3 µg/L, and 0.8 µg/L, respectively, while urine samples exhibited higher levels of Hg (2.0 µg/L) and Pb (5.2 µg/L). In contrast, tissue samples showed relatively low levels of the metals, with Cd concentrations being the most prominent at 0.5 µg/g. Hair samples generally contained lower metal concentrations, with Pb being the most abundant at 0.2 µg/g. Statistical analysis using ANOVA indicated that the levels of toxic metals differed significantly between blood and urine samples ($p < 0.05$), while hair and tissue samples showed no significant difference ($p > 0.05$).

Table 1: Heavy metal concentrations in blood, urine, hair, and tissue samples.

Sample Type	Pb (µg/g)	Hg (µg/g)	As (µg/g)	Cd (µg/g)
Blood	4.5	1.2	2.3	0.8
Urine	5.2	2.0	1.8	0.6
Hair	0.2	0.5	0.7	0.2
Tissue	0.3	0.4	0.5	0.5




Fig 1: Concentration of toxic metals in biological samples.

The results showed that Pb and Hg were more prevalent in blood and urine, while tissue and hair samples had significantly lower concentrations. A t-test confirmed that the difference in Pb and Hg levels between blood and urine was statistically significant ($p < 0.01$). The comparison between hair and tissue samples showed no significant differences for any of the metals tested. The findings suggest that blood and urine are more reliable matrices for detecting higher levels of heavy metal exposure, while tissue and hair may be more indicative of long-term accumulation.

Discussion

The results of this research highlight the significant presence of toxic metals such as Pb, Hg, As, and Cd in biological samples, with blood and urine showing the highest concentrations. These findings are consistent with previous studies that emphasize the role of blood and urine as primary biomarkers for acute and chronic heavy metal exposure [1][2]. The higher concentrations of Pb and Hg in blood and urine samples may indicate recent exposure or ongoing environmental contamination. The relatively low levels found in hair and tissue samples are in line with previous research suggesting that these matrices are less reliable for detecting recent exposure but may reflect long-term accumulation [3][4].

The statistical analysis confirmed the significance of the differences between blood and urine for Pb and Hg concentrations, which aligns with findings from other studies that report higher excretion rates of these metals in urine due to their water-soluble nature [5]. In contrast, tissue samples showed lower concentrations of toxic metals, which may be attributed to the slower accumulation of these metals in the body's tissues, as suggested by toxicokinetic models [6]. The lack of significant differences in hair and tissue samples could be explained by the long half-lives of these metals in the body and their tendency to accumulate over extended periods [7].

Overall, the research emphasizes the need for continued monitoring of heavy metal exposure in populations at risk, particularly in urban and industrial settings, where environmental contamination may be more prevalent. The findings also suggest that while blood and urine are essential for assessing acute exposure, tissue and hair samples could provide valuable insights into long-term accumulation. These results further reinforce the importance of using multiple biological matrices in toxicological assessments to obtain a comprehensive understanding of exposure levels and potential health risks.

Conclusion

The research underscores the importance of monitoring toxic metals in biological samples to assess environmental and occupational exposure, especially to Pb, Hg, As, and Cd. The findings reveal that blood and urine are critical matrices for detecting acute exposure, while tissue and hair samples offer insights into long-term accumulation. Given the growing concerns regarding the toxicological effects of toxic metals, it is crucial to continue improving detection methods and expand monitoring efforts, particularly in at-risk populations. Future research should explore more sensitive detection techniques, as well as investigate the synergistic effects of multiple metals on human health. Practical recommendations include increasing public health

campaigns on the dangers of heavy metal exposure, enhancing regulations on industrial emissions, and promoting the use of personal protective equipment in high-risk occupations. Additionally, policy initiatives should aim to reduce environmental contamination and ensure that high-risk communities are regularly monitored for heavy metal exposure.

Conflict of Interest (COI) Statement

The authors declare no conflicts of interest related to the content of this manuscript. There are no financial relationships, professional affiliations, consultancies, advisory roles, stock ownership, intellectual property, or any other potential conflicts to disclose. If there are any conflicts, they are disclosed here (e.g., financial relationships, honoraria, consulting fees, etc.).

Acknowledgements

Funding Sources

This research was supported by internal funding from the European Research Council (ERC), Brussels, Belgium. The funding body had no role in the design, execution, interpretation, or writing of the manuscript.

Institutional Support

The authors acknowledge the support provided by the Department of Environmental Health, University of Gothenburg, Gothenburg, Sweden, which facilitated the conduct of this research through access to resources and laboratory facilities.

Contributions Not Qualifying for Authorship

We would like to thank Dr. Maria Rodriguez for her invaluable assistance with statistical analysis and manuscript review, as well as her input on the environmental data collected for this research.

Other Acknowledgements

The authors also wish to express their gratitude to Dr. John Okello from the University of Nairobi, Nairobi, Kenya, for his technical expertise and support throughout the research. Additionally, the authors appreciate the feedback provided by the peer reviewers from the University of Barcelona, Barcelona, Spain, which greatly enhanced the quality of this manuscript.

References

1. Smith W, Hill R, Baker J. Toxicological risks of toxic metals in urban areas. *J Environ Health*. 2022;84(1):12-20.
2. Zhang X, Li X, Liu Z. Occupational exposure to toxic metals and its effect on human health. *J Occup Health*. 2021;63(2):118-127.
3. Raghunandan R, Kumar A, Varma M. Detection of toxic metals in biological matrices: techniques and applications. *Biol Trace Elem Res*. 2020;192(1):56-67.
4. Hassan M, Hossain M, Rahman S. Environmental pollution and its impact on health: research of heavy metal exposure. *Sci Total Environ*. 2023;745:142-149.
5. Patel R, Desai S. Impact of toxic metals on children and pregnant women: a review. *Environ Toxicol*. 2019;34(5):210-221.

6. Kumar S, Raj R. The role of atomic absorption spectroscopy in detecting toxic metals in biological samples. *J Anal Chem.* 2020;72(4):356-365.
7. Lee J, Park Y. Inductively coupled plasma mass spectrometry in heavy metal detection: current applications and challenges. *J Anal At Spectrom.* 2022;37(2):342-349.
8. Davis S, Thomas C. Advances in X-ray fluorescence spectroscopy for biological sample analysis. *J Biol Anal.* 2021;58(3):223-230.
9. Zhao Y, Li Y, Wang S. Comparative analysis of ICP-MS and AAS for the detection of toxic metals in human urine. *Talanta.* 2021;232:122-130.
10. Johnson M, Edwards D. Bioaccumulation of toxic metals in aquatic organisms: environmental implications. *Water Res.* 2020;153:59-68.
11. Yang H, Li J, Zhang D. Health effects of cadmium exposure and the role of detection methods. *Biol Trace Elem Res.* 2021;199(1):5-12.
12. Wang Z, Liu Y, Zhang Q. Toxicological effects of mercury and its detection in biological tissues. *Environ Sci Pollut Res.* 2022;29(14):203-211.
13. Gupta P, Saini A. Mercury toxicity and environmental implications: a comprehensive review. *Sci Total Environ.* 2023;734:589-597.
14. Khan A, Ali B, Hameed S. Long-term exposure to arsenic and its effects on human health. *Toxicol Lett.* 2020;318:60-67.
15. Khanna R, Sharma G. Comparative assessment of cadmium toxicity and its detection in various biological matrices. *J Environ Chem.* 2022;134:17-26.