



International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
 ISSN Online: 2617-4707
 NAAS Rating (2025): 5.29
 IJABR 2025; 9(7): 1852-1855
www.biochemjournal.com
 Received: 04-06-2025
 Accepted: 07-07-2025

Ana María Rodríguez
 Institute of Agricultural
 Biotechnology, University of
 Córdoba, Córdoba, Argentina

José Luís Gómez
 Institute of Agricultural
 Biotechnology, University of
 Córdoba, Córdoba, Argentina

Sofia Petrova
 Institute of Agricultural
 Biotechnology, University of
 Córdoba, Córdoba, Argentina

Plant biochemistry: Exploring the mechanisms of plant defence systems against pathogens

Ana María Rodríguez, José Luís Gómez and Sofia Petrova

DOI: <https://www.doi.org/10.33545/26174693.2025.v9.i7w.6823>

Abstract

Plants have evolved a complex array of defence mechanisms to protect themselves from various biotic stresses, particularly pathogens. The biochemical pathways that underpin these defence systems play crucial roles in maintaining plant health and productivity. Plant defence mechanisms are categorized into pre-existing (constitutive) and inducible responses, with inducible defences often activated upon pathogen detection. These mechanisms include the production of secondary metabolites, signaling molecules such as jasmonic acid and salicylic acid, and the activation of specific plant immune receptors. One of the key elements in plant defence is the role of pattern recognition receptors (PRRs) that detect pathogen-associated molecular patterns (PAMPs), triggering a cascade of immune responses. The biosynthesis of defence-related compounds such as phytoalexins, pathogenesis-related proteins, and reactive oxygen species (ROS) is a central part of the plant's immune strategy. Additionally, recent advancements in molecular techniques have provided deeper insights into the genetic regulation of plant immune responses. This paper explores the biochemical mechanisms of plant defence systems, emphasizing the role of secondary metabolites, signaling molecules, and molecular patterns in pathogen recognition and response. We aim to highlight the interaction between plant metabolic pathways and the immune response, offering potential insights for improving plant resistance through biotechnological approaches. Understanding these biochemical processes is critical for developing crops with enhanced resistance to pathogens, thereby ensuring food security and agricultural sustainability in the face of growing environmental challenges.

Keywords: Plant defence, pathogens, biochemical mechanisms, secondary metabolites, pattern recognition receptors, immune response, phytoalexins, jasmonic acid, salicylic acid, reactive oxygen species

Introduction

Plants are continuously exposed to various pathogens such as bacteria, fungi, viruses, and nematodes, which pose significant threats to agricultural production worldwide. To survive these challenges, plants have developed intricate biochemical defence mechanisms. These defence systems can be broadly categorized into constitutive and inducible responses. Constitutive defences are always present in the plant, while inducible defences are activated in response to pathogen attack ^[1]. The biochemical pathways involved in plant defence are highly complex and involve the synthesis of secondary metabolites, the production of defence proteins, and the activation of immune signaling networks ^[2].

A central component of plant immune systems is the recognition of pathogen-associated molecular patterns (PAMPs) by pattern recognition receptors (PRRs) on the plant's surface. This recognition triggers a series of signaling cascades, including the activation of mitogen-activated protein kinase (MAPK) pathways and the production of reactive oxygen species (ROS) ^[3]. These responses serve to limit pathogen growth and initiate the expression of defence-related genes. Another important aspect of plant defence is the biosynthesis of secondary metabolites, such as phytoalexins, which are produced in response to pathogen invasion. These metabolites not only directly inhibit pathogen growth but also play a role in reinforcing the plant's immune barriers ^[4].

The objective of this paper is to review the biochemical mechanisms underlying plant defence systems, focusing on the role of secondary metabolites, signaling molecules, and molecular patterns in the recognition of pathogens. We hypothesize that understanding these

Corresponding Author:
Ana María Rodríguez
 Institute of Agricultural
 Biotechnology, University of
 Córdoba, Córdoba, Argentina

biochemical pathways can lead to the development of genetically engineered plants with improved resistance to pathogens, offering a sustainable solution for crop protection [5].

By exploring the molecular and biochemical facets of plant defence, this paper aims to contribute to the growing body of knowledge necessary for enhancing agricultural productivity and resilience against biotic stressors.

Materials and Methods

Materials

The research examined the biochemical response of *Arabidopsis thaliana* and *Solanum lycopersicum* to pathogen attack. Plants were grown in controlled greenhouse conditions and exposed to *Pseudomonas syringae* (bacterial) and *Botrytis cinerea* (fungal). Chemical elicitors such as methyl jasmonate (MeJA) and salicylic acid were used to trigger systemic acquired resistance. High-performance liquid chromatography (HPLC) and gas chromatography-mass spectrometry (GC-MS) were used to quantify secondary metabolites. Gene expression was analyzed via quantitative PCR (qPCR), and ROS production was measured using fluorometric assays with DCFDA.

Methods

To examine the plant's biochemical responses to pathogen attack, various defence-related metabolites were quantified using high-performance liquid chromatography (HPLC) and gas chromatography-mass spectrometry (GC-MS). Gene expression analysis of defence-related genes was conducted using quantitative PCR (qPCR) with primers specific to PR1, PR2, and LOX genes [4]. The levels of reactive oxygen species (ROS) were measured by fluorometric assays using DCFDA (2',7'-dichlorodihydrofluorescein diacetate) [5]. Enzyme activity assays were performed to assess the activity of key enzymes involved in plant defence, including phenylalanine ammonia-lyase (PAL) and peroxidases [6]. Statistical analysis was performed using one-way ANOVA, followed by Tukey's post-hoc test to compare treatment means for significant differences [7]. Data from three

biological replicates per treatment group were analyzed, and the results were expressed as means $\pm$ standard deviation.

Results

The analysis of secondary metabolite levels revealed a significant increase in the concentration of phytoalexins in *S. lycopersicum* plants after pathogen exposure. A 2-fold increase in phytoalexin production was observed in plants treated with methyl jasmonate compared to untreated controls ($p < 0.05$). Similarly, salicylic acid treatment resulted in a significant increase in the expression of pathogenesis-related (PR) genes, with a 1.5-fold increase in PR1 expression, indicating the activation of systemic acquired resistance pathways [2]. Enzyme activity assays also revealed that pathogen infection led to a significant increase in PAL and peroxidase activities, with higher levels observed in methyl jasmonate-treated plants. The ROS assay demonstrated that both *P. syringae* and *B. cinerea* infection induced ROS production, but higher levels were observed in plants treated with salicylic acid, indicating enhanced oxidative stress tolerance [5].

Statistical analysis of gene expression data indicated that salicylic acid treatment significantly upregulated PR2 and LOX genes compared to the control group ($p < 0.05$). Furthermore, ANOVA results showed that treatment with chemical elicitors, especially salicylic acid, significantly enhanced the overall plant defence response, reflected by increased levels of secondary metabolites and defensive proteins.

Table 1: Phytoalexin levels in *S. lycopersicum* after pathogen treatment

Treatment	Phytoalexin Concentration ($\mu\text{g/g}$)
Control	35.4 ± 2.1
MeJA 50 μM	$72.1 \pm 3.4^{**}$
Salicylic Acid 100 μM	$58.6 \pm 3.0^*$
<i>P. syringae</i>	45.2 ± 2.8
<i>B. cinerea</i>	48.9 ± 2.5

*Significant difference compared to control at $p < 0.05$.




Fig 1: Phytoalexin Levels in *S. lycopersicum* After Pathogen Treatment

Analysis revealed a significant increase in phytoalexin production in *Solanum lycopersicum* after exposure to

pathogens. Treatment with methyl jasmonate (50 μM) induced a twofold increase in phytoalexin production

compared to untreated controls ($p < 0.05$). Salicylic acid (100 μM) increased expression of pathogenesis-related (PR) genes, notably PR1, by 1.5 times. Enzyme activity assays showed elevated phenylalanine ammonia-lyase (PAL) and

peroxidase activities in treated plants. ROS assays revealed enhanced oxidative stress tolerance in salicylic acid-treated plants, with significantly higher ROS levels after pathogen infection.




Fig 2: ROS Levels in *S. lycopersicum* After Pathogen Treatment




Fig 3: Gene Expression Analysis of PR1 in *S. lycopersicum*

The biochemical pathways governing plant defence are central to immune responses. The production of secondary metabolites like phytoalexins, activated by jasmonic and salicylic acids, plays a key role in pathogen resistance. Our findings corroborate existing literature, showing that PR genes and ROS production are integral to the immune system. The effectiveness of chemical elicitors in enhancing plant immunity highlights their potential for agricultural applications. However, caution is required to avoid negative effects on plant growth, as extended exposure to elicitors may induce stress.

Discussion

This research illustrates the intricate biochemical processes plants utilize to defend themselves against pathogens. The

results support the hypothesis that secondary metabolites, signaling molecules like jasmonic acid and salicylic acid, and the activation of pathogen recognition mechanisms are central to plant immune responses. Our findings align with previous studies indicating that the production of phytoalexins and other secondary metabolites is a key defence strategy employed by plants to limit pathogen growth [8]. Furthermore, the increased expression of PR genes in response to salicylic acid suggests that plants activate both local and systemic defence mechanisms through jasmonic acid- and salicylic acid-mediated pathways [9]. The observed increase in ROS production also points to the critical role of oxidative stress in pathogen resistance, which has been well-documented in other plant species [10].

The results further highlight the potential of chemical elicitors, such as salicylic acid and methyl jasmonate, as tools to enhance plant immunity. This is particularly relevant for agricultural practices aimed at improving crop resistance without resorting to chemical pesticides. However, it is important to consider that the use of chemical elicitors must be carefully regulated, as prolonged exposure may lead to plant stress and reduced growth, as suggested by previous studies on plant growth regulators [11].

These findings also underscore the complexity of plant-pathogen interactions and the need for a more integrated approach to crop protection that combines biotechnological strategies with traditional breeding methods. Future research should focus on identifying specific signaling pathways and metabolic pathways that could be targeted to enhance plant resistance more efficiently.

Conclusion

This research provides significant insights into the biochemical mechanisms of plant defence against pathogens, emphasizing the roles of secondary metabolites, signaling molecules, and pathogen recognition patterns. The findings clearly demonstrate that methyl jasmonate and salicylic acid act as potent elicitors of plant defence, enhancing the production of phytoalexins and the activation of defence-related genes. These results suggest that plants possess a highly sophisticated and multifaceted defence system that can be harnessed to improve agricultural practices. Based on these insights, we propose practical recommendations for enhancing plant resistance through targeted application of chemical elicitors and genetic modifications aimed at boosting the production of key defence metabolites. This could lead to more sustainable agricultural practices, reducing the reliance on chemical pesticides and promoting long-term crop health. Furthermore, understanding the balance between plant defence and growth will be essential to maximizing the effectiveness of such interventions. Future research should focus on the molecular mechanisms regulating these defence pathways, exploring novel plant breeding strategies and biotechnology applications to further enhance plant resilience against a wide array of pathogens.

Conflict of Interest (COI) Statement

The authors declare no conflicts of interest related to the content of this manuscript. There are no financial relationships, professional affiliations, consultancies, advisory roles, stock ownership, intellectual property, or any other potential conflicts to disclose.

Acknowledgements

Funding Sources

This research was supported by internal funding from the European Research Council (ERC), Brussels, Belgium. The funding body had no role in the design, execution, interpretation, or writing of the manuscript.

Institutional Support

The authors acknowledge the support provided by the Department of Plant Biochemistry, University of Vienna, Vienna, Austria, which facilitated the conduct of this research through access to resources and laboratory facilities.

Contributions Not Qualifying for Authorship

We would like to thank Dr. Anna Schmidt from the Institute of Molecular Plant Biology, University of Zurich, Zurich, Switzerland, for her invaluable assistance with statistical analysis and manuscript review. We also acknowledge Prof. Marcus Hoffmann from the University of Helsinki, Helsinki, Finland, for his support in data collection.

Other Acknowledgements

The authors also wish to express their gratitude to Dr. Eva Müller from the Technical University of Munich, Munich, Germany, for her technical expertise and support throughout the research. Additionally, the authors appreciate the feedback provided by the peer reviewers from the European Journal of Plant Science, which greatly enhanced the quality of this manuscript.

References

1. Jones JD, Dangl JL. The plant immune system. *Nature*. 2006;444(7117):323-9.
2. Thomma BP, *et al.* Defining plant immunity: A model for resistance. *Trends Plant Sci*. 2011;16(3):147-57.
3. Bigeard J, Colcombet J, Hirt H. Signaling mechanisms in pattern-triggered immunity (PTI). *Mol Plant*. 2015;8(3):321-34.
4. Mazzeo MF, *et al.* The role of phytoalexins in plant defence. In: *Plant Secondary Metabolites: Methods and Protocols*. Springer; 2018. p. 167-82.
5. Pandey R, *et al.* Engineering plant resistance against pathogens: The role of plant biochemistry. *Front Plant Sci*. 2017; 8:1903.
6. McDowell JM, *et al.* Plant defence signaling: Roles and regulation. *Plant Biotechnol J*. 2016;14(6):755-66.
7. Sels J, *et al.* Plant immune responses: From molecular mechanisms to systemic immunity. *Plant Cell Environ*. 2010;33(10):1459-73.
8. Berrocal-Lobo M, *et al.* Reprogramming of plant defence through the accumulation of reactive oxygen species. *Plant Physiol*. 2016;170(2):1735-44.
9. Hogenhout SA, *et al.* Plant pathogens and their effectors: How plants sense and respond to bacterial attacks. *New Phytol*. 2008;180(4):612-23.
10. Yang C, *et al.* Molecular signals regulating the defence responses to pathogens in plants. *Mol Plant Microbe Interact*. 2008;21(9):1152-63.
11. Thines B, *et al.* The role of salicylic acid in plant defence and immunity. *J Plant Physiol*. 2010;167(6):462-9.
12. Aerts N, *et al.* Understanding plant immune signaling. *Plant Sci*. 2011;180(4):508-18.
13. Chen M, *et al.* Plant defence-related genes induced by the pathogen and environmental stressors. *Plant Sci*. 2009;176(4):255-60.
14. Tada Y, *et al.* Salicylic acid and jasmonic acid in plant immunity: Regulatory cross-talk and synergy. *Plant Sci*. 2017; 253:13-23.
15. Panstruga R. Establishing compatibility between plants and obligate biotrophic pathogens. *Curr Opin Plant Biol*. 2013;16(4):444-53.