


International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
 ISSN Online: 2617-4707
 NAAS Rating: 5.29
 IJABR 2025; 9(5): 1208-1211
www.biochemjournal.com
 Received: 20-04-2025
 Accepted: 22-05-2025

Dr. Maria Lopez
 Department of Pharmaceutical
 Sciences, University of Zurich,
 Zurich, Switzerland

Dr. Markus Schreiber
 Department of Pharmaceutical
 Sciences, University of Zurich,
 Zurich, Switzerland

Free radical research: The role of free radicals in aging and age-related diseases

Maria Lopez and Markus Schreiber

DOI: <https://www.doi.org/10.33545/26174693.2025.v9.i5m.6821>

Abstract

Free radicals, particularly reactive oxygen species (ROS) and reactive nitrogen species (RNS), are byproducts of cellular metabolism that play a significant role in aging and the pathogenesis of age-related diseases. These highly reactive molecules can cause oxidative damage to cellular components, including proteins, lipids, and DNA, contributing to the decline in physiological functions with age. The accumulation of oxidative damage over time is widely believed to be a key factor in aging and the development of various diseases such as Alzheimer's, Parkinson's, cardiovascular diseases, and cancer. The free radical theory of aging posits that free radicals are a primary cause of aging due to their ability to induce cellular damage, which accelerates the aging process. Furthermore, oxidative stress can alter the redox balance within cells, impairing cellular functions and leading to inflammation, mitochondrial dysfunction, and cell death. In recent years, research has focused on developing antioxidant therapies and lifestyle interventions to mitigate the effects of oxidative damage and promote healthy aging. However, the complex relationship between free radicals, aging, and disease remains an area of active investigation. This paper aims to explore the mechanisms through which free radicals contribute to aging and age-related diseases and evaluate current therapeutic approaches aimed at reducing oxidative stress. By understanding the role of free radicals in aging, researchers may uncover novel strategies for extending health span and combating age-associated diseases. This review provides a comprehensive overview of the current literature, highlighting the molecular mechanisms underlying oxidative damage and the potential of antioxidants and other interventions in counteracting the harmful effects of free radicals.

Keywords: Free radicals, aging, oxidative stress, reactive oxygen species, age-related diseases, antioxidants, mitochondria, cellular damage, redox balance, inflammation

Introduction

Free radicals, primarily reactive oxygen species (ROS) and reactive nitrogen species (RNS), are highly reactive molecules that play a crucial role in cellular signaling and metabolism. However, when present in excess, they can cause oxidative damage to essential cellular components such as proteins, lipids, and DNA. This oxidative damage is implicated in the aging process and the development of various age-related diseases, including neurodegenerative diseases, cardiovascular conditions, and cancer [1]. The free radical theory of aging, proposed by Denham Harman in the 1950s, suggests that the accumulation of free radical-induced damage to cells over time accelerates the aging process and the onset of age-related diseases [2]. The theory is supported by evidence showing that oxidative stress, resulting from an imbalance between the production of free radicals and the body's antioxidant defenses, can lead to cellular dysfunction and the aging of tissues [3]. The problem of free radical damage is compounded by mitochondrial dysfunction. Mitochondria, the powerhouses of the cell, generate ROS as a natural byproduct of cellular respiration [4]. However, with age, mitochondrial function declines, leading to an increased production of ROS and further oxidative damage [5]. In addition to their role in aging, free radicals contribute to inflammation and cell death, exacerbating the progression of age-related diseases [6].

The hypothesis that targeting oxidative stress can mitigate the effects of aging has led to the development of various antioxidant therapies. Antioxidants, both endogenous and exogenous, neutralize free radicals, thereby preventing oxidative damage [7]. However, while antioxidants have shown promise in preclinical studies, clinical evidence supporting their

Corresponding Author:
Dr. Maria Lopez
 Department of Pharmaceutical
 Sciences, University of Zurich,
 Zurich, Switzerland

efficacy in slowing aging or preventing age-related diseases is still limited [8]. Thus, the objective of this paper is to review the mechanisms by which free radicals contribute to aging and disease, and to evaluate the potential of antioxidant interventions in promoting healthy aging. The complex relationship between free radicals and aging warrants further research to identify effective therapeutic strategies to combat oxidative damage.

Materials and Methods

Materials

The research utilized a variety of materials to explore the role of free radicals in aging and age-related diseases. The primary materials included cultured human cell lines, biological samples such as blood and tissue from elderly individuals, and various antioxidant compounds for experimental treatments. The human cell lines were chosen for their relevance to age-related diseases such as Alzheimer's and cardiovascular diseases, which are commonly associated with oxidative stress and free radical activity [1, 3]. Blood samples were collected from both elderly individuals and younger controls to analyze the levels of reactive oxygen species (ROS) and antioxidant markers. The antioxidant compounds used in the research included vitamin C, vitamin E, and polyphenolic extracts, which have previously been shown to mitigate oxidative stress and enhance cellular resilience [5, 7].

Methods

The experimental design included both *in vitro* and *in vivo* analyses. *In vitro* experiments involved exposing cultured human cell lines to various concentrations of oxidative

stress-inducing agents, such as hydrogen peroxide (H_2O_2), followed by treatment with antioxidant compounds to assess their protective effects on cell viability and oxidative damage. The cell viability was measured using the MTT assay, a standard method to assess cell metabolic activity [8]. ROS levels were quantified using flow cytometry, with the fluorescent probe DCFDA to detect ROS production in treated and untreated cells [9].

For *in vivo* studies, blood and tissue samples from elderly and control subjects were analyzed for oxidative damage markers such as lipid peroxidation (malondialdehyde levels) and protein oxidation (carbonyl content) [10]. The antioxidant capacity of these samples was evaluated using assays such as the Trolox equivalent antioxidant capacity (TEAC) and ferric reducing ability of plasma (FRAP) [11]. Statistical analysis was performed using ANOVA to compare ROS levels, antioxidant activity, and oxidative damage between the elderly and control groups, with a p-value of < 0.05 considered statistically significant [12]. Regression analysis was also applied to explore the relationship between oxidative stress markers and age-related diseases [13].

Results

The results of the research showed a significant increase in the levels of reactive oxygen species (ROS) in blood samples from elderly individuals compared to the control group, confirming the hypothesis that oxidative stress increases with age [1, 12]. Table 1 shows the mean ROS levels in elderly and control subjects, with elderly individuals exhibiting a 35% higher ROS level ($p < 0.05$). This was consistent with findings from previous studies that demonstrated the relationship between ROS and aging [14].

Table 1: ROS Levels in Blood Samples of Elderly vs. Control Groups

Group	ROS Level (mean $\pm$ SD)	p-value
Elderly	12.7 $\pm$ 2.3	<0.05
Control	9.4 $\pm$ 1.8	

Additionally, antioxidant markers, including vitamin C and vitamin E, were significantly lower in elderly individuals compared to controls, suggesting a decrease in the body's

natural defense mechanisms against oxidative stress with aging [15]. The data in Table 2 shows a 40% reduction in antioxidant levels in elderly subjects ($p < 0.05$).

Table 2: Antioxidant Levels in Blood Samples of Elderly vs. Control Groups

Group	Vitamin C (μ mol/L)	Vitamin E (μ mol/L)	p-value
Elderly	22.4 $\pm$ 4.1	16.3 $\pm$ 3.2	<0.05
Control	38.1 $\pm$ 5.2	27.6 $\pm$ 4.5	

The regression analysis revealed a strong negative correlation ($r = -0.72$) between antioxidant levels and oxidative damage markers, indicating that lower antioxidant capacity is associated with higher levels of oxidative

damage in elderly individuals [16, 17]. Figure 1 depicts this relationship, with a clear downward trend in antioxidant levels as oxidative damage markers increase.




Fig 1: A negative correlation between antioxidant levels and oxidative damage markers was observed in elderly individuals




Fig 2: Antioxidant treatment significantly reduced ROS levels in cultured cells exposed to H_2O_2

Discussion

The findings from this research support the hypothesis that oxidative stress is a major contributor to aging and age-related diseases. The significant increase in ROS levels in elderly individuals, coupled with a decrease in antioxidant markers, suggests that the body's ability to counteract oxidative damage diminishes with age. This aligns with the free radical theory of aging, which posits that the accumulation of oxidative damage over time accelerates the aging process and contributes to the onset of age-related diseases such as Alzheimer's, cardiovascular diseases, and cancer [2, 6].

The regression analysis demonstrated a significant relationship between antioxidant depletion and increased oxidative damage, underscoring the importance of maintaining an effective antioxidant defense system in aging individuals. These findings are consistent with previous studies that have shown reduced antioxidant capacity in the elderly, making them more susceptible to oxidative damage [15, 17].

The results also highlighted the potential of antioxidant treatments, such as vitamin C and E, in mitigating oxidative damage. The reduction in ROS levels observed in cell cultures treated with antioxidants supports the therapeutic use of these compounds in age-related disease management.

However, while preclinical studies have shown promising results, further clinical trials are needed to confirm the efficacy of antioxidants in human populations and determine the optimal dosages for long-term use [8, 9].

In summary, the findings from this research provide compelling evidence of the role of free radicals in aging and age-related diseases. The data suggest that antioxidant supplementation may offer a promising strategy for counteracting oxidative stress, but further research is required to develop effective antioxidant-based therapies.

Conclusion

This research has provided valuable insights into the role of free radicals in aging and age-related diseases. The findings demonstrate that oxidative stress, driven by an imbalance between free radicals and antioxidants, is a key factor in the aging process and the development of various diseases. As individuals age, the body's ability to neutralize free radicals diminishes, leading to increased oxidative damage to cellular structures and tissues. This oxidative damage is implicated in the pathogenesis of several age-related diseases, including neurodegenerative disorders, cardiovascular diseases, and cancer.

Based on these findings, it is recommended that interventions aimed at increasing antioxidant capacity be

explored as potential therapeutic strategies for preventing or delaying the onset of age-related diseases. Antioxidant-rich diets or supplements may help alleviate oxidative stress and enhance the body's defense mechanisms. Additionally, lifestyle modifications, such as regular physical exercise and reduced exposure to environmental stressors, can also contribute to reducing oxidative damage and promoting healthy aging. Furthermore, clinical trials focusing on the long-term efficacy of antioxidant supplementation are necessary to determine the most effective dosages and identify potential risks or side effects. A combination of dietary interventions, exercise, and antioxidant therapy may provide a holistic approach to managing oxidative stress and promoting healthy aging. Ultimately, understanding the complex relationship between free radicals and aging will pave the way for developing targeted therapies that can extend health span and improve the quality of life for the aging population.

Conflict of Interest (COI) Statement

The authors declare no conflicts of interest related to the content of this manuscript. There are no financial relationships, professional affiliations, consultancies, advisory roles, stock ownership, intellectual property, or any other potential conflicts to disclose. If there are any conflicts, they will be disclosed here (e.g., financial relationships, honoraria, consulting fees, etc.).

Acknowledgements

Funding Sources

This research was supported by internal funding from the European Research Council (ERC), Brussels, Belgium. The funding body had no role in the design, execution, interpretation, or writing of the manuscript.

Institutional Support

The authors acknowledge the support provided by the Department of Biochemistry, University of Amsterdam, Amsterdam, The Netherlands, which facilitated the conduct of this research through access to resources and laboratory facilities.

Contributions Not Qualifying for Authorship

We would like to thank Dr. Elisabeth Müller and Mr. Thomas Jensen for their invaluable assistance with data collection, statistical analysis, and manuscript review. Their contributions were essential for the completion of this research but did not meet the criteria for authorship.

Other Acknowledgements

The authors also wish to express their gratitude to Dr. Sarah Lambert from the Institute of Advanced Biotechnology, University of Zurich, Zurich, Switzerland, for her technical expertise and support throughout the research. Additionally, the authors appreciate the feedback provided by the peer reviewers from the International Journal of Biochemistry, which greatly enhanced the quality of this manuscript.

References

- Halliwell B. Free radicals and antioxidants: A personal view. *Nutr Rev.* 1994;52(8):253-265.
- Harman D. Aging: A theory based on free radical and radiation chemistry. *J Gerontol.* 1956;11(3):298-300.
- Stadtman ER, Levine RL. Free radical-mediated oxidation of proteins in aging and disease. *Drug Aging.* 2000;17(3):161-168.
- Murphy MP. How mitochondria produce reactive oxygen species. *Biochem J.* 2009;417(1):1-13.
- Bratic A, Larsson NG. The role of mitochondria in aging. *J Clin Invest.* 2013;123(3):951-957.
- Alessio HM. Exercise-induced oxidative stress. *Med Sci Sports Exerc.* 2000;32(6):1002-1013.
- Beckman KB, Ames BN. The free radical theory of aging matures. *Physiol Rev.* 1998;78(2):547-581.
- Ristow M, Zarse K, Oberbach A, *et al.* Antioxidants prevent health-promoting effects of physical exercise in humans. *Proc Natl Acad Sci U S A.* 2009;106(21):8665-8670.
- Sies H. Oxidative stress: A concept in redox biology and medicine. *Redox Biol.* 2015; 4:180-183.
- Finkel T. The longevity of health. *Cell.* 2007;129(4):855-857.
- Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev.* 2010;4(8):118-126.
- Packer L, Cadenas E, Vina J. Mitochondrial free radical generation, oxidative stress, and aging. *Free Radic Biol Med.* 2001;32(9):1333-1342.
- Poljsak B, Milisav I. Modulation of oxidative stress by antioxidants. In: Tewari D, editor. *Oxidative Stress and Antioxidant Defenses.* 1st ed. Springer; 2017. p. 47-66.
- Cortes-Rojo C, Delgado C, Garcia-Gimenez JL, *et al.* Antioxidants and neuroprotection: Mechanisms of action and role in the prevention of neurodegenerative diseases. *Free Radic Res.* 2020;54(8):698-720.
- Kayani S, Khan MI, Suleria HA, *et al.* Phytochemicals and free radicals: The importance of antioxidants in human health. *J Antioxidant Res.* 2020;9(1):49-59.
- Monnier VM, Boulton AJM, Clark JE, *et al.* The mechanisms of action of advanced glycation end products in diabetic complications. *Diabetes.* 2005;54(4):809-817.
- Singh R, Madhu R, Dhaliwal A. Role of oxidative stress in aging. *Int J Med Sci Public Health.* 2016;5(3):518-524.
- Ray PD, Huang BW, Tsuji Y. Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling. *Cell Signal.* 2012;24(5):981-990.