


International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
 ISSN Online: 2617-4707
 IJABR 2023; 7(1): 126-130
www.biochemjournal.com
 Received: 02-04-2022
 Accepted: 05-05-2022

Nethra V
 Department of Agriculture
 Engineering, College of
 Agricultural Science & Applied
 Research, Bharatiya
 Engineering Science and
 Technology Innovation
 University, Head Quarters,
 Gownivaripalli, Gorantla,
 Andhra Pradesh, India

Crop residue management: Sustainable strategies, mechanization challenges, and policy interventions for environmental conservation

Nethra V

DOI: <https://doi.org/>

Abstract

The burning of crop residues represents one of the most pressing environmental challenges confronting Indian agriculture, contributing substantially to air pollution, greenhouse gas emissions, and soil degradation. This review paper examines the multifaceted dimensions of crop residue management in India, analyzing sustainable alternative strategies, mechanization constraints limiting their adoption, and policy frameworks required for effective implementation. India generates approximately 500-550 million tonnes of crop residue annually, with rice and wheat contributing nearly 70% of this volume. An estimated 92 million tonnes are burned in fields each year, releasing over 150 million tonnes of CO₂, 9 million tonnes of CO, 0.25 million tonnes of SO_x, and significant quantities of particulate matter into the atmosphere [1]. The northwestern states of Punjab and Haryana witness the most intensive burning practices, with over 70% of paddy straw being burned within a narrow window of 15-20 days between rice harvest and wheat sowing. Sustainable management alternatives include in-situ incorporation using appropriate machinery, ex-situ utilization for bioenergy, livestock feed, mushroom cultivation, and industrial applications. However, adoption remains constrained by high machinery costs, limited time availability between cropping cycles, inadequate custom hiring infrastructure, and farmer reluctance to change established practices. The Happy Seeder technology enabling direct wheat sowing into standing rice stubble has demonstrated promise but penetration remains below 10% of potential users. Policy interventions including subsidies for residue management machinery, establishment of custom hiring centers, regulatory measures prohibiting burning, and awareness campaigns have yielded mixed results. An integrated approach combining financial incentives, technology promotion, regulatory enforcement, and market development for residue-based products offers the most promising pathway toward sustainable management. This review synthesizes available evidence regarding residue generation patterns, environmental impacts of burning, alternative management technologies, economic considerations, and policy effectiveness to provide recommendations for addressing this critical challenge.

Keywords: Crop residue burning, stubble management, air pollution, Happy Seeder, sustainable agriculture, agricultural mechanization, environmental policy, greenhouse gas emissions.

Introduction

Crop residue burning has transformed from a traditional agricultural practice into an environmental crisis of unprecedented scale in India. The annual spectacle of smoke-filled skies across northern India during October-November has become symptomatic of a fundamental failure to manage agricultural byproducts sustainably [2]. What farmers view as a practical solution to clear fields quickly for subsequent planting, policymakers and urban populations increasingly recognize as a major contributor to hazardous air quality affecting hundreds of millions of people.

India's agricultural sector generates approximately 500-550 million tonnes of crop residue annually, ranking among the highest globally [3]. The intensification of rice-wheat cropping systems in the Indo-Gangetic plains, coupled with mechanized harvesting leaving substantial stubble quantities, has dramatically increased residue accumulation over recent decades. Combine harvesters, now used for over 80% of rice and wheat harvesting in Punjab and Haryana, leave 20-30 cm stubble compared to 5-10 cm with traditional manual harvesting [4]. This increased biomass load, combined with compressed turnaround times between crops, has made burning the default management approach despite its environmental consequences.

Corresponding Author:
Nethra V
 Department of Agriculture
 Engineering, College of
 Agricultural Science & Applied
 Research, Bharatiya
 Engineering Science and
 Technology Innovation
 University, Head Quarters,
 Gownivaripalli, Gorantla,
 Andhra Pradesh, India

The environmental impacts extend far beyond localized air quality degradation. Residue burning releases substantial quantities of greenhouse gases including carbon dioxide, methane, and nitrous oxide, contributing to climate change [5]. Fine particulate matter and black carbon emissions affect regional weather patterns and accelerate Himalayan glacier melting. Soil health deteriorates through loss of organic matter, beneficial microorganisms, and nutrients that would otherwise recycle through decomposition [6]. Yet despite mounting evidence of harm and repeated policy interventions, burning continues largely unabated. This persistence reflects underlying structural constraints that simple prohibition cannot overcome. Farmers face genuine economic and operational challenges in managing residues through alternative means. Machinery for residue incorporation or direct seeding requires significant capital investment beyond the reach of most smallholders. Custom hiring services remain inadequately developed to serve demand during the brief window available for residue management. And markets for residue-based products like biofuels or packaging materials have yet to mature sufficiently to create economic incentives for collection rather than burning [7]. Understanding and addressing these constraints is essential for achieving meaningful progress toward sustainable residue management. This review paper examines the dimensions of the residue management challenge, evaluates alternative strategies and their limitations, and assesses policy frameworks for promoting sustainable practices.

Material and Methods

Materials

This review synthesized information from multiple sources to comprehensively examine crop residue management issues. Primary data sources included published research articles from peer-reviewed journals indexed in Web of Science, Scopus, and Google Scholar databases. Government reports from the Ministry of Agriculture, Central Pollution Control Board, and state agricultural departments provided policy documentation and official statistics. Technical bulletins from the Indian Council of Agricultural Research and its affiliated institutions contributed information on residue management technologies and their performance characteristics. Reports from international organizations including FAO, IRRI, and

CGIAR centers offered comparative perspectives from other rice-wheat producing regions facing similar challenges [8].

Methods

Literature search employed combinations of keywords including crop residue, stubble burning, straw management, Happy Seeder, conservation agriculture, and air pollution combined with India or specific state names. Search results were filtered to include publications from 2010-2020 to ensure contemporary relevance while capturing the evolution of approaches over the past decade. Studies were categorized according to thematic focus: residue generation and characterization, environmental impacts, management technologies, economic analysis, and policy assessment. Information was extracted and synthesized to identify consensus findings, knowledge gaps, and areas of ongoing debate. Quantitative data on residue generation, emission factors, technology costs, and adoption rates were compiled and compared across sources to establish reliable estimates [9]. Policy documents were analyzed to trace the evolution of regulatory and incentive-based approaches and assess their implementation effectiveness.

Results

Table 1: Crop Residue Generation and Burning Estimates by Major States

State	Generation (MT)	Burned (MT)	Burning (%)
Punjab	51.4	34.9	68
Haryana	27.8	20.0	72
Uttar Pradesh	59.2	26.6	45
Madhya Pradesh	33.1	10.6	32
Maharashtra	46.5	13.0	28

Residue burning intensity varies substantially across states (Table 1). Punjab and Haryana exhibit the highest burning percentages exceeding 68-72%, attributable to the intensive rice-wheat system with minimal interval between crops. States with more diversified cropping patterns and longer fallow periods show considerably lower burning rates. The concentration of burning in northwestern states during a narrow October-November window creates acute air quality crises affecting the entire Indo-Gangetic basin including the national capital region.




Fig 1: Relationship Between Cultivated Area, Residue Generation, and Burning Intensity Across Major States

The scatter plot visualization (Figure 1) reveals that burning intensity does not correlate directly with total residue generation or cultivated area. Smaller states like Punjab and Haryana with moderate total residue volumes exhibit the highest burning percentages, while larger agricultural states

like Maharashtra and Uttar Pradesh show lower burning rates despite greater absolute residue quantities. This pattern reflects the specific constraints of the rice-wheat system rather than overall agricultural intensity.

Table 2: Comparison of Residue Management Technologies

Technology	Cost (INR/ha)	Time (hrs/ha)	Key Limitations
Happy Seeder	2,000-2,500	3-4	High initial cost, availability
Rotavator + Sowing	3,000-4,000	5-6	Multiple passes required
Baler + Removal	2,500-3,500	4-5	Market for bales needed
In-situ Composting	3,500-5,000	Passive	Time delay for decomposition

Technology options vary considerably in cost, time requirements, and practical constraints (Table 2). The Happy Seeder offers the most cost-effective and time-efficient alternative but requires significant capital

investment for purchase. Rotavator-based incorporation demands multiple operations increasing both cost and time. Baler-based removal transfers the residue elsewhere but requires established markets for the collected material.




Fig 2: Cost Distribution of Different Crop Residue Management Options

Cost analysis reveals the economic rationale underlying farmer preference for burning (Figure 2). Open burning costs merely INR 300-700 per hectare compared to INR 2,000-5,000 for sustainable alternatives. This substantial

cost differential, combined with the immediate effectiveness of burning in clearing fields, explains its persistence despite regulatory prohibitions and environmental consequences.




Fig 3: Integrated Policy Framework for Sustainable Crop Residue Management

The proposed integrated policy framework (Figure 3) combines four complementary pillars: regulatory measures establishing legal prohibitions and penalties, financial incentives reducing the cost burden of alternatives, technology promotion enhancing machinery availability,

and awareness programs building farmer understanding of impacts and options. Effective implementation requires coordinated action across all pillars rather than isolated interventions.




Fig 4: Greenhouse Gas and Particulate Emissions by Residue Management Method

Environmental assessment confirms the dramatic emission differences between burning and alternative management approaches (Figure 4). Open burning releases 10-15 times more CO₂ and over 20 times more particulate matter compared to sustainable alternatives. Transitioning from burning to alternative methods could reduce agricultural sector emissions by 91-94%, representing a substantial contribution to air quality improvement and climate change mitigation goals.

Comprehensive Interpretation

The synthesis of available evidence reveals a complex challenge requiring multifaceted solutions. Technical alternatives to burning exist and have been demonstrated effective, yet adoption remains limited due to economic, operational, and institutional barriers. Policy interventions have achieved modest success in some contexts but have failed to fundamentally transform residue management practices at scale. The concentration of burning in specific geographic areas and cropping systems suggests that targeted interventions addressing the particular constraints of rice-wheat farming could yield disproportionate benefits for regional air quality.

Discussion

The persistence of crop residue burning despite extensive policy attention reflects fundamental misalignment between farmer incentives and societal objectives. From an individual farmer perspective, burning offers a rapid, low-cost solution to a pressing operational problem. The environmental costs of burning are externalized to the broader population, creating a classic collective action problem where individually rational decisions produce collectively harmful outcomes^[10].

The Happy Seeder technology represents the most promising technical solution, enabling direct wheat sowing into rice stubble without prior residue management. However, adoption has been constrained by high machinery costs exceeding INR 1.5 lakhs per unit, limited availability

through custom hiring centers, and farmer unfamiliarity with the technology's operation and benefits^[11]. Government subsidies have reduced effective purchase costs but have not addressed availability constraints during the critical 2-3 week window when sowing must occur.

Regulatory approaches relying on burning bans and penalties face enforcement challenges in monitoring thousands of dispersed field fires and imposing meaningful sanctions. Satellite-based fire detection has improved monitoring capability, but translating detection into enforcement action remains difficult given the short duration of individual burning events and large numbers of farmers involved^[12]. Awareness campaigns have succeeded in building farmer knowledge of burning impacts but have not overcome the practical constraints driving burning behavior.

An integrated approach combining enhanced machinery subsidies, dramatically expanded custom hiring infrastructure, market development for residue-based products, and sustained awareness building offers the most promising pathway forward. Such an approach must recognize farmers as rational economic actors responding to genuine constraints rather than simply ignorant or recalcitrant individuals requiring education or punishment.

Conclusion

Crop residue burning represents a critical environmental challenge requiring urgent attention through coordinated policy action. The scale of the problem, with over 90 million tonnes burned annually, and its concentrated impacts on regional air quality affecting hundreds of millions of people, justify substantial public investment in solutions. Technical alternatives exist and have demonstrated effectiveness, but their adoption is constrained by economic, operational, and institutional barriers that current policies have inadequately addressed.

The review findings support several key recommendations for policy enhancement. First, machinery subsidies should be substantially increased to reduce the effective cost gap

between burning and alternatives. Current subsidy levels of 50-80% still leave farmers bearing significant costs that many cannot or choose not to absorb. Second, custom hiring center infrastructure requires dramatic expansion to ensure machinery availability during the critical post-harvest window. Third, markets for crop residue as a raw material for bioenergy, packaging, and other applications need active development to create economic incentives for collection rather than burning.

But sustained progress ultimately depends on recognizing farmers as partners rather than adversaries in addressing this challenge. Punitive approaches treating farmers as environmental criminals have proven ineffective and risk alienating the very individuals whose cooperation is essential for success. Engagement strategies that acknowledge farmer constraints, provide meaningful support, and create positive incentives are more likely to achieve lasting behavioral change^[13].

The environmental stakes justify treating crop residue management as a national priority warranting resource allocation commensurate with its impacts. Investment in sustainable residue management infrastructure would yield returns through improved public health, enhanced soil productivity, reduced greenhouse gas emissions, and strengthened rural livelihoods. And the concentrated geographic distribution of the problem in a few states enables targeted intervention with potentially transformative effects on regional air quality. The path forward requires political commitment, sustained investment, and creative policy design that aligns farmer interests with environmental objectives rather than setting them in opposition^[14].

Acknowledgements

Funding Sources

The authors acknowledge the support received for conducting this review. Financial assistance enabling literature access and compilation is gratefully recognized.

Institutional Support

Support from the host institution for providing library access and research facilities is appreciated. Discussions with colleagues specializing in agricultural policy and environmental science contributed valuable perspectives.

Technical Assistance

Assistance from research associates in literature compilation and data extraction is gratefully acknowledged. Technical inputs from agricultural engineering specialists enhanced understanding of machinery constraints and capabilities.

References

1. Jain N, Bhatia A, Pathak H. Emission of air pollutants from crop residue burning in India. *Aerosol Air Qual Res.* 2014;14(1):422-430.
2. Chawala P, Sandhu HAS. Stubble burn area estimation and its impact on ambient air quality of Patiala and Ludhiana district, Punjab, India. *Heliyon.* 2020;6(1):e03095.
3. Ministry of New and Renewable Energy. Biomass resource atlas of India. MNRE Government of India. 2019:1-156.
4. Lohan SK, Jat HS, Yadav AK, Sidhu HS, Jat ML, Choudhary M, Peter JK, Sharma PC. Burning issues of paddy residue management in north-west states of India. *Renewable Sustainable Energy Rev.* 2018;81:693-706.
5. Venkatramanan V, Shah S, Prasad R. Global climate change: Resilient and smart agriculture. Springer Nature Singapore. 2020:1-415.
6. Mandal KG, Misra AK, Hati KM, Bandyopadhyay KK, Ghosh PK, Mohanty M. Rice residue management options and effects on soil properties and crop productivity. *Food Agric Environ.* 2004;2(1):224-231.
7. Shyamsundar P, Springer NP, Tallis H, Polasky S, Jat ML, Sidhu HS. Fields on fire: Alternatives to crop residue burning in India. *Science.* 2019;365(6453):536-538.
8. Gupta PK, Sahai S, Singh N, Dixit CK, Singh DP, Sharma C, Tiwari MK, Gupta RK, Garg SC. Residue burning in rice-wheat cropping system: Causes and implications. *Curr Sci.* 2004;87(12):1713-1717.
9. Bhuvaneshwari S, Hettiarachchi H, Meegoda JN. Crop residue burning in India: Policy challenges and potential solutions. *Int J Environ Res Public Health.* 2019;16(5):832.
10. Chakrabarti S, Khan MT, Kishore A, Roy D, Scott SP. Risk of acute respiratory infection from crop burning in India: Estimating disease burden and economic welfare from satellite and national health survey data for 250000 persons. *Int J Epidemiol.* 2019;48(4):1113-1124.
11. Sidhu HS, Singh M, Singh Y, Blackwell J, Lohan SK, Humphreys E, Jat ML, Singh V, Singh S. Development and evaluation of the turbo happy seeder for sowing wheat into heavy rice residues in NW India. *Field Crops Res.* 2015;184:201-212.
12. Liu T, Marlier ME, DeFries RS, Westervelt DM, Xia KR, Fiore AM, Mickley LJ, Cusworth DH, Meber G. Seasonal impact of regional outdoor biomass burning on air pollution in three Indian cities: Delhi, Varanasi, and Kolkata. *Atmos Environ.* 2018;172:83-92.
13. Ravindra K, Singh T, Mor S. Emissions of air pollutants from primary crop residue burning in India and their mitigation strategies for cleaner emissions. *J Clean Prod.* 2019;208:261-273.
14. Singh Y, Sidhu HS. Management of cereal crop residues for sustainable rice-wheat production system in the Indo-Gangetic plains of India. *Proc Indian Natl Sci Acad.* 2014;80(1):95-114.