

International Journal of Agriculture Extension and Social Development

Volume 6; Issue 1; Jan-Jun 2023; Page No. 123-128

Received: 18-11-2023
Accepted: 25-12-2023

Indexed Journal
Peer Reviewed Journal

Soil erosion and its effect on agricultural sustainability

Pallavi CN

Department of Soil Science and Agricultural Chemistry, College of Agricultural Science & Applied Research, Bharatiya Engineering Science and Technology Innovation University, Head Quarters, Gownivaripalli, Gorantla, Andhra Pradesh, India

DOI: <https://www.doi.org/10.33545/26180723.2023.v6.i1b.2993>

Corresponding Author: Pallavi CN

Abstract

Soil erosion threatens agricultural sustainability by removing fertile topsoil and depleting the nutrient capital essential for crop production. This five-year research quantified soil loss rates and associated nutrient depletion under different land management practices on sloping agricultural land. Erosion monitoring employed standard runoff plots installed across five slope classes ranging from 2% to over 20% gradient. Five management treatments were evaluated: bare fallow (control), conventional tillage, reduced tillage, no-till with residue retention, and contour cultivation with cover crops. Soil loss was measured following each erosive rainfall event, with sediment samples analyzed for nitrogen, phosphorus, potassium, and organic carbon content. Results revealed dramatic differences in erosion rates among management practices. Bare fallow plots experienced annual soil losses increasing from 42.8 t/ha in year one to 54.7 t/ha by year five, substantially exceeding the tolerable soil loss threshold of 11 t/ha/year. Conventional tillage reduced losses to 28-37 t/ha annually but still remained above tolerance levels. Conservation practices achieved progressively greater erosion control: reduced tillage declined from 18.6 to 13.2 t/ha over the five years, no-till with residue retention from 12.4 to 6.4 t/ha, and contour cultivation with cover crops from 8.6 to 3.8 t/ha. Nutrient losses paralleled soil loss patterns with strong correlations between erosion magnitude and nutrient depletion. Nitrogen losses on steep slopes exceeding 20% gradient reached 72.6 kg/ha annually under conventional management, representing substantial fertilizer replacement costs. Phosphorus and potassium losses showed similar patterns, with steep slopes losing 18.8 kg P/ha and 42.4 kg K/ha respectively. Organic carbon losses proved particularly concerning, reaching 1, 148 kg C/ha annually on steep slopes, compromising long-term soil productivity and carbon sequestration potential. Slope gradient emerged as the dominant factor controlling erosion intensity, with losses increasing exponentially above 10% slope. However, conservation practices effectively mitigated slope effects, maintaining erosion below tolerance levels even on steeper terrain. Economic analysis indicated that conservation measures provided positive returns through reduced input requirements and maintained yields despite initial implementation costs. These findings demonstrate that soil erosion poses serious threats to agricultural sustainability on sloping lands, but appropriate conservation practices can effectively control losses while maintaining productivity. Adoption of conservation tillage combined with contour cultivation and cover cropping offers a practical pathway for sustainable land management in erosion-prone agricultural landscapes.

Keywords: Soil erosion, conservation tillage, nutrient loss, slope gradient, runoff plots, sustainable agriculture, cover crops, organic carbon depletion, tolerable soil loss

Introduction

What happens when the very foundation of agricultural production gradually washes away? Soil erosion represents one of the most pervasive threats to long-term agricultural sustainability, silently removing the nutrient-rich topsoil accumulated over millennia ^[1]. Unlike dramatic natural disasters, erosion proceeds incrementally, often escaping notice until productivity declines become unmistakable and remediation costs prohibitive ^[2].

Global estimates suggest that agricultural lands lose approximately 75 billion tonnes of soil annually through water and wind erosion ^[3]. This massive redistribution of earth materials carries profound consequences for food security, as eroded topsoil contains disproportionate concentrations of organic matter, nutrients, and beneficial microorganisms compared to underlying subsoil layers ^[4]. The selective removal of fine particles and organic matter during erosion events further concentrates losses of the most agronomically valuable soil constituents.

Erosion processes operate through several distinct mechanisms depending on rainfall characteristics, slope geometry, and surface conditions. Sheet erosion removes thin layers of soil relatively uniformly across field surfaces during rainfall events ^[5]. Rill erosion concentrates flow into small channels that deepen progressively during storms. Gully erosion represents the most severe form, creating deep incisions that fragment fields and remove massive soil volumes ^[6]. Understanding these processes provides the foundation for designing effective control measures.

Slope gradient stands among the most influential factors determining erosion intensity. Steeper slopes generate faster runoff velocities with greater sediment transport capacity ^[7]. The relationship between slope and erosion follows non-linear patterns, with losses increasing disproportionately once critical thresholds are exceeded. Morgan ^[8] demonstrated that doubling slope gradient can increase erosion rates by factors of three to five depending on soil properties and rainfall characteristics.

Land management practices profoundly influence erosion susceptibility by modifying surface conditions that control runoff generation and sediment detachment. Conventional tillage leaves soil surfaces bare and compacted, maximizing exposure to raindrop impact and overland flow ^[9]. Conservation tillage approaches that maintain crop residues on the surface dramatically reduce erosion by protecting soil from direct raindrop impact and slowing runoff velocity ^[10]. Cover crops provide similar protection during fallow periods while adding organic matter that improves soil structure.

The concept of tolerable soil loss, designated as the T value, establishes upper limits for erosion that can be sustained without compromising long-term productivity ^[11]. For most agricultural soils, tolerable losses range from 5 to 12 tonnes per hectare annually, rates assumed to be balanced by soil formation processes. However, actual formation rates typically fall far below these thresholds, suggesting that even tolerable losses may gradually deplete soil resources over extended timeframes ^[12].

Nutrient losses accompanying eroded soil impose direct economic costs through increased fertilizer requirements to maintain yields. Nitrogen, phosphorus, and potassium removed with sediment must be replaced through external inputs, while organic matter losses reduce the soil's capacity to retain and cycle nutrients ^[13]. These hidden costs often escape farm-level accounting but accumulate substantially over time. Pimentel *et al.* ^[14] estimated that erosion-related nutrient losses cost farmers billions of dollars annually in replacement fertilizers alone.

Beyond on-farm productivity impacts, eroded sediments degrade downstream water resources through siltation of reservoirs and nutrient enrichment of surface waters ^[15]. Phosphorus transported with sediment contributes to eutrophication of lakes and coastal zones, while sediment deposition reduces storage capacity of water supply reservoirs. These off-site impacts generate societal costs that extend far beyond individual farm boundaries.

Climate change projections suggest increasing erosion risks in many agricultural regions as rainfall intensity increases and seasonal patterns shift ^[16]. Higher intensity storms generate greater runoff and erosive energy even without changes in total precipitation. These evolving conditions heighten the urgency of implementing effective erosion control measures on vulnerable agricultural lands.

This research aimed to quantify soil erosion rates and nutrient losses under different land management practices across a range of slope conditions. Specific objectives included comparing erosion control effectiveness of conventional versus conservation management approaches, quantifying nutrient losses associated with different erosion intensities, and evaluating the influence of slope gradient on erosion outcomes under various management systems.

Study Area Description

The research site occupied undulating terrain in a sub-humid subtropical region with pronounced wet and dry seasons. Mean annual precipitation totaled 924 mm, with 78% occurring during the June through September monsoon period when erosion risks peak. Temperature ranged from 8°C mean minimum in January to 38°C mean maximum in May. The landscape comprised rolling hills with slopes

ranging from nearly level to exceeding 25% gradient, representative of extensive agricultural areas in the region. Soils were classified as Typic Haplustalfs (Alfisols) with clay loam texture containing 32% clay, 38% silt, and 30% sand. These soils exhibited moderate erodibility with K-factor values of 0.32 as determined by standard nomograph procedures. Prior land use included rainfed cultivation of cereals and pulses under conventional tillage for over two decades, a history that had already produced visible erosion features including rill networks on steeper slopes.

Materials and Methods

Materials

Runoff plots measuring 22 m length by 4 m width were established on slopes representing five gradient classes: 2-5%, 5-10%, 10-15%, 15-20%, and greater than 20%. Plot boundaries consisted of galvanized metal sheets inserted 15 cm into soil to prevent lateral water movement. Collection troughs at the downslope end directed runoff into calibrated tanks for volume measurement. Sediment collection tanks allowed settling of coarse particles while overflow captured fine suspended sediments. Cover crop seeds included cowpea (*Vigna unguiculata*) and horsegram (*Macrotyloma uniflorum*) for summer and winter seasons respectively. Maize (*Zea mays* var. DHM-117) served as the test crop across all treatments ^[1, 2].

Field Experimental Design

A split-plot design was employed with slope classes as main plots and management treatments as subplots, replicated three times. Five management treatments were evaluated: T₁-bare fallow maintained through periodic herbicide application; T₂-conventional tillage with complete residue incorporation; T₃-reduced tillage with partial residue retention; T₄-no-till with complete residue retention on soil surface; and T₅-contour cultivation combined with cover crop rotation during fallow periods. Plots receiving crop cultivation followed maize-fallow rotation with cultivation during monsoon season and fallow or cover crops during dry months. Conventional tillage involved moldboard plowing to 20 cm depth followed by two passes of disc harrow. Reduced tillage used chisel plow only to 15 cm depth. No-till plots received direct seeding through surface residues without prior tillage ^[3, 4].

Methods

Soil loss measurement followed standard runoff plot methodology. Following each erosive rainfall event exceeding 12.5 mm, runoff volume was recorded from calibrated tanks and sediment concentration determined gravimetrically from representative samples. Total soil loss was calculated as the product of runoff volume and sediment concentration. Sediment samples were composited monthly for nutrient analysis including total nitrogen by micro-Kjeldahl digestion, available phosphorus by Olsen extraction, exchangeable potassium by ammonium acetate extraction, and organic carbon by Walkley-Black wet oxidation ^[5, 6]. Rainfall data were recorded using automatic tipping-bucket gauges at the experimental site. Erosivity indices were calculated for each storm event using standard methodology. Statistical analysis employed analysis of variance for split-plot design with means separation by least

significant difference at 5% probability ^[7].

Soil Quality Analysis

Soil quality assessments were conducted annually using samples collected from the 0-15 cm depth. Physical parameters included bulk density by core method, aggregate stability by wet sieving, and infiltration rate by double-ring infiltrometer. Chemical analyses determined pH, electrical conductivity, organic carbon, available nitrogen,

phosphorus, and potassium following standard protocols. Biological indicators included microbial biomass carbon by fumigation-extraction and dehydrogenase activity by triphenyltetrazolium chloride reduction. These parameters collectively assessed erosion impacts on soil productive capacity beyond simple mass loss measurements ^[8, 9].

Results

Table 1: Annual soil loss (t/ha) under different management treatments over five-year monitoring period

Treatment	Year 1	Year 2	Year 3	Year 4	Year 5
Bare Fallow	42.8	45.2	48.6	51.3	54.7
Conventional Tillage	28.4	30.1	32.8	34.2	36.5
Reduced Tillage	18.6	17.2	15.8	14.6	13.2
No-Till + Residue	12.4	10.8	9.2	7.8	6.4
Contour + Cover	8.6	7.2	5.8	4.6	3.8

Values represent means across all slope classes. Tolerable soil loss (T value) = 11 t/ha/year. Treatment × Year interaction significant at $p < 0.01$.

Soil loss rates differed dramatically among management treatments and showed divergent temporal trends. Bare fallow and conventional tillage exhibited increasing erosion over the five-year period, reflecting progressive soil degradation and increased erodibility. In contrast,

conservation practices showed declining erosion trends as surface protection improved through residue accumulation and soil structure enhancement. By year five, contour cultivation with cover crops achieved 93% reduction in soil loss compared to bare fallow conditions.




Figure 1: Annual soil loss trends under five land management treatments over the five-year experimental period showing divergent trajectories between degrading and improving systems

Table 2: Nutrient losses (kg/ha/year) associated with soil erosion across slope classes under conventional tillage

Slope Class	Soil Loss (t/ha)	N Loss	P Loss	K Loss	OC Loss
2-5%	12.4	8.4	2.1	5.2	124
5-10%	24.6	18.6	4.8	11.4	286
10-15%	38.2	32.4	8.6	19.8	512
15-20%	52.8	48.2	12.4	28.6	768
>20%	68.4	72.6	18.8	42.4	1148

OC = Organic Carbon (kg C/ha/year). Values are five-year means. All correlations between slope and losses significant at $p < 0.001$.




Fig 2: Conceptual diagram illustrating soil erosion types, degradation pathways, agricultural consequences, and conservation approaches for sustainable land management

Nutrient losses increased exponentially with slope gradient, closely tracking soil mass losses. Nitrogen losses reached 72.6 kg/ha annually on slopes exceeding 20%, equivalent to substantial fertilizer inputs. Organic carbon losses proved

particularly severe on steep slopes, with over one tonne of carbon removed per hectare annually. These losses not only reduce current soil fertility but compromise long-term productive capacity and carbon sequestration potential.



Figure 3: Stacked bar chart showing nutrient losses (N, P, K, and organic carbon) associated with soil erosion across five slope gradient classes under conventional tillage management

Comprehensive Interpretation

The interaction between management practice and slope gradient revealed that conservation measures effectively mitigated topographic effects on erosion. While

conventional tillage produced soil losses exceeding tolerance levels on all slopes above 5%, contour cultivation with cover crops maintained losses below tolerance even on the steepest slopes monitored. This demonstrates that

appropriate management can extend productive cultivation to slopes that would otherwise require retirement from agriculture. Economic calculations indicated that conservation practices generated positive returns beginning in year three through reduced fertilizer requirements and maintained yields, despite initial establishment costs. The cumulative value of retained nutrients and organic matter substantially exceeded implementation expenses over the five-year period, supporting economic viability of conservation adoption.

Discussion

The erosion rates documented under conventional management substantially exceeded tolerable limits, confirming serious sustainability concerns for sloping agricultural lands under current practices. Lal ^[17] reported comparable erosion rates of 25-50 t/ha annually on similar slopes under conventional cultivation in tropical regions, suggesting the present findings reflect widespread conditions rather than site-specific anomalies.

The progressive increase in erosion under bare fallow and conventional tillage contrasted with declining trends under conservation practices illustrates divergent soil health trajectories. Degrading systems lose structural integrity as organic matter declines, increasing erodibility and runoff generation in a self-reinforcing cycle ^[18]. Conservation systems interrupt this cycle through surface protection and organic matter accumulation that improve infiltration and aggregate stability over time.

Nutrient enrichment ratios in eroded sediment exceeded those of source soils by factors of 1.5 to 3.0, confirming selective removal of fine particles and organic matter during erosion events. This enrichment effect amplifies productivity impacts beyond what simple mass loss figures would suggest ^[19]. The topsoil fraction removed during erosion contains disproportionate shares of the nutrients, organic matter, and biological activity that support crop production.

Organic carbon losses merit particular concern given the lengthy timeframes required for accumulation and the multiple soil functions that organic matter supports. Montgomery ^[12] estimated that replacing eroded organic matter through natural accumulation processes would require centuries to millennia, effectively rendering erosion losses permanent from a management perspective. The over 1,000 kg C/ha annual losses on steep slopes under conventional management represent significant depletion of this irreplaceable resource.

The effectiveness of combined conservation practices exceeded expectations based on individual practice contributions, suggesting synergistic interactions. Contour cultivation reduces runoff velocity and increases infiltration opportunity, while cover crops provide surface protection and organic matter inputs ^[20]. Their combination addresses multiple erosion mechanisms simultaneously, achieving control levels that neither practice alone could attain.

Some limitations of this research should be acknowledged. The five-year duration captured treatment establishment effects but may not fully represent long-term equilibrium conditions. Extrapolation to soil types with different erodibility characteristics requires caution. And the specific conservation practice combinations evaluated may require

adaptation for different cropping systems and regional conditions.

Policy implications arising from these findings support stronger promotion of conservation practices on erosion-prone lands. The demonstrated economic viability removes one barrier to adoption, but farmers may still require technical assistance and possibly initial cost-sharing to implement unfamiliar practices ^[21]. Watershed-level planning that considers both on-farm productivity and downstream sediment impacts would optimize societal benefits from erosion control investments.

Conclusion

This five-year investigation demonstrated that soil erosion on sloping agricultural land poses serious threats to sustainability under conventional management but can be effectively controlled through appropriate conservation practices. The research quantified erosion rates ranging from 3.8 t/ha annually under combined conservation measures to 54.7 t/ha under bare fallow conditions, representing more than ten-fold differences attributable to management choices.

Conventional tillage failed to maintain erosion below tolerable limits on any slope class above 5% gradient, indicating that current practices cannot sustain productivity on substantial portions of the agricultural landscape. In contrast, conservation practices including reduced tillage, no-till with residue retention, and contour cultivation with cover crops achieved progressive erosion reductions that brought losses below tolerance levels on all slope classes evaluated.

Nutrient losses accompanying erosion reached levels that impose substantial economic costs through fertilizer replacement requirements. Nitrogen losses of 72.6 kg/ha annually on steep slopes exceed typical fertilizer application rates, meaning that erosion can remove more nutrients than farmers apply. Phosphorus and potassium losses similarly represent significant depletion of soil fertility capital that accumulated over geological timeframes.

Organic carbon losses proved particularly concerning given the central role of organic matter in soil productivity and the lengthy timeframes required for accumulation. Annual losses exceeding 1,000 kg C/ha on steep slopes under conventional management represent irreversible depletion from a practical management perspective. Conservation practices that reduced erosion simultaneously protected this critical soil component.

The temporal trends observed carry important implications for long-term sustainability. Conventional systems showed increasing erosion over time as soil structure degraded, suggesting accelerating decline without intervention. Conservation systems exhibited opposite trends with declining erosion as surface protection and soil quality improved, indicating capacity for continued productivity improvement with sustained practice implementation.

Practical recommendations from this research include adopting conservation tillage as a minimum erosion control measure on all sloping lands, implementing contour cultivation on slopes exceeding 5% gradient, and establishing cover crops during fallow periods to maintain continuous surface protection. These practices can be combined progressively as farmers gain experience and

confidence with conservation approaches.

Future research should evaluate conservation practice performance across broader ranges of soil types and climate conditions, develop improved methods for assessing erosion impacts on long-term productivity, and investigate economic incentive mechanisms for promoting conservation adoption among risk-averse farmers. These efforts would strengthen the evidence base supporting sustainable land management on erosion-prone agricultural landscapes.

References

1. Pimentel D. Soil erosion: a food and environmental threat. *Environment, Development and Sustainability*. 2006;8(1):119-137.
2. Bakker MM, Govers G, Rounsevell MDA. The crop productivity-erosion relationship: an analysis based on experimental work. *Catena*. 2004;57(1):55-76.
3. Quinton JN, Govers G, Van Oost K, Bardgett RD. The impact of agricultural soil erosion on biogeochemical cycling. *Nature Geoscience*. 2010;3(5):311-314.
4. Gregorich EG, Greer KJ, Anderson DW, Liang BC. Carbon distribution and losses: erosion and deposition effects. *Soil and Tillage Research*. 1998;47(3-4):291-302.
5. Wischmeier WH, Smith DD. Predicting rainfall erosion losses: a guide to conservation planning. *USDA Agriculture Handbook*. 1978;537:1-58.
6. Poesen J, Nachtergaele J, Verstraeten G, Valentin C. Gully erosion and environmental change: importance and research needs. *Catena*. 2003;50(2-4):91-133.
7. Liu BY, Nearing MA, Risse LM. Slope gradient effects on soil loss for steep slopes. *Transactions of the ASAE*. 1994;37(6):1835-1840.
8. Morgan RPC. *Soil erosion and conservation*. 3rd ed. Blackwell Publishing. 2005:1-304.
9. Rhoton FE, Shipitalo MJ, Lindbo DL. Runoff and soil loss from midwestern and southeastern US silt loam soils as affected by tillage practice and soil organic matter content. *Soil and Tillage Research*. 2002;66(1):1-11.
10. Scopel E, Triomphe B, Affholder F, Da Silva FAM, Corbeels M, Xavier JHV. Conservation agriculture cropping systems in temperate and tropical conditions, performances and impacts: a review. *Agronomy for Sustainable Development*. 2013;33(1):113-130.
11. Renard KG, Foster GR, Weesies GA, McCool DK, Yoder DC. Predicting soil erosion by water: a guide to conservation planning with the Revised Universal Soil Loss Equation (RUSLE). *USDA Agriculture Handbook*. 1997;703:1-404.
12. Montgomery DR. Soil erosion and agricultural sustainability. *Proceedings of the National Academy of Sciences*. 2007;104(33):13268-13272.
13. Hao Y, Lal R, Owens LB, Izaurralde RC, Post WM, Hothem DL. Effect of cropland management and slope position on soil organic carbon pool at the North Appalachian Experimental Watersheds. *Soil and Tillage Research*. 2002;68(2):133-142.
14. Pimentel D, Harvey C, Resosudarmo P, Sinclair K, Kurz D, McNair M. Environmental and economic costs of soil erosion and conservation benefits. *Science*. 1995;267(5201):1117-1123.
15. Carpenter SR, Caraco NF, Correll DL, Howarth RW, Sharpley AN, Smith VH. Nonpoint pollution of surface waters with phosphorus and nitrogen. *Ecological Applications*. 1998;8(3):559-568.
16. Nearing MA, Pruski FF, O'Neal MR. Expected climate change impacts on soil erosion rates: a review. *Journal of Soil and Water Conservation*. 2004;59(1):43-50.
17. Lal R. Soil degradation by erosion. *Land Degradation and Development*. 2001;12(6):519-539.
18. Le Bissonnais Y. Aggregate stability and assessment of soil crustability and erodibility: I. Theory and methodology. *European Journal of Soil Science*. 1996;47(4):425-437.
19. Jacinthe PA, Lal R, Kimble JM. Carbon budget and seasonal carbon dioxide emission from a central Ohio Luvisol as influenced by wheat residue amendment. *Soil and Tillage Research*. 2002;67(2):147-157.
20. Dabney SM, Delgado JA, Reeves DW. Using winter cover crops to improve soil and water quality. *Communications in Soil Science and Plant Analysis*. 2001;32(7-8):1221-1250.
21. Knowler D, Bradshaw B. Farmers' adoption of conservation agriculture: a review and synthesis of recent research. *Food Policy*. 2007;32(1):25-48.