






International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
IJABR 2022; 6(2): 198-203
www.biochemjournal.com
Received: 26-09-2022
Accepted: 28-10-2022

Dr. Chidanandh AR
Department of Biotechnology,
College of Agricultural Science
and Applied Research,
Bharatiya Engineering Science
and Technology Innovation
University, Head Quarters,
Gownivaripalli, Gorantla,
Andhra Pradesh, India

How do tardigrades survive extreme conditions? Studying cryptobiosis mechanisms and applying trehalose-based preservation techniques to vaccine storage and organ transplantation

Chidanandh AR

DOI: <https://www.doi.org/10.33545/26174693.2022.v6.i2c.7262>

Abstract

Tardigrades, microscopic eight-legged organisms, demonstrate extraordinary survival capabilities under extreme environmental conditions through cryptobiosis a state of suspended animation. These remarkable creatures withstand desiccation, extreme temperatures, radiation, and pressure that would destroy most life forms. Understanding the molecular mechanisms underlying tardigrade survival offers tremendous potential for biopreservation applications. This research investigates cryptobiosis mechanisms in tardigrades and evaluates trehalose-based preservation strategies inspired by tardigrade biology for vaccine storage and organ transplantation. Through experimental investigations and molecular analyses, we examined survival responses to desiccation, freezing, radiation, and pressure across 120-day exposure periods. Results demonstrate exceptional survival rates: 94.3% following complete desiccation, 89.7% after prolonged freezing, and 78.2% following radiation exposure. Molecular characterization identified trehalose accumulation as central protective mechanism, with intracellular concentrations reaching 18.7% during tun formation. Late embryogenesis abundant proteins showed 12-fold upregulation during cryptobiosis induction. Application testing revealed that trehalose-supplemented preservation solutions significantly enhanced vaccine stability, maintaining 94.7% efficacy at -80 °C and 87.3% at room temperature compared to 73.2% and 31.4% controls respectively. Organ preservation experiments demonstrated that trehalose treatment extended viable storage from 6 hours to 18.4 hours at 4°C while maintaining 83.6% cellular viability. These findings establish trehalose-based biopreservation as promising technology for medical and pharmaceutical applications.

Keywords: Tardigrades, cryptobiosis, trehalose, biopreservation, vaccine storage, organ transplantation, desiccation tolerance, suspended animation, molecular protection, cryopreservation

Introduction

Tardigrades, commonly known as water bears, represent one of nature's most resilient organisms. These microscopic invertebrates inhabit diverse environments from Antarctic ice to tropical rainforests [1]. Their most remarkable characteristic is cryptobiosis the ability to enter suspended animation where metabolic activity becomes virtually undetectable. In this state, tardigrades survive conditions that would instantly kill most organisms, including exposure to space vacuum, extreme radiation, and temperatures approaching absolute zero [2]. Cryptobiosis occurs when tardigrades encounter adverse environmental conditions. As moisture disappears, tardigrades contract into a compact form called a tun, reducing body water content from approximately 85% to less than 3% [3]. They can persist for years in this state, resuming normal activity within hours when water becomes available. This extraordinary survival capacity has captivated researchers seeking to understand and harness these mechanisms for practical applications. The molecular basis of tardigrade survival involves multiple protective mechanisms. Trehalose, a disaccharide sugar, accumulates to extraordinarily high concentrations during tun formation, replacing water molecules around cellular structures and maintaining protein and membrane integrity during dehydration [4]. Additionally, tardigrades express unique proteins including tardigrade-specific intrinsically disordered proteins and late embryogenesis abundant proteins that provide molecular protection [5]. Medical and pharmaceutical industries face persistent biopreservation challenges. Vaccine storage typically requires continuous cold chain maintenance, limiting

Corresponding Author:
Dr. Chidanandh AR
Department of Biotechnology,
College of Agricultural Science
and Applied Research,
Bharatiya Engineering Science
and Technology Innovation
University, Head Quarters,
Gownivaripalli, Gorantla,
Andhra Pradesh, India

distribution to resource-poor regions [6]. Organ transplantation confronts severe time constraints; most organs remain viable for only hours outside the body [7]. Current cryopreservation methods for cells frequently cause ice crystal damage. Inspiration from tardigrade biology offers potential solutions. This research examines tardigrade cryptobiosis mechanisms and translates findings into practical biopreservation applications.

Materials and Methods

Materials

This research was conducted from January 2018 to December 2019. Tardigrade specimens (*Hypsibius dujardini*) were cultured in laboratory conditions at 18°C with algal food supply. Ethical approval was obtained from the institutional ethics committee. Research materials included analytical grade trehalose from commercial suppliers, preservation buffers, vaccine samples from standard immunization programs, porcine organ samples from certified facilities, and human cell lines from established repositories. All materials were obtained following ethical guidelines and standard protocols.

Methods: Tardigrades were subjected to controlled desiccation by gradual humidity reduction over 48 hours. Survival was assessed by rehydration and observation of movement. Molecular analyses used mass spectrometry for trehalose quantification and quantitative RT-PCR for protein

expression analysis. Vaccine preservation involved mixing standard vaccines with trehalose solutions at concentrations ranging from 0-20% and monitoring efficacy over time using enzyme-linked immunosorbent assay. Organ preservation used porcine liver segments treated with trehalose-supplemented solutions, measuring cellular viability through methylthiazolyldiphenyl-tetrazolium bromide assays. Cell cryopreservation compared trehalose-containing versus conventional dimethyl sulfoxide-based freezing protocols. Statistical analyses employed analysis of variance and t-tests using statistical software.

Ethical Considerations

This research adhered to rigorous ethical standards throughout all phases. All animal-derived materials were obtained from approved sources with proper ethical oversight. Vaccine samples were surplus materials from clinical programs, obtained with appropriate permissions. Human cell lines were commercially available with documented ethical provenance. No human subjects were directly involved in this research. Laboratory procedures followed institutional biosafety and animal welfare guidelines. Data management maintained confidentiality and security. The research received institutional ethics committee approval and complied with relevant national and international guidelines for biological research.

Results

Table 1. Tardigrade Survival Rates under Different Stress Conditions

Stress Condition	Duration (days)	Survival Rate (%)	Recovery Time (hours)	n
Complete Desiccation	30	94.3	2.4	120
Complete Desiccation	90	87.6	4.1	120
Freezing (-20°C)	30	89.7	3.2	100
Freezing (-80°C)	30	82.4	5.8	100
Radiation (5000 Gy)	1	78.2	6.3	80
Pressure (600 MPa)	1	71.8	8.7	80

Table 1 presents tardigrade survival rates under various extreme conditions. Complete desiccation for 30 days resulted in 94.3% survival with rapid recovery (2.4 hours average). Extended desiccation (90 days) maintained high survival (87.6%) though recovery time increased. Freezing at -20°C yielded 89.7% survival, while ultra-low

temperature (-80 °C) resulted in 82.4% survival. Radiation exposure at 5000 Gray, a dose lethal to most organisms, produced 78.2% survival. Extreme pressure (600 megapascals) resulted in 71.8% survival. These results demonstrate tardigrades' extraordinary resilience across multiple stress types.




Fig 1: Tardigrade Survival under Different Extreme Conditions over Time

Figure 1 illustrates survival rates across different stress conditions and exposure durations. Desiccation shows gradual decline but maintains high survival even after 120 days. Freezing demonstrates moderate decline with temperature-dependent effects. Radiation and pressure show

steeper initial declines. The data reveal that desiccation tolerance represents tardigrades' most robust survival mechanism, likely reflecting evolutionary adaptation to terrestrial environments where water availability fluctuates.




Fig 2: Preservation Effectiveness at Different Trehalose Concentrations

Figure 2 displays preservation effectiveness across trehalose concentration gradients. Control treatments (0% trehalose)

showed baseline effectiveness of approximately 42%. Addition of 5% trehalose increased effectiveness to 67%,

while 10% achieved 81% effectiveness. Optimal effectiveness (89%) occurred at 15% trehalose concentration. Higher concentrations (20%) showed slight decline to 87%, suggesting potential osmotic stress. These

results indicate that 15% trehalose represents optimal concentration for biopreservation applications, balancing protective benefits against potential negative effects.




Fig 3: Cryptobiosis Mechanism in Tardigrades

Figure 3 illustrates the molecular mechanisms underlying tardigrade cryptobiosis. Normal state maintains 85% water content with active metabolism and normal protein function. Upon stress trigger, tardigrades initiate molecular changes including trehalose synthesis, late embryogenesis abundant protein expression, DNA protection mechanisms, and membrane stabilization. These coordinated responses

produce the tun state characterized by less than 3% water content, suspended metabolism, trehalose-mediated protection, and glass-like cellular state. Rehydration reverses these changes, allowing recovery to normal function. This schematic synthesizes current understanding of cryptobiosis molecular basis.




Fig 4: Trehalose-Based Preservation Effectiveness across Applications

Figure 4 compares preservation effectiveness between control and trehalose treatments across multiple applications. Vaccine storage at -80 °C improved from 73.2% (control) to 94.7% (trehalose treatment). Room temperature vaccine storage showed dramatic improvement from 31.4% to 87.3%. Organ preservation at 4 °C increased from 51.7% to 83.6%. Cell cryopreservation improved from 68.3% to 91.2%. Blood product storage enhanced from 64.1% to 88.9%. These consistent improvements across diverse applications demonstrate trehalose's broad utility for biopreservation, potentially revolutionizing cold chain logistics and expanding access to life-saving medical materials.

Comprehensive Interpretation

Molecular analysis revealed that trehalose accumulation represents the central mechanism of tardigrade cryptobiosis protection. During tun formation, intracellular trehalose concentrations reached 18.7% of dry weight, substantially higher than most organisms. This sugar acts as water replacement, preserving protein structure and membrane integrity during extreme dehydration. Late embryogenesis abundant proteins showed 12-fold upregulation during cryptobiosis induction, providing additional molecular protection. DNA damage repair mechanisms activated during recovery, explaining high survival rates despite radiation exposure. The successful translation of these mechanisms to biopreservation applications validates the biomimicry approach, demonstrating that understanding extreme organism adaptations can yield practical technological innovations with significant societal benefits.

Discussion

This research demonstrates that tardigrade cryptobiosis mechanisms can be effectively applied to biopreservation challenges. The 94.7% vaccine efficacy maintained at -80°C with trehalose treatment represents significant improvement over conventional methods. More dramatically, the 87.3% efficacy at room temperature could revolutionize vaccine distribution in resource-limited settings lacking reliable refrigeration infrastructure. Organ preservation extension from 6 to 18.4 hours significantly expands transplant logistics, potentially reducing organ waste and improving recipient outcomes. The 91.2% post-thaw viability in cell cryopreservation surpasses conventional dimethyl sulfoxide methods, benefiting reproductive medicine and cellular therapy applications. These findings align with previous research demonstrating trehalose's protective properties, while extending applications to practical medical contexts. Future research should optimize delivery methods, investigate long-term stability, and conduct clinical trials to establish regulatory approval pathways.

Limitations

Several limitations warrant acknowledgment. First, laboratory-cultured tardigrades may differ from wild populations in stress tolerance, potentially affecting generalizability. Second, while trehalose showed effectiveness across tested applications, optimal concentrations may vary for specific vaccines, organs, or cell types requiring further optimization. Third, long-term stability testing extended to 120 days; longer-term effects remain unknown. Fourth, the research used porcine organs rather than human organs due to ethical constraints;

translation to human transplantation requires validation. Fifth, economic analysis was limited; comprehensive cost-effectiveness studies comparing trehalose preservation to conventional methods would inform implementation decisions. Finally, regulatory approval pathways for trehalose-based preservation formulations remain undefined, requiring substantial additional work before clinical deployment.

Conclusion

Tardigrade research reveals profound insights into biological resilience and offers practical solutions for biopreservation challenges. The cryptobiosis mechanisms centered on trehalose accumulation and protective proteins demonstrate nature's sophisticated approaches to surviving extreme conditions. Translation of these mechanisms into medical applications shows tremendous promise, with significant improvements in vaccine stability, organ preservation duration, and cell cryopreservation viability. Trehalose-based preservation could revolutionize vaccine distribution to remote regions, extend organ transplant logistics, and improve reproductive medicine outcomes. While challenges remain in optimizing concentrations, delivery methods, and regulatory approval, the fundamental viability of this approach is established. Future research should focus on scaling production, conducting clinical trials, and addressing regulatory pathways. The humble tardigrade, microscopic and often overlooked, may hold keys to solving major challenges in medicine and public health. This research demonstrates that studying extreme organisms can yield practical benefits for humanity, reinforcing the value of basic biological research and biomimicry in technological innovation.

References

1. Nelson DR, Guidetti R, Rebecchi L. Phylum Tardigrada. In: Thorp JH, Covich AP, editors. *Thorp and Covich's Freshwater Invertebrates*. Academic Press; 2015. p. 347-380.
2. Jönsson KI, Rabbow E, Schill RO, Harms-Ringdahl M, Rettberg P. Tardigrades survive exposure to space in low Earth orbit. *Current Biology*. 2008;18(17):R729-R731.
3. Crowe JH, Hoekstra FA, Crowe LM. Anhydrobiosis. *Annual Review of Physiology*. 1992;54:579-599.
4. Elbein AD, Pan YT, Pastuszak I, Carroll D. New insights on trehalose: a multifunctional molecule. *Glycobiology*. 2003;13(4):17R-27R.
5. Hashimoto T, Horikawa DD, Saito Y, Kuwahara H, Kozuka-Hata H, Shin-I T, *et al.* Extremotolerant tardigrade genome and improved radiotolerance of human cultured cells by tardigrade-unique protein. *Nature Communications*. 2016;7:12808.
6. Chen D, Kristensen D. Opportunities and challenges of developing thermostable vaccines. *Expert Review of Vaccines*. 2009;8(5):547-557.
7. Monbaliu D, Brassil J. Machine perfusion of the liver: past, present and future. *Current Opinion in Organ Transplantation*. 2010;15(2):160-166.
8. Crowe JH, Carpenter JF, Crowe LM. The role of vitrification in anhydrobiosis. *Annual Review of Physiology*. 1998;60:73-103.
9. Rebecchi L, Guidetti R, Borsari S, Altiero T, Bertolani R. Dynamics of long-term anhydrobiotic survival of

- lichen-dwelling tardigrades. *Hydrobiologia*. 2006;558:23-30.
10. Wharton DA. *Life at the limits: organisms in extreme environments*. Cambridge: Cambridge University Press; 2002.
 11. Alberts B, Johnson A, Lewis J, Morgan D, Raff M, Roberts K, *et al*. *Molecular biology of the cell*. 6th ed. New York: Garland Science; 2014.
 12. Guidetti R, Jönsson KI. Long-term anhydrobiotic survival in semi-terrestrial micrometazoans. *Journal of Zoology*. 2002;257(2):181-187.
 13. Horikawa DD, Sakashita T, Katagiri C, Watanabe M, Kikawada T, Nakahara Y, *et al*. Radiation tolerance in the tardigrade *Milnesium tardigradum*. *International Journal of Radiation Biology*. 2006;82(12):843-848.
 14. Tanaka S, Tanaka J, Miwa Y, Horikawa DD, Katayama T, Arakawa K, *et al*. Novel mitochondria-targeted heat-soluble proteins identified in the anhydrobiotic tardigrade improve osmotic tolerance of human cells. *PLoS One*. 2015;10(2):e0118272.