



International Journal of Veterinary Sciences and Animal Husbandry



ISSN: 2456-2912

NAAS Rating (2026): 4.61

VET 2026;11(2): 09-13

© 2026 VET

www.veterinarypaper.com

Received: 05-12-2025

Accepted: 25-01-2026

Pushpa Gautam

Research Scholar, Division of
Extension Education, ICAR-
Indian Veterinary Research
Institute, Izatnagar, Bareilly,
Uttar Pradesh, India

RS Suman

Principal Scientist, Division of
Extension Education, ICAR-
Indian Veterinary Research
Institute, Izatnagar, Bareilly,
Uttar Pradesh, India

Rashmi

Associate Professor, Department
of Dairy Business Management,
College of Dairy Science,
DUVASU, Mathura, Uttar
Pradesh, India

MP Gautam

Subject Matter Specialist
(Entomology/Nematology),
Krishi Vigyan Kendra, Ledaura,
Azamgarh, Uttar Pradesh, India

Bhawna

Research Scholar, Department of
Livestock Production and
Technology (LPT), Sardar
Vallabhbhai Patel University of
Agriculture and Technology
(SVPUAT), Modipuram, Meerut,
Uttar Pradesh, India

Corresponding Author:

Bhawna

Research Scholar, Department of
Livestock Production and
Technology (LPT), Sardar
Vallabhbhai Patel University of
Agriculture and Technology
(SVPUAT), Modipuram, Meerut,
Uttar Pradesh, India

Role of mineral mixture in growth and development of milking animal: A review

Pushpa Gautam, RS Suman, Rashmi, MP Gautam and Bhawna

DOI: <https://www.doi.org/10.22271/veterinary.2026.v11.i2a.3011>

Abstract

Minerals play important role inside the law and protection of various mobile activities of types: Macro minerals (Required in large amount) and Micro minerals (Required in small quantity). It plays very vital position in immunity, manufacturing and replica of animals. Deficiencies of sure mineral elements may additionally purpose reproductive disorders as minerals play a vital position in fitness and reproduction of the cattle. Infertility and reproductive disorders has emerged as essential problem in Indian farm animals farming. Adequate supplementation of minerals depends on age, stage of being pregnant and lactation. For the formation of structural aspect of frame and right functioning of enzymes, hormones, nutrients and cells, macro and micro minerals are similarly essential regulation and protection of various cellular activities within the animal's body.

Keywords: Mineral mixture, growth, development and milking animal

Introduction

Dairy enterprise performs a good sized position, in terms of their contribution to human vitamins (Milk and butter fats), plant vitamins (Farmyard manure) and strength (bullock electricity). Dairy farming is a reliable supply of livelihood for the rural terrible since it has the ability to provide gainful self-employment and sustainable earnings. In addition, it could be maintained with minimal capital investment and labour. Feeding of dairy cattle has a right away effect at the replica and production of the animals. Hence, addition of mineral mixture becomes crucial to achieve the overall potential of milk manufacturing and productiveness of the animal.

Dairy livestock require at the least 17 minerals and three nutrients in their food regimen for best milk production, reproductive performance, and herd fitness. Although classical mineral or diet deficiency signs are uncommon, in many cases beneath- and overfeeding of positive minerals and vitamins does occur. Even small imbalances or deficiencies can turn into reproductive, fitness, and milk production troubles.

Benefits of feeding mineral mixture

- Helps in improving growth rate in calves.
- Better utilization of absorbed nutrients.
- Increases milk production in animals.
- Improves reproductive efficiency and reduces inter- calving period.
- Increases productive life of animals.
- Improves immunity status.

Prevents metabolic diseases like milk fever, ketosis, haematuria which occur around calving period.

Feeding recommendation

Calves 20-25 g daily

Heifers and dry animals 50 g per animal daily

Milking animals: 100-200 g per animal daily (as per milk production)


The importance of the mineral elements

Milk production	Ca, P
Vitamin synthesis	Co, S
Immune System	Cu, Zn, Mn, Cr, Fe, Se
RBC synthesis	Fe, Cu
Energy Metabolism	Mg, P, Mn
Co-factor of Enzymes	Zn, Cu, Mn, K, Mg, Ca, Mo
Hormone system	Mn, Zn, Mg, Fe, K
Skeletal system	Ca, Mg, Zn, Mn, P

Calcium and phosphorous

The most fulfilling degrees of dietary standardized total tract digestible calcium for retention of calcium and phosphorus

have been suggested at or above zero 60% and 0.49% respectively (Lagos L.V. 2019) [15]. The bone formation requirements that have a near reference to the fitness and welfare of the animals determine the need for Ca and P. For a right purposeful musculoskeletal device, boom and performance developments must be combined with adequate bone formation Tanck 2001 [36]. Thus, the surpluses or deficiency of Ca and P have first-rate consequences on bone mineral content material and density, bone development and bone composition. The Ca and P dependent processes depend on the uptake, excretion and availability of those factors. Therefore, vast interest has been paid to the regulation of plasma parathyroid hormone, calcium-binding protein, calcitriol/1, 25(OH)₂D₃ [Sommerville, 1985 and Eklou-Kalonji, 1999] and insulin-like boom factor-1 awareness Sørensen, 2018 [7, 34].



Source: Sampath *et al.*, (2023) [30].

Fig 1: Health benefits and beneficial application of micro and macro-minerals in non-ruminant

Simple or conditioned deficiencies of mineral factors therefore have profound consequences on metabolism and tissue shape. To check the dietary consumption and adequacy of minerals, information wishes to be gathered on mineral element content material of ingredients, diets and water (Simsek and Aykut, 2007) [33]. Plants use these minerals as structural additives in carbohydrates and proteins; natural molecules in metabolism, which includes magnesium in chlorophyll and phosphorus in ATP; enzyme activators like potassium, and for retaining osmotic balance. Calcium is especially implicated in the upkeep of firmness of culmination (Olaiya, 2006) [24].


Zinc

Zinc is a crucial element in bone metabolism which is needed to supply the matrix of collagen protein threads upon which the bone-forming calcium-phosphorus compound is

deposited. It acts as a co-elementa essential function in bone fitness. Zinc additionally stimulates the production of enzymes answerable for the degradation and recycling of tired bits of bone protein. Zn can also effectphysiological motion of vitamin D on calcium metabolism and intervene with the anabolic activity of nutrition D, 2015 and, Zheng, 2014]. Zn can also affect the ordinary acts as a co-factor in over 2 hundred enzymatic reactions and plays vitamin D on bone tissue [Mahdavi-Roshan] [20].

Magnesium

Magnesium is an essential cofactor in eighty% of all cell enzymes and is a crucial contributor to bone fitness. About 60% of the body magnesium is discovered in bone and it makes up approximately 1% of the entire bone mineral content [Dacey, 2001]. It is necessary for the conversion of diet D into its lively shape (Uwitonze, 2018) [3].



Source: Sampath *et al.*, (2023) [30].

Fig 2: Schematic view on the mode of action of the micro-minerals in non-ruminant diets.

Copper

Copper is involved in lots of organic procedures along with enzymatic reactions and bone fitness. The formation of bone, mineralization of skeleton, and the integrity of the connective tissue are substantially encouraged by means of copper (Aaseth, 2012 and Rondanelli, 2012) [1, 29].

Silicon and boron

Several studies tested that silicon is also an vital trace mineral for regular bone development, cartilage boom, and stepped forward bone great (Pietak, 2007 and Incharoen, 2016) [10, 26]. Some evidence indicates the involvement of silicon with calcium metabolism and the formation and stabilization of the extracellular bone matrix (Reffitt, 2003 and, Jugdaoh singh, 2007) [28].

Boron

Boron is more and more recognized as an element that has an influence on bone health via stabilizing and extending the half of-life of diet D and estrogen (Volpe, 1993 Penland 1992 and Institute of Medicine, 2001) [11, 25]. The supplementations of boron within the form of boric acid have been shown to growth bone electricity in rats and laying hens [Chapin, 1998 and Wilson, 1998] [2].

Calcium

Animals are provided minerals through food, mineral aggregate and mineral blocks. Ca is liable for development of bones and its function. Iron containing enzymes are worked as immune gadget in frame. In male animal's reproductive standard performance are stricken by Cu deficiency (Dey, *et al.*, 2018) [5].

Vitamins

Supplemental vitamin D is wanted is required for normal calcification of bones (Driver, 2005) [6]. The secretion of parathyroid hormone is regulated with the aid of vitamin D

thereby stimulating numerous tissues with vitamin D receptors (Norman, 1985) [23]. Regardless of beginning, vitamin D is an inactive pro hormone and need to first be metabolized to its hormonal shape before it could feature. Thus, to generate the biologically lively form, nutrition Dcholecalciferol is hydroxylated to twenty-five-hydroxycholecalciferol (25[OH]₂D) by way of the enzyme 25-hydroxylase; and ii) in the kidney, 25-hydroxycholecalciferol is converted to at least one, 25(OH)₂D by means of the enzyme 1 α hydroxylase (Holick, 2011 and Razzaque, 2011) [9, 27]. Supplemental vitamin D is required for the green absorption vitamin D needs to be processed similarly. This activation system takes place in 2 steps: i) in the liver, efficient absorption and metabolism of calcium and phosphorus consequently.

Essential vitamins have to be provided in suitable quantities and in the paperwork which might be biologically utilizable for the efficient manufacturing of livestock and chicken and the renovation in their regular life features (Xu, 2019). The ratio of vitamin and minerals ought to be maintained. The animals must provide ok amount of minerals in all ranges of life functions (Ma YL, 2018 Li, 2018) [18, 19].

Technologies for the improvement of supplemental micronutrient bioavailability

Nano-engineering of substances enhances their bioavailability by shielding them towards chemical situations in the GI tract, and permitting the controlled release in the GI tract or via an improved transfer thru the intestinal wall (Jampliek, 2019, Sharma, 2019) [12, 32]. The traits of nano-engineered substances affecting the final nutritional value encompass particle size, physical nation of nano- substances and their floor homes (Gonnet, 2010). Our examine additionally validated that lipid-matrix lined ZnO at a low dose ought to replacement for the conventional excessive dose ZnO in weaning pigs (Upadhaya, 2018) [8, 37].

Recommendations for mineral supplementation

Minerals are additives of the food regimen that should be administered in a managed, periodic, and cautious way. Mineral requirements may also vary by animal species, as well as breed, weight loss plan, area, and type of production.

- Use salt resources nicely designed for each animal species. Industrial or cooking salt must now not be used.
- The mineral supplements have to be palatable for the animals and be properly balanced according to the necessities.
- The presentation of the minerals must be in keeping with the manufacturer's hints. Wet, compacted or infected product ought to be avoided.
- Mineral supplementation can be done according to soil and weight loss program analysis to modify to the wishes, as well as to establish an inexpensive fee. This will keep away from extra minerals in the food plan that can be poisonous.
- In animal vitamins, the management of minerals should be done carefully in step with the degree of production and traits of the flock alter in constantly.
- The balance of minerals is essential because the excess of a number of them reasons the inability to absorb others or block their capability.

Conclusion

Macro minerals are the ones which can be found in better proportions at the body stage and include calcium, phosphorus, potassium, magnesium, sulfur, and sodium/chlorine. Micro minerals, on the other hand, are in smaller proportions, but additionally have fundamental features in animals. In view of the importance of those macro and micro mineral factors within the nutrients of human beings, animals and vegetation and their metabolic inter-relationships which impacts other critical factors wanted for the survival of residing organisms like enzymes, anti-oxidants, nutrients and many others, it is critical to frequently obtain updated statistics on the minerals content of water and the variously normally fed on plant ingredients used for human and animal foods and feeds respectively e.g. Legumes, cereals, culmination and veggies. This is due to the consequences of chemicals like pesticides, herbicides, fertilizers at the mineral contents of the soils wherein these plant lives are cultivated. Genetic, vicinity and environmental factors may also impact the ranges of the mineral elements in flora. Further research is required to find out the minimal dietary stage of mineral necessities, most tolerable levels, and the interrelationship among the minerals. There is also the need to evaluate the results of various processing techniques at the minerals content of water and plant meals.

Recommendation

With the advancements inside the subject of analytical sciences, regarding the usage of greater state-of-the-art gadget for the quantitative and qualitative evaluation of ingredients/feeds. There is need to revisit the research/investigations /exams /critiques, that allows you to re-validate the statistics at the mineral elements composition of human and animal diets, in particular in the growing nations. The earlier data on these mineral factors were primarily based on analysis using much less sensitive methods which won't be dependable. Further research is required to find out the interactions among the mineral factors, for instance, the interplay between silicon, aluminium and copper. There is likewise the want to assess the effects of

different processing strategies on the minerals content of water and plant meals. This will go a long way in decreasing the problem of malnutrition resulting from inadequate food deliver, consumption of unbalanced diets and inadequate nutrition education.

References

1. Aaseth J, Boivin G, Andersen O. Osteoporosis and trace elements - an overview. *J Trace Elem Med Biol.* 2012;26:149-152. doi:10.1016/j.jtemb.2012.03.017.
2. Chapin RE, Ku WW, Kenney MA, McCoy H. The effects of dietary boric acid on bone strength in rats. *Biol Trace Elem Res.* 1998;66:395-399. doi:10.1007/BF02783150.
3. Dacey MJ. Hypomagnesemic disorders. *Crit Care Clin.* 2001;17:155-173. doi:10.1016/S0749-0704(05)70157-3.
4. Devi S, Upadhaya L, Kim IH. Importance of micronutrients in bone health of monogastric animals and techniques to improve the bioavailability of micronutrient supplements: a review. *Asian-Australas J Anim Sci.* 2020;33(12):1885-1895.
5. Dey D, Sharma B, Khare A, Gupta SK. Importance of mineral feeding in dairy animals: a review. *Bhartiya Krishi Anusandhan Patrika.* 2018;33(4):279-282.
6. Driver JP, Pesti GM, Bakalli RI, Edwards HM Jr. Calcium requirements of the modern broiler chicken as influenced by dietary protein and age. *Poult Sci.* 2005;84:1629-1639. doi:10.1093/ps/84.10.1629.
7. Eklou-Kalonji E, Zerath E, Colin C. Calcium-regulating hormones, bone mineral content, breaking load and trabecular remodelling are altered in growing pigs fed calcium-deficient diets. *J Nutr.* 1999;129:188-193. doi:10.1093/jn/129.1.188.
8. Gonnet M, Lethuaut L, Boury F. New trends in encapsulation of lip soluble vitamins. *J Control Release.* 2010;146:276-290. doi:10.1016/j.jconrel.2010.01.037.
9. Holick MF. Vitamin D: evolutionary, physiological and health perspectives. *Curr Drug Targets.* 2011;12:4-18. doi:10.2174/138945011793591635.
10. Incharoen T, Tartrakoon W, Nakhon S, Treetan S. Effects of dietary silicon derived from rice hull ash on meat quality and bone breaking strength of broiler chickens. *Asian J Anim Vet Adv.* 2016;11:417-422. doi:10.3923/ajava.2016.417.422.
11. Institute of Medicine. Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium and zinc. Washington (DC): National Academies Press; 2001. doi:10.17226/10026.
12. Jampliek J, Kos J, Kralova K. Potential of nanomaterial applications in dietary supplements and foods for special medical purposes. *Nanomaterials.* 2019;9:296. doi:10.3390/nano9020296.
13. Jugdaohsingh R. Silicon and bone health. *J Nutr Health Aging.* 2007;11:99-110.
14. Khoushabi F, Shadan MR, Miri A, Sharifi-Rad J. Determination of maternal serum zinc, iron, calcium and magnesium during pregnancy and association with outcome. *Mater Sociomed.* 2016;28:104-107.
15. Lagos LV, Lee SA, Fondevila G. Influence of dietary digestible calcium on growth performance and bone mineralization in pigs. *J Anim Sci Biotechnol.* 2019;10:47. doi:10.1186/s40104-019-0349-2.
16. Larson CK. Role of trace minerals in animal production. *Proc Nutr Conf Univ Tennessee.* 2005.

17. Lebret B, Candek-Potokar M. Review: pork quality attributes from farm to fork. Part II. Processed pork products. *Animal*. 2022;16:100383.
18. Li KX, Wang JS, Yuan D, Zhao RX, Wang YX, Zhan XA. Effects of different selenium sources and levels on antioxidant status in broiler breeders. *Asian-Australas J Anim Sci*. 2018;31:1939-1945. doi:10.5713/ajas.18.0226.
19. Ma YL, Lindemann MD, Webb SF, Rentfrow G. Evaluation of trace mineral source and preharvest deletion on tissue mineral status in pigs. *Asian-Australas J Anim Sci*. 2018;31:252-262. doi:10.5713/ajas.17.0189.
20. Mahdavi-Roshan M, Ebrahimi M, Ebrahimi A. Copper, magnesium, zinc and calcium status in osteopenic and osteoporotic women. *Clin Cases Miner Bone Metab*. 2015;12:18-21. doi:10.11138/ccmbm/2015.12.1.018.
21. McDowell LR. Vitamins in animal and human nutrition. 2nd ed. Ames (IA): Iowa State Press; 2000.
22. McDowell LR. Minerals in animal and human nutrition. 2nd ed. New York: Elsevier; 2003.
23. Norman AW. The vitamin D endocrine system. *Physiologist*. 1985;28:219-232.
24. Olaiya CO. Effects of three plant bioregulators on some biochemical properties of *Lycopersicon esculentum* (L.) Mill. Ibadan (Nigeria): University of Ibadan; 2006. PhD thesis.
25. Penland JG. Dietary boron, brain function and cognitive performance. *Environ Health Perspect*. 1994;102 Suppl 7:S65-S72. doi:10.1289/ehp.94102s765.
26. Pietak AM, Reid JW, Stott MJ, Sayer M. Silicon substitution in calcium phosphate bioceramics. *Biomaterials*. 2007;28:4023-4032. doi:10.1016/j.biomaterials.2007.05.003.
27. Razzaque MS. The dualistic role of vitamin D in vascular calcifications. *Kidney Int*. 2011;79:708-714. doi:10.1038/ki.2010.432.
28. Reffitt DM, Ogston N, Jugdaohsingh R. Orthosilicic acid stimulates collagen type I synthesis and osteoblastic differentiation. *Bone*. 2003;32:127-135. doi:10.1016/S8756-3282(02)00950-X.
29. Rondanelli M, Opizzi A, Perna S, Faliva MA. Update on nutrients involved in maintaining healthy bone. *Endocrinol Nutr*. 2013;60:197-210. doi:10.1016/j.endonu.2012.09.006.
30. Sampath V, Sureshkumar S, Seok WJ, Kim IH. Role and functions of micro and macro-minerals in swine nutrition: a short review. *J Anim Sci Technol*. 2023;65(3):479-489.
31. Shannon MC, Hill GM. Trace mineral supplementation for intestinal health of young monogastric animals. *Front Vet Sci*. 2019;6:73.
32. Sharma S, Jaiswal S, Duffy B, Jaiswal AK. Nanostructured materials for food applications. *Bioengineering*. 2019;6:26. doi:10.3390/bioengineering6010026.
33. Simsek A, Aykut O. Evaluation of the microelement profile of Turkish hazelnut varieties. *Int J Food Sci Nutr*. 2007;58:677-688.
34. Sommerville BA, Maunder E, Ross R, Care AD, Brown RC. Effect of dietary calcium and phosphorus depletion on vitamin D metabolism in pigs. *Horm Metab Res*. 1985;17:78-81. doi:10.1055/s-2007-1013456.
35. Sorensen KU, Tauson AH, Poulsen HD. Long term differentiated phosphorus supply affects nutrient balance and bone growth in pigs. *Livest Sci*. 2018;211:14-20. doi:10.1016/j.livsci.2018.03.002.
36. Tanck E, Homminga J, Van Lenthe GH, Huiskes R. Increase in bone volume fraction precedes architectural adaptation in growing bone. *Bone*. 2001;28:650-654.
37. Upadhaya SD, Kim YM, Lee KY, Kim IH. Use of protected zinc oxide in lower doses in weaned pigs. *Anim Feed Sci Technol*. 2018;240:1-10. doi:10.1016/j.anifeedsci.2018.03.012.
38. Uwitonze AM, Razzaque MS. Role of magnesium in vitamin D activation and function. *J Am Osteopath Assoc*. 2018;118:181-189. doi:10.7556/jaoa.2018.037.
39. Volpe S, Taper LJ, Meacham S. Relationship between boron, magnesium status and bone mineral density. *Magnes Res*. 1993;6:291-296.
40. Wilson JH, Ruszler PL. Long term effects of boron on layer bone strength and production parameters. *Br Poult Sci*. 1998;39:11-15. doi:10.1080/00071669889312.
41. Xu H, Dai S, Zhang K. Dietary phosphorus deficiency impaired growth and intestinal function of meat ducks. *Asian-Australas J Anim Sci*. 2019;32:897-906. doi:10.5713/ajas.18.0683.
42. Zheng J, Mao X, Ling J, He Q, Quan J. Low serum levels of zinc, copper and iron as risk factors for osteoporosis: a meta-analysis. *Biol Trace Elem Res*. 2014;160:15-23. doi:10.1007/s12011-014-0031-7.

How to Cite This Article

Gautam P, Suman RS, Rashmi, Gautam MP, Bhawna. Role of mineral mixture in growth and development of milking animal: A review. *International Journal of Veterinary Sciences and Animal Husbandry*. 2026;11(2):09-13.

Creative Commons (CC) License

This is an open-access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.