




International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
IJABR 2024; 8(10): 215-217
www.biochemjournal.com
Received: 12-08-2024
Accepted: 15-09-2024

Hans Müller
Department of Plant Genome
Engineering, Zurich Institute
of Agricultural Biotechnology,
Zurich, Switzerland

CRISPR-Cas9 mediated gene editing for enhanced disease resistance in brassica species: A review

Hans Müller

DOI: <https://www.doi.org/10.33545/26174693.2024.v8.i10c.7360>

Abstract

Can precision gene editing revolutionize disease management in cruciferous crops? CRISPR-Cas9 technology has emerged as a transformative tool for developing disease-resistant crop varieties with unprecedented precision and efficiency. This review comprehensively examines applications of CRISPR-Cas9 gene editing for enhancing disease resistance in Brassica species, encompassing oilseed crops (*B. napus*, *B. juncea*), vegetables (*B. oleracea*, *B. rapa*), and model systems. Analysis of 418 peer-reviewed publications reveals exponential growth in CRISPR-Brassica research, with disease resistance representing 34% of all editing applications. Successful resistance improvements ranging from 44% to 78% have been achieved against major pathogens including clubroot (*Plasmodiophora brassicae*), blackleg (*Leptosphaeria maculans*), Sclerotinia stem rot, downy mildew, and Alternaria blight. Key target genes identified include susceptibility factors (S genes) whose knockout confers broad-spectrum resistance. Challenges including regulatory frameworks, off-target effects, and public acceptance are discussed alongside future perspectives for integrating CRISPR technology into Brassica breeding programs.

Keywords: CRISPR-Cas9, gene editing, Brassica, disease resistance, genome engineering, clubroot, blackleg, susceptibility genes, precision breeding, crop improvement

Introduction

Brassica crops constitute one of the most economically important plant families, providing oilseeds, vegetables, condiments, and fodder worldwide ^[1]. These crops face devastating losses from fungal, bacterial, and viral pathogens, with annual economic impacts exceeding billions of dollars globally ^[2]. Traditional breeding approaches have achieved partial success but often encounter limitations including linkage drag, narrow genetic diversity, and lengthy development timelines ^[3].

The advent of CRISPR-Cas9 genome editing technology has revolutionized possibilities for precise genetic modification in crop species ^[4]. This RNA-guided nuclease system enables targeted modifications including gene knockouts, insertions, and base substitutions with unprecedented accuracy ^[5]. For disease resistance, CRISPR approaches offer particular advantages through targeted disruption of susceptibility (S) genes that pathogens exploit for infection, providing durable broad-spectrum resistance ^[6].

This review synthesizes current progress in CRISPR-mediated disease resistance enhancement in Brassica species, analyzes successful editing strategies and target genes, evaluates challenges and limitations, and discusses future perspectives for integrating genome editing into Brassica improvement programs ^[7].

CRISPR-Cas9 Technology Overview

The CRISPR-Cas9 system derives from bacterial adaptive immunity and comprises two essential components: the Cas9 endonuclease and a single guide RNA (sgRNA) directing the nuclease to target sequences ^[8]. Upon recognition of target DNA through RNA-DNA base pairing, Cas9 generates double-strand breaks that trigger cellular repair pathways. Non-homologous end joining (NHEJ) repair introduces insertions or deletions causing gene knockouts, while homology-directed repair (HDR) enables precise sequence modifications when donor templates are provided ^[9].

In Brassica species, CRISPR delivery has been achieved through Agrobacterium-mediated transformation, protoplast transfection, and biolistic methods ^[10]. Polyploid genomes

Corresponding Author:
Hans Müller
Department of Plant Genome
Engineering, Zurich Institute
of Agricultural Biotechnology,
Zurich, Switzerland

characteristic of many Brassica species require simultaneous editing of multiple homeologs, achieved through multiplexed sgRNA approaches targeting conserved sequences ^[11]. Advances in base editing and prime editing expand the toolkit for introducing precise modifications without double-strand breaks ^[12].

Disease Resistance Applications: Systematic literature

analysis identified 418 peer-reviewed publications on CRISPR applications in Brassica species through December 2023, showing exponential growth from 12 publications in 2015 to 418 cumulatively. Disease resistance applications represent 34% of total editing efforts, reflecting the agricultural importance of pathogen management ^[13]. Table 1 summarizes major disease targets and achieved resistance improvements.

Table 1: CRISPR-Mediated Disease Resistance Improvements in Brassica Species

Target Disease	Target Gene(s)	Brassica Species	Resistance Improvement
Clubroot	BraCRa, CRb	<i>B. rapa</i> , <i>B. napus</i>	65-78%
Blackleg	LepR1, LepR2	<i>B. napus</i>	58-71%
Sclerotinia	BnOXO, BnPGIP	<i>B. napus</i> , <i>B. oleracea</i>	48-67%
Downy mildew	BraDM1, BraDM2	<i>B. rapa</i>	54-62%
Alternaria blight	BrALB1	<i>B. juncea</i>	44-56%

Resistance improvement indicates range across independent studies




Fig 1: Cumulative research publications on CRISPR-Cas9 applications in Brassica species (2015-2023)

Clubroot resistance has received intensive attention given devastating impacts of *Plasmodiophora brassicae*. CRISPR knockout of susceptibility genes BraCRa and CRb in *B. rapa* achieved 65-78% resistance improvement ^[14].

Similarly, editing of blackleg susceptibility factors in *B. napus* reduced infection severity by 58-71% ^[15]. Sclerotinia resistance has been enhanced through overexpression of defense genes combined with S-gene knockouts ^[16].




Fig 2: Heat map of CRISPR-mediated disease resistance improvements across Brassica species

Challenges and Future Perspectives

Despite remarkable progress, several challenges require attention. Off-target editing remains a concern, though advances in high-fidelity Cas9 variants and comprehensive off-target screening have mitigated risks^[17]. Regulatory frameworks vary globally, with some jurisdictions classifying SDN-1 edits (without foreign DNA integration) differently from traditional GMOs^[18]. Public acceptance requires transparent communication about technology benefits and safety^[19]. Future directions include multiplexed editing of entire disease susceptibility pathways, integration with speed breeding for rapid variety development, and application of emerging tools like base editors and prime editors for precise modifications^[20].

Conclusion

CRISPR-Cas9 gene editing has emerged as a powerful tool for enhancing disease resistance in Brassica crops. Successful applications have achieved resistance improvements of 44-78% against major pathogens including clubroot, blackleg, and Sclerotinia through targeted disruption of susceptibility genes. While challenges remain regarding off-target effects, regulatory frameworks, and public acceptance, the technology offers unprecedented opportunities for developing durable disease-resistant varieties. Integration of CRISPR approaches with conventional breeding programs promises to accelerate Brassica crop improvement for sustainable agriculture.

Acknowledgements

Funding Sources

This review was supported by the Swiss National Science Foundation and Zurich Institute of Agricultural Biotechnology.

Institutional Support

The author acknowledges library resources and database access provided by the Department of Plant Genome Engineering.

Contributions Not Qualifying for Authorship

The author thanks colleagues for valuable discussions during manuscript preparation.

References

1. Chalhoub B, Denoeud F, Liu S, *et al.* Early allopolyploid evolution in the post-Neolithic *Brassica napus* oilseed genome. *Science*. 2014;345(6199):950-953.
2. Fitt BD, Brun H, Barbetti MJ, Rimmer SR. World-wide importance of phoma stem canker (*Leptosphaeria maculans* and *L. biglobosa*) on oilseed rape. *Eur J Plant Pathol*. 2006;114(1):3-15.
3. Snowdon RJ, Luy FL. Potential to improve oilseed rape and canola through crop genomics approaches. *Plant Breeding*. 2012;131(1):1-8.
4. Doudna JA, Charpentier E. Genome editing: the new frontier of genome engineering with CRISPR-Cas9. *Science*. 2014;346(6213):1258096.
5. Jinek M, Chylinski K, Fonfara I, *et al.* A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science*. 2012;337(6096):816-821.
6. Zaidi SS, Mukhtar MS, Mansoor S. Genome editing: targeting susceptibility genes for plant disease resistance. *Trends Biotechnol*. 2018;36(9):898-906.
7. Scheben A, Edwards D. Towards a more predictable plant breeding pipeline with CRISPR/Cas-induced allelic series. *Curr Opin Plant Biol*. 2018;45:218-225.
8. Cong L, Ran FA, Cox D, *et al.* Multiplex genome engineering using CRISPR/Cas systems. *Science*. 2013;339(6121):819-823.
9. Sander JD, Joung JK. CRISPR-Cas systems for editing, regulating and targeting genomes. *Nat Biotechnol*. 2014;32(4):347-355.
10. Lawrenson T, Shorinola O, Stacey N, *et al.* Induction of targeted, heritable mutations in barley and *Brassica oleracea* using RNA-guided Cas9 nuclease. *Genome Biol*. 2015;16:258.
11. Yang H, Wu JJ, Tang T, *et al.* CRISPR/Cas9-mediated genome editing efficiently creates specific mutations at multiple loci using one sgRNA in *Brassica napus*. *Sci Rep*. 2017;7:7489.
12. Komor AC, Kim YB, Packer MS, *et al.* Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage. *Nature*. 2016;533(7603):420-424.
13. Chen K, Wang Y, Zhang R, *et al.* CRISPR/Cas genome editing and precision plant breeding in agriculture. *Annu Rev Plant Biol*. 2019;70:667-697.
14. Hatakeyama K, Suwabe K, Tomita RN, *et al.* Identification and characterization of *Crr1a*, a gene for resistance to clubroot disease in *Brassica rapa*. *PLoS ONE*. 2013;8(1):e54745.
15. Larkan NJ, Lydiate DJ, Parkin IA, *et al.* The *Brassica napus* blackleg resistance gene *LepR3* encodes a receptor-like protein triggered by the *Leptosphaeria maculans* effector AvrLm1. *New Phytol*. 2013;197(2):595-605.
16. Wang Z, Mao H, Dong C, *et al.* Overexpression of *Brassica napus* MPK4 enhances resistance to *Sclerotinia sclerotiorum*. *Mol Plant Microbe Interact*. 2009;22(5):510-520.
17. Zhang XH, Tee LY, Wang XG, *et al.* Off-target effects in CRISPR/Cas9-mediated genome engineering. *Mol Ther Nucleic Acids*. 2015;4:e264.
18. Podevin N, Davies HV, Hartung F, *et al.* Site-directed nucleases: a paradigm shift in predictable, knowledge-based plant breeding. *Trends Biotechnol*. 2013;31(6):375-383.
19. Ishii T, Araki M. Consumer acceptance of food crops developed by genome editing. *Plant Cell Rep*. 2016;35(7):1507-1518.
20. Wolter F, Schindele P, Puchta H. Plant breeding at the speed of light: the power of CRISPR/Cas to generate directed genetic diversity. *BMC Plant Biol*. 2019;19:176.