



International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
IJABR 2024; 8(12): 133-136
www.biochemjournal.com
Received: 08-10-2024
Accepted: 12-11-2024

David Harrison
Department of Molecular
Biotechnology, Manchester
Institute of Applied Sciences,
Manchester, United Kingdom

Emma Richardson
Department of Molecular
Biotechnology, Manchester
Institute of Applied Sciences,
Manchester, United Kingdom

CRISPR-based editing in animal cell lines and protocol optimization

David Harrison and Emma Richardson

DOI: <https://www.doi.org/10.33545/26174693.2024.v8.i12b.7354>

Abstract

CRISPR-Cas9 technology represents a programmable molecular scissors system enabling precise genome editing through RNA-guided DNA cleavage. This research systematically optimized CRISPR-Cas9 protocols for efficient gene editing in commonly used animal cell lines including HEK293, HeLa, CHO, and NIH3T3. The investigation was conducted at Manchester Institute of Applied Sciences between April 2022 and January 2024. Multiple parameters were evaluated including guide RNA design algorithms, ribonucleoprotein complex formation conditions, transfection methods, cell confluence at treatment, and post-transfection recovery protocols. Editing efficiency was quantified using T7 endonuclease I assays and next-generation sequencing of target loci. Optimization experiments identified several critical parameters substantially affecting editing outcomes. Ribonucleoprotein complex pre-formation at 37 °C for 15 minutes improved efficiency by 34% compared to room temperature assembly. Transfection at 70-80% cell confluence yielded 28% higher editing rates than treatment at lower densities. Nucleofection achieved the highest mean efficiency at 74.3%, followed by optimized electroporation at 68.4% and optimized lipofection at 52.1%. A combined optimized protocol incorporating these refinements achieved 82.7% editing efficiency in HEK293 cells with minimal cytotoxicity. Cell type-specific variations were observed, with CHO cells requiring modified electrical parameters for optimal results. These findings provide an evidence-based framework for maximizing CRISPR-Cas9 editing efficiency across diverse animal cell lines.

Keywords: CRISPR-Cas9, gene editing, protocol optimization, transfection, ribonucleoprotein, cell lines, nucleofection, electroporation, genome engineering, biotechnology

Introduction

CRISPR-Cas9 is defined as a bacterial adaptive immune system repurposed for programmable genome editing, wherein the Cas9 endonuclease is directed to specific genomic sequences by complementary guide RNAs to create targeted double-strand breaks [1, 1]. Since its initial application for mammalian genome editing in 2013, this technology has transformed biological research by enabling precise genetic modifications previously achievable only through laborious homologous recombination approaches.

The efficiency of CRISPR-Cas9 editing depends on multiple interconnected factors that must be optimized for each experimental system [2]. Guide RNA design influences both on-target cleavage efficiency and off-target activity, with various algorithms offering different predictive accuracies [2]. Delivery of editing components to target cells presents additional challenges, with lipofection, electroporation, and nucleofection each offering distinct advantages depending on cell type and experimental requirements [3].

Ribonucleoprotein delivery has emerged as a preferred approach for many applications, offering reduced off-target effects compared to plasmid-based expression due to transient Cas9 presence [4, 3]. However, optimal conditions for RNP complex formation and cellular delivery vary considerably across cell types and have not been systematically characterized for commonly used animal cell lines.

Post-transfection cell handling represents another underexplored area where protocol refinements may substantially impact editing outcomes [4]. Parameters including recovery medium composition, timing of analysis, and cell density during editing likely influence results but are inconsistently reported and rarely optimized [5]. Standardized protocols accounting for these variables would benefit the research community.

Corresponding Author:
David Harrison
Department of Molecular
Biotechnology, Manchester
Institute of Applied Sciences,
Manchester, United Kingdom

This research aimed to systematically optimize CRISPR-Cas9 editing protocols across multiple commonly used animal cell lines through evaluation of guide RNA design, RNP formation conditions, transfection methods, and recovery protocols [5]. We hypothesized that systematic optimization would substantially improve editing efficiency compared to standard manufacturer recommendations.

Material and Methods

Materials

This research utilized four commonly used animal cell lines: HEK293 (human embryonic kidney), HeLa (human cervical carcinoma), CHO-K1 (Chinese hamster ovary), and NIH3T3 (mouse embryonic fibroblast) [6]. All lines were obtained from the European Collection of Authenticated Cell Cultures with authentication confirmed by STR profiling. The research was conducted at Manchester Institute of Applied Sciences between April 2022 and January 2024 [6]. Streptococcus pyogenes Cas9 protein was obtained from commercial sources (Integrated DNA Technologies) [7]. Synthetic guide RNAs targeting the AAVS1 safe harbor locus (human cells) or Rosa26 locus (mouse cells) were synthesized with chemical modifications for enhanced stability. Transfection reagents included Lipofectamine CRISPRMAX, Neon Transfection System consumables, and Lonza Nucleofector kits [7].

Methods

Guide RNAs were designed using multiple algorithms including CRISPOR, Benchling, and CHOPCHOP, with sequences selected based on predicted on-target efficiency and minimal off-target potential [8]. RNP complexes were formed by incubating Cas9 protein with guide RNA at molar ratios ranging from 1:1.2 to 1:3, with temperatures and durations systematically varied [8]. Transfection optimization compared lipofection, electroporation, and nucleofection across cell confluence ranges of 50-90% [9]. Post-transfection recovery conditions

evaluated medium composition, supplementation with recovery enhancers, and timing before passage. Editing efficiency was quantified at 72 hours post-transfection using T7 endonuclease I assays with confirmation by targeted amplicon sequencing [9].

Statistical analyses employed GraphPad Prism version 9. [10]. Comparisons used one-way ANOVA with Tukey post-hoc tests for multiple groups [10]. Optimization experiments utilized factorial designs to identify parameter interactions. Significance required $p < 0.05$.

Results

Initial experiments established baseline editing efficiencies using standard manufacturer protocols, yielding mean values of 28.4% for lipofection, 45.2% for electroporation, and 58.7% for nucleofection in HEK293 cells [11]. Systematic optimization of individual parameters revealed substantial potential for improvement across all delivery methods.

Table 1: Optimization of RNP Complex Formation Conditions

Temperature	Duration (min)	Efficiency (%)	p-value vs RT
Room temp (22 °C)	10	45.2±6.8	Reference
Room temp (22 °C)	15	47.8±7.2	0.634
Room temp (22 °C)	30	46.4±8.1	0.812
37 °C	10	58.4±5.4	0.008
37 °C	15	68.7±4.8	<0.001
37 °C	30	64.2±6.2	<0.001

Values expressed as Mean ± SD (n = 6 independent experiments) [12]. RT: room temperature. Optimal condition highlighted: 37 °C for 15 minutes.

RNP complex formation at 37 °C for 15 minutes produced optimal editing efficiency of 68.7%, representing a 52% improvement over room temperature assembly (Table 1) [13, 12]. This enhancement likely reflects improved Cas9 conformational dynamics and guide RNA loading at physiological temperature [10].




Fig 1: Comparison of CRISPR-Cas9 editing efficiency across different transfection protocols in HEK293 cells. Optimized protocols significantly outperform standard conditions. Box plots show median, quartiles, and range. *** $p < 0.001$; ns: not significant.

Table 2: Cell Type-Specific Editing Efficiency with Optimized Protocol

Cell Line	Standard (%)	Optimized (%)	Improvement
HEK293	58.7±8.4	82.7±5.2	+40.9%
HeLa	42.3±9.6	71.4±7.8	+68.8%
CHO-K1	38.6±11.2	64.8±8.4	+67.9%
NIH3T3	51.2±7.8	76.3±6.2	+49.0%

Values expressed as Mean ± SD (n = 6) ^[14]. Standard: manufacturer protocol with nucleofection; Optimized: combined protocol incorporating all refinements.




Fig 2: Schematic representation of the optimized CRISPR-Cas9 protocol workflow. Key optimization points include RNP formation at 37 °C, transfection at 70-80% confluence, and 24-hour recovery period before analysis.

Comprehensive Interpretation

The combined optimized protocol incorporating 37 °C RNP formation, 70-80% cell confluence, nucleofection delivery, and standardized recovery conditions achieved substantial improvements across all tested cell lines (Table 2) ^[15, 13]. HEK293 cells reached 82.7% efficiency, representing 40.9% improvement over standard protocols. HeLa and CHO cells showed the greatest relative improvements at 68.8% and 67.9% respectively, suggesting these harder-to-transfect lines benefit most from optimization ^[11].

Discussion

This research establishes that systematic optimization of CRISPR-Cas9 protocols can substantially improve editing efficiency across diverse animal cell lines. ^[14]. The combined effect of individually modest improvements produced dramatic overall enhancement, with final efficiencies approaching or exceeding 80% in optimized conditions ^[12]. The benefit of 37 °C RNP complex formation likely reflects temperature-dependent conformational changes in Cas9 that enhance guide RNA loading and subsequent DNA recognition. ^[15]. Previous structural studies demonstrated that Cas9 undergoes substantial rearrangement upon guide

RNA binding, a process that may be kinetically favored at physiological temperature ^[13]. Cell confluence at transfection emerged as a critical parameter, with optimal results at 70-80% density. This observation aligns with the cell cycle dependency of DNA repair pathway choice, as homology-directed repair predominates in S/G2 phases when sister chromatids are available as templates. Cells at optimal confluence likely represent a more homogeneous population regarding cell cycle distribution ^[14].

Limitations

Several limitations warrant acknowledgment. Optimization focused on a single target locus per species, and results may vary for other genomic locations with different chromatin accessibility. The research did not comprehensively evaluate off-target effects of optimized protocols. Cost considerations for nucleofection consumables may limit applicability in resource-constrained settings ^[15].

Practical Applications

The optimized protocols described here are immediately applicable to research laboratories performing CRISPR-Cas9 experiments in common cell lines. The systematic

framework for optimization can guide adaptation to additional cell types not tested in this research. Protocol standardization should improve reproducibility across laboratories and facilitate comparison of results.

Conclusion

This research demonstrates that systematic optimization of CRISPR-Cas9 protocols substantially improves editing efficiency in animal cell lines. Key refinements include RNP complex formation at 37 °C for 15 minutes, transfection at 70-80% cell confluence, and standardized 24-hour recovery periods. The combined optimized protocol achieved 82.7% efficiency in HEK293 cells with improvements of 40-69% across all tested lines. These findings provide an evidence-based framework for maximizing CRISPR-Cas9 editing outcomes in research applications.

References

1. Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E. A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science*. 2012;337(6096):816-821.
2. Doench JG, Fusi N, Sullender M, *et al.* Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. *Nat Biotechnol*. 2016;34(2):184-191.
3. Liang X, Potter J, Kumar S, *et al.* Rapid and highly efficient mammalian cell engineering via Cas9 protein transfection. *J Biotechnol*. 2015;208:44-53.
4. Kim S, Kim D, Cho SW, Kim J, Kim JS. Highly efficient RNA-guided genome editing in human cells via delivery of purified Cas9 ribonucleoproteins. *Genome Res*. 2014;24(6):1012-1019.
5. Liu C, Zhang L, Liu H, Cheng K. Delivery strategies of the CRISPR-Cas9 gene-editing system for therapeutic applications. *J Control Release*. 2017;266:17-26.
6. Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. Genome engineering using the CRISPR-Cas9 system. *Nat Protoc*. 2013;8(11):2281-2308.
7. Zuris JA, Thompson DB, Shu Y, *et al.* Cationic lipid-mediated delivery of proteins enables efficient protein-based genome editing *in vitro* and *in vivo*. *Nat Biotechnol*. 2015;33(1):73-80.
8. Haeussler M, Schönig K, Eckert H, *et al.* Evaluation of off-target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. *Genome Biol*. 2016;17(1):148.
9. Sentmanat MF, Peters ST, Flober CP, Bhaya D, Bhattacharyya S. A survey of validation strategies for CRISPR-Cas9 editing. *Sci Rep*. 2018;8(1):888.
10. Sternberg SH, Redding S, Jinek M, Greene EC, Doudna JA. DNA interrogation by the CRISPR RNA-guided endonuclease Cas9. *Nature*. 2014;507(7490):62-67.
11. Lin S, Staahl BT, Alla RK, Doudna JA. Enhanced homology-directed human genome engineering by controlled timing of CRISPR/Cas9 delivery. *eLife*. 2014;3:e04766.
12. Miyaoka Y, Berman JR, Cooper SB, *et al.* Systematic quantification of HDR and NHEJ reveals effects of locus, nuclease, and cell type on genome-editing. *Sci Rep*. 2016;6:23549.
13. Nishimasu H, Ran FA, Hsu PD, *et al.* Crystal structure of Cas9 in complex with guide RNA and target DNA. *Cell*. 2014;156(5):935-949.
14. Gutschner T, Haemmerle M, Genovese G, Draetta GF, Chin L. Post-translational regulation of Cas9 during G1 enhances homology-directed repair. *Cell Rep*. 2016;14(6):1555-1566.
15. Vakulskas CA, Dever DP, Rettig GR, *et al.* A high-fidelity Cas9 mutant delivered as a ribonucleoprotein complex enables efficient gene editing in human hematopoietic stem and progenitor cells. *Nat Med*. 2018;24(8):1216-1224.