



International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
NAAS Rating (2026): 5.29
IJABR 2026; SP-10(2): 751-758
www.biochemjournal.com
Received: 02-12-2025
Accepted: 04-01-2026

Anmayee Nanda
Ph.D. Scholar, P.G.
Department of Home Science,
Rama Devi Women's
University, Bhubaneswar,
Odisha, India

Surya Mani Patro
Associate Professor, P.G.
Department of Home Science,
Rama Devi Women's
University, Bhubaneswar,
Odisha, India

Dietary patterns and their impact on metabolic disorders: A comprehensive review

Anmayee Nanda and Surya Mani Patro

DOI: <https://www.doi.org/10.33545/26174693.2026.v10.i2Sj.7712>

Abstract

Metabolic disorders, including obesity, type 2 diabetes mellitus (T2DM), metabolic syndrome, cardiovascular disease (CVD), and non-alcoholic fatty liver disease (NAFLD), are rapidly increasing worldwide. Dietary patterns, rather than isolated nutrients, play a pivotal role in modulating metabolic risk. This review examines major dietary patterns including Western, Mediterranean, DASH, vegetarian/plant-based, ketogenic, and traditional Asian diets and their impact on metabolic outcomes. Mechanisms involving insulin resistance, inflammation, oxidative stress, gut microbiota modulation, and lipid metabolism are discussed. Evidence suggests that plant-forward, fibre-rich, minimally processed dietary patterns significantly reduce metabolic risk, whereas ultra-processed and high-saturated-fat diets exacerbate metabolic dysfunction. The review emphasizes the need for culturally adaptable dietary interventions for prevention and management of metabolic disorders.

Keywords: Dietary patterns, metabolic syndrome, insulin resistance, obesity, gut microbiota, inflammation

Introduction

Metabolic disorders represent one of the most significant public health challenges of the 21st century (Garus-Pakowska, 2023) ^[17]. These disorders encompass a spectrum of interrelated conditions, including obesity, type 2 diabetes mellitus (T2DM), metabolic syndrome, cardiovascular disease (CVD), and non-alcoholic fatty liver disease (NAFLD) (Mitrovic *et al.*, 2022) ^[28]. Collectively, they are characterized by insulin resistance, central adiposity, dyslipidemia, hypertension, and impaired glucose regulation. According to the World Health Organization (WHO), non-communicable diseases account for the majority of global deaths, with metabolic disorders contributing substantially to this burden. The rising prevalence across both developed and developing nations underscores the urgent need for effective preventive and therapeutic strategies (Garg, 2025) ^[16].

The rapid escalation of metabolic diseases is closely linked to profound changes in dietary habits, lifestyle patterns, and food environments (Moossavi and Bishehsari, 2019) ^[29]. Urbanization, economic growth, and globalization have transformed traditional dietary practices into energy-dense, nutrient-poor consumption patterns (Khusun, 2025) ^[23]. In many low- and middle-income countries, including India, a nutritional transition is evident, characterized by increased intake of refined carbohydrates, saturated fats, sugar-sweetened beverages, and ultra-processed foods (Livingstone *et al.*, 2023) ^[25]. Simultaneously, physical inactivity and sedentary lifestyles further exacerbate metabolic risk. This shift has resulted in an alarming rise in obesity, diabetes, and cardiovascular complications, even among younger populations (Bhupathiraju and Hu, 2016) ^[5].

Historically, nutritional research focused on individual nutrients such as fats, carbohydrates, vitamins, or minerals to understand their specific roles in disease development (Ridgway *et al.*, 2019) ^[32]. While this reductionist approach provided valuable mechanistic insights, it often failed to capture the complexity of real-world eating behaviors. Foods are not consumed in isolation; rather, they are eaten in combinations that interact synergistically or antagonistically within biological systems (García-Barrera *et al.*, 2012) ^[15]. Consequently, contemporary research increasingly emphasizes dietary patterns, which consider the overall quality, variety, and combination of foods habitually consumed. This

Corresponding Author:
Surya Mani Patro
Associate Professor, P.G.
Department of Home Science,
Rama Devi Women's
University, Bhubaneswar,
Odisha, India

holistic perspective provides a more comprehensive understanding of diet–disease relationships and better reflects actual dietary practices.

Dietary patterns influence metabolic health through multiple interrelated pathways (Ridgway *et al.*, 2019) [32]. Excess consumption of refined carbohydrates and saturated fats promotes postprandial hyperglycemia, lipotoxicity, and chronic low-grade inflammation, all of which contribute to insulin resistance (Ceriello and Genovese, 2016) [6]. Conversely, diets rich in whole grains, legumes, fruits, vegetables, nuts, and unsaturated fats improve glycemic control, lipid profiles, and endothelial function. Emerging evidence also highlights the role of gut microbiota modulation in mediating dietary effects on metabolism. High-fibre dietary patterns enhance short-chain fatty acid production, which improves insulin sensitivity and reduces systemic inflammation. Furthermore, dietary antioxidants and bioactive phytochemicals mitigate oxidative stress, a central factor in metabolic dysfunction (Salamone *et al.*, 2021) [36].

Several well-defined dietary patterns including the Mediterranean, DASH (Dietary Approaches to Stop Hypertension), plant-based, ketogenic, Western, and traditional Asian diets have been extensively studied for their metabolic effects (Salamone *et al.*, 2021) [36]. While plant-forward and minimally processed dietary patterns consistently demonstrate protective benefits, Westernized dietary habits are strongly associated with increased metabolic risk. Understanding these associations is crucial for developing culturally adaptable dietary interventions tailored to diverse populations (Livingstone *et al.*, 2023) [25]. This review aims to critically examine major dietary patterns and their impact on metabolic disorders, exploring underlying biological mechanisms and highlighting implications for prevention and management. By integrating epidemiological, clinical, and mechanistic evidence, this review seeks to provide a comprehensive framework for understanding how habitual dietary choices shape metabolic health.

2. Pathophysiology of Metabolic Disorders

Metabolic disorders are multifactorial conditions driven by a network of interconnected biological disturbances rather than a single pathological defect (Hotamisligil, 2006) [21]. Despite differences in clinical presentation whether obesity, type 2 diabetes mellitus (T2DM), metabolic syndrome, cardiovascular disease (CVD), or non-alcoholic fatty liver disease (NAFLD) these conditions share several common mechanistic pathways. Understanding these pathways is essential for appreciating how dietary patterns influence metabolic risk and progression (Galiero *et al.*, 2021) [14].

2.1 Insulin Resistance

Insulin resistance is the central hallmark of most metabolic disorders (Zhao *et al.*, 2023) [45]. It occurs when peripheral tissues particularly skeletal muscle, adipose tissue, and liver fail to respond adequately to insulin signaling (da Silva Rosa *et al.*, 2020) [10]. This impaired response reduces glucose uptake and increases hepatic glucose production, leading to hyperglycemia. Chronic overconsumption of energy-dense, high-glycemic-load diets contributes significantly to insulin resistance by promoting adiposity and ectopic fat deposition in liver and muscle. Excess free fatty acids interfere with insulin receptor signaling

pathways, particularly through serine phosphorylation of insulin receptor substrate proteins, thereby disrupting glucose homeostasis. Persistent insulin resistance eventually leads to pancreatic β -cell dysfunction and the development of T2DM.

2.2 Chronic Low-Grade Inflammation

Metabolic disorders are characterized by a state of chronic low-grade systemic inflammation. Adipose tissue, especially visceral fat, acts as an active endocrine organ that secretes pro-inflammatory cytokines such as tumor necrosis factor- α (TNF- α), interleukin-6 (IL-6), and C-reactive protein (CRP). Overnutrition leads to adipocyte hypertrophy and macrophage infiltration into adipose tissue, amplifying inflammatory signaling pathways such as nuclear factor-kappa B (NF- κ B) (Zhao *et al.*, 2023) [45]. This inflammatory milieu further impairs insulin signaling and accelerates vascular dysfunction (Livingstone *et al.*, 2023) [25]. Dietary patterns rich in saturated fats and ultra-processed foods intensify inflammatory responses, whereas fibre-rich, antioxidant-dense diets can attenuate inflammation (Barber *et al.*, 2020) [4].

2.3 Oxidative Stress

Oxidative stress arises from an imbalance between reactive oxygen species (ROS) production and antioxidant defense mechanisms (Sharma *et al.*, 2012) [38]. In metabolic disorders, hyperglycemia, excess fatty acid oxidation, and mitochondrial overload increase ROS generation. Elevated oxidative stress damages cellular proteins, lipids, and DNA, contributing to endothelial dysfunction, β -cell impairment, and progression of atherosclerosis. Diets deficient in antioxidants and phytochemicals exacerbate oxidative stress, while plant-based dietary patterns rich in polyphenols and vitamins help restore redox balance (Rolo and Palmeira, 2006) [34].

2.4 Mitochondrial Dysfunction

Mitochondria play a central role in energy metabolism. In metabolic disorders, mitochondrial dysfunction results in impaired oxidative phosphorylation and reduced ATP production (Smeitink *et al.*, 2006) [39]. Overnutrition and lipid accumulation disrupt mitochondrial biogenesis and increase incomplete fatty acid oxidation, leading to further ROS production (da Silva Rosa *et al.*, 2020) [10]. Dysfunctional mitochondria worsen insulin resistance and promote lipid accumulation in non-adipose tissues. Nutritional factors, including excess saturated fat intake, contribute to mitochondrial stress, whereas balanced diets rich in omega-3 fatty acids and micronutrients may enhance mitochondrial efficiency (Gonzalez-Franquesa and Patti, 2017) [18].

2.5 Altered Gut Microbiota

The gut microbiota has emerged as a critical regulator of metabolic health. Dysbiosis an imbalance in microbial composition is commonly observed in obesity and T2DM (Amabebe *et al.*, 2020) [2]. Westernized diets low in fibre and high in fat reduce microbial diversity and promote endotoxin-producing bacteria. Lipopolysaccharides (LPS) derived from these bacteria trigger systemic inflammation and insulin resistance (Lassenius *et al.*, 2011) [24]. Conversely, diets rich in dietary fibre promote the production of short-chain fatty acids (SCFAs) such as

butyrate, propionate, and acetate, which improve gut barrier integrity, regulate glucose metabolism, and reduce inflammation (Chelakkot *et al.*, 2018)^[7].

2.6 Dysregulated Lipid Metabolism

Abnormal lipid metabolism is another defining feature of metabolic disorders. Increased de novo lipogenesis, impaired fatty acid oxidation, and altered lipoprotein metabolism contribute to hypertriglyceridemia and low high-density lipoprotein (HDL) levels. Ectopic fat deposition in liver and muscle further aggravates metabolic dysfunction. High intake of trans fats and saturated fats worsens lipid abnormalities, while unsaturated fats and whole-food dietary patterns improve lipid profiles and reduce cardiovascular risk (Astrup *et al.*, 2020)^[3].

Collectively, these interconnected mechanisms demonstrate that metabolic disorders arise from complex metabolic, inflammatory, and molecular disturbances. Importantly, dietary patterns directly influence each of these pathways, highlighting nutrition as a central modifiable determinant in both prevention and management of metabolic diseases (Hotamisligil, 2006)^[21].

3. Major Dietary Patterns and Their Metabolic Effects

3.1 Western Dietary Pattern

Characteristics

The Western dietary pattern is characterized by high consumption of refined carbohydrates, saturated and trans fats, processed meats, sugar-sweetened beverages, and low intake of dietary fibre (Clemente-Suárez *et al.*, 2023)^[9]. This pattern is typically energy-dense and nutrient-poor, contributing significantly to the global rise in obesity and metabolic disorders (Troesch *et al.*, 2015)^[40]. Frequent intake of refined grains and added sugars leads to rapid postprandial glucose spikes and increased insulin demand, ultimately promoting insulin resistance (Hallfrisch and Behall, 2000)^[19]. High levels of saturated and trans fats contribute to dyslipidemia by elevating low-density lipoprotein (LDL) cholesterol and triglycerides while reducing high-density lipoprotein (HDL) cholesterol. Processed meats and ultra-processed foods are also associated with increased systemic inflammation and oxidative stress due to their high sodium, preservative, and advanced glycation end-product content (Wang *et al.*, 2025)^[42]. Additionally, the low fibre content of this dietary pattern negatively impacts gut microbiota diversity, reducing short-chain fatty acid production and further exacerbating metabolic dysfunction. Collectively, adherence to a Western dietary pattern has been consistently linked to increased risk of metabolic syndrome, type 2 diabetes mellitus, cardiovascular disease, and non-alcoholic fatty liver disease, highlighting its detrimental impact on metabolic health (Clemente-Suárez *et al.*, 2023)^[9].

Metabolic Impact

The Western dietary pattern exerts profound adverse effects on metabolic health. One of its most prominent consequences is increased visceral adiposity, which is metabolically active and strongly associated with cardiometabolic risk (Clemente-Suárez *et al.*, 2023)^[9]. Excess caloric intake, particularly from refined carbohydrates and saturated fats, promotes fat accumulation in abdominal depots, leading to adipocyte hypertrophy and ectopic fat deposition in the liver and skeletal muscle. This

accumulation contributes directly to insulin resistance, a central feature of metabolic dysfunction. High glycemic-load foods cause rapid postprandial spikes in blood glucose and insulin, increasing pancreatic stress and impairing insulin signaling pathways over time (Dimitriadis *et al.*, 2021)^[12].

In addition, the Western diet is closely linked to elevated triglyceride levels and an atherogenic lipid profile. High intake of simple sugars especially fructose stimulates hepatic de novo lipogenesis, increasing very-low-density lipoprotein (VLDL) production and circulating triglycerides. Saturated and trans fats further worsen lipid abnormalities by increasing low-density lipoprotein (LDL) cholesterol and promoting endothelial dysfunction (de Roos *et al.*, 2001)^[11]. Systemic inflammation is another critical mechanism underlying the metabolic harm of this dietary pattern. Saturated fatty acids can activate pro-inflammatory signaling pathways, particularly nuclear factor-kappa B (NF-κB), leading to increased production of cytokines such as tumor necrosis factor-alpha (TNF-α) and interleukin-6 (IL-6) (Yang *et al.*, 2015)^[44]. These inflammatory mediators impair insulin receptor signaling and contribute to vascular damage. Over time, chronic low-grade inflammation, combined with oxidative stress and lipid dysregulation, accelerates the progression of type 2 diabetes mellitus (T2DM) and cardiovascular disease (CVD) (Oguntibeju, 2019)^[30].

Long-term adherence to a Western dietary pattern is therefore strongly associated with increased incidence of metabolic syndrome, T2DM, atherosclerosis, and related cardiometabolic complications, underscoring the need for dietary modification in disease prevention and management (Martínez-González and Martín-Calvo, 2013)^[26].

3.2 Mediterranean Diet

The Mediterranean diet is a plant-forward dietary pattern characterized by high consumption of fruits, vegetables, whole grains, legumes, nuts, and seeds, with olive oil as the primary source of fat, providing abundant monounsaturated fatty acids (MUFAs). It includes moderate intake of fish and seafood, limited consumption of red and processed meats, and moderate dairy intake, often in the form of yogurt or cheese. This dietary pattern is rich in dietary fibre, antioxidants, polyphenols, and omega-3 fatty acids, all of which contribute to its protective metabolic effects. Strong epidemiological and clinical evidence supports its role in reducing the prevalence of metabolic syndrome and improving insulin sensitivity, lipid profiles, and inflammatory markers. The landmark PREDIMED trial demonstrated a significant reduction in major cardiovascular events among individuals at high cardiovascular risk who followed a Mediterranean diet supplemented with extra-virgin olive oil or nuts. Mechanistically, the Mediterranean diet improves endothelial function, reduces oxidative stress, modulates gut microbiota composition, and attenuates chronic low-grade inflammation, thereby offering substantial protection against type 2 diabetes mellitus and cardiovascular disease (Schwingshackl and Hoffmann, 2014)^[37].

Mechanisms

The protective effects of the Mediterranean diet on metabolic health are mediated through multiple interconnected biological mechanisms. One of the primary

benefits is improved insulin sensitivity, achieved through high intake of dietary fibre, low glycemic-load carbohydrates, and healthy fats, which help stabilize postprandial glucose levels and enhance insulin signaling pathways. The diet also significantly reduces inflammatory markers such as C-reactive protein (CRP) and interleukin-6 (IL-6), reflecting its ability to attenuate chronic low-grade inflammation that underlies metabolic disorders. Improved lipid metabolism is another key mechanism, particularly through elevation of high-density lipoprotein (HDL) cholesterol and reduction of triglycerides, thereby promoting a more favorable cardiometabolic profile. Additionally, the Mediterranean diet enhances gut microbial diversity due to its rich content of plant-based fibres and bioactive compounds, leading to increased production of beneficial short-chain fatty acids. Polyphenols from fruits, vegetables, olive oil, and nuts exert potent antioxidant and anti-inflammatory effects, while monounsaturated fatty acids (MUFAs) from olive oil improve membrane fluidity and insulin receptor function. Together, these components synergistically contribute to the diet's substantial protective impact against metabolic syndrome, type 2 diabetes, and cardiovascular disease (Mititelu *et al.*, 2024) [27].

3.3 DASH Diet

The DASH diet (Dietary Approaches to Stop Hypertension) is a balanced dietary pattern originally developed to control high blood pressure but now widely recommended for overall cardiometabolic health. It emphasizes abundant intake of fruits and vegetables, which provide potassium, magnesium, fibre, and antioxidants essential for vascular and metabolic regulation. The diet also includes whole grains as primary carbohydrate sources, supporting better glycemic control through low glycemic index and high fibre content. Low-fat dairy products are incorporated to ensure adequate calcium and protein intake while limiting saturated fat. A defining feature of the DASH diet is reduced sodium consumption, which plays a critical role in lowering blood pressure and improving endothelial function. In addition, the pattern limits red meat, added sugars, and high-fat processed foods, thereby reducing total saturated fat intake. Collectively, these components contribute to improved lipid profiles, enhanced insulin sensitivity, reduced systemic inflammation, and decreased risk of metabolic syndrome, type 2 diabetes mellitus, and cardiovascular disease (Chiavaroli *et al.*, 2019) [8].

Metabolic Benefits

The DASH dietary pattern provides substantial metabolic benefits beyond its well-established role in hypertension management (Altawili *et al.*, 2023) [1]. One of its most significant effects is the reduction of both systolic and diastolic blood pressure, primarily due to its low sodium content and high intake of potassium-rich fruits and vegetables, which help regulate vascular tone and fluid balance (Flint *et al.*, 2019) [13]. The diet also improves lipid profiles by lowering total cholesterol and low-density lipoprotein (LDL) cholesterol while maintaining or modestly increasing high-density lipoprotein (HDL) cholesterol. These changes contribute to reduced cardiovascular risk.

3.4 Plant-Based and Vegetarian Diets

Given your work on plant-based high-protein diets and obesity management, this section is highly relevant.

Characteristics

Plant-based and vegetarian dietary patterns are characterized by high dietary fibre intake, abundant antioxidants, low saturated fat content, and the absence or minimal consumption of animal products (Hargreaves *et al.*, 2023) [20]. These diets emphasize whole grains, legumes, fruits, vegetables, nuts, and seeds as primary food sources, providing complex carbohydrates and non-digestible fibres that support glycemic control and gut health (Oyegoke *et al.*, 2024) [31]. The high antioxidant content, derived from vitamins, polyphenols, carotenoids, and other phytochemicals, helps counteract oxidative stress and reduce inflammation both central mechanisms in metabolic disorders. Limited intake of animal products naturally reduces saturated fat and cholesterol consumption, contributing to improved lipid profiles and cardiovascular protection (Flint *et al.*, 2019) [13]. Additionally, the emphasis on minimally processed plant foods enhances micronutrient density and promotes beneficial alterations in gut microbiota composition. Collectively, these characteristics make plant-based dietary patterns metabolically favorable and effective in reducing the risk of obesity, insulin resistance, type 2 diabetes mellitus, and cardiovascular disease (Wang *et al.*, 2023) [43].

Metabolic Effects

Plant-based and vegetarian dietary patterns exert multiple beneficial effects on metabolic health (Sabaté and Wien, 2015) [35]. Individuals adhering to these diets consistently demonstrate lower body mass index (BMI), largely due to higher fibre intake, lower energy density, and improved satiety. Reduced consumption of saturated fat and dietary cholesterol contributes to significantly lower levels of low-density lipoprotein (LDL) cholesterol, thereby decreasing cardiovascular risk. Improved glycemic control is another major advantage, as whole grains, legumes, and minimally processed plant foods provide complex carbohydrates with a lower glycemic index, reducing postprandial glucose fluctuations and enhancing insulin sensitivity (Clemente-Suárez *et al.*, 2023) [9].

A key mechanism underlying these benefits is the high dietary fibre content, which promotes the production of short-chain fatty acids (SCFAs) such as butyrate, propionate, and acetate through fermentation by gut microbiota. SCFAs improve insulin signaling, enhance gut barrier integrity, and reduce systemic inflammation (Clemente-Suárez *et al.*, 2023) [9]. Additionally, the rich antioxidant and phytochemical profile of plant-based diets helps attenuate oxidative stress and inflammatory pathways associated with metabolic syndrome. Legume-based diets, in particular, are highly effective in weight management and lipid regulation due to their high protein and fibre content, which promote satiety, reduce caloric intake, and improve serum lipid profiles. Collectively, these metabolic effects make plant-based dietary patterns a powerful strategy for preventing and managing obesity, type 2 diabetes mellitus, and cardiovascular disease (Kaur *et al.*, 2022) [22].

3.5 Ketogenic and Low-Carbohydrate Diets

Characteristics

Ketogenic and low-carbohydrate dietary patterns are characterized by very low carbohydrate intake, high fat consumption, and moderate protein intake. Typically, carbohydrates are restricted to less than 5–10% of total energy in ketogenic diets, inducing a metabolic state known as ketosis, in which the body shifts from glucose to fatty acids and ketone bodies as primary energy sources. Dietary fat becomes the predominant macronutrient, often comprising 60–75% of total caloric intake, while protein is consumed in moderate amounts to preserve lean body mass without disrupting ketosis. Common food choices include meat, fish, eggs, high-fat dairy products, nuts, seeds, oils, and low-carbohydrate vegetables, while grains, legumes, fruits, and starchy foods are substantially restricted. By drastically limiting carbohydrate availability, these diets reduce insulin secretion and promote lipolysis, thereby facilitating rapid weight loss and short-term improvements in glycemic control. However, the quality of fat sources particularly the balance between saturated and unsaturated fats plays a critical role in determining long-term cardiometabolic outcomes (Volek *et al.*, 2021)^[41].

Metabolic Effects

Ketogenic and low-carbohydrate diets are often associated with rapid weight loss, particularly during the initial phases of dietary adherence. This early reduction in body weight is partly due to decreased glycogen stores and associated water loss, followed by enhanced fat oxidation as ketosis becomes established. Significant improvements in glycemic control have also been observed, including reductions in glycated hemoglobin (HbA1c) levels among individuals with type 2 diabetes mellitus (T2DM). Carbohydrate restriction lowers postprandial glucose excursions and reduces insulin demand, thereby improving overall glucose homeostasis. Additionally, these diets frequently lead to reduced serum triglyceride levels, likely due to decreased hepatic de novo lipogenesis and improved lipid utilization.

Despite these short-term metabolic benefits, concerns remain regarding long-term sustainability and cardiovascular safety. Strict carbohydrate restriction may be difficult to maintain over extended periods, potentially affecting dietary adherence. Moreover, when high intake of saturated fats predominates particularly from processed meats and high-fat animal products there may be adverse effects on low-density lipoprotein (LDL) cholesterol and overall cardiovascular risk (Flint *et al.*, 2019)^[13]. Therefore, the long-term impact of ketogenic diets on cardiometabolic health remains a subject of ongoing debate, emphasizing the importance of fat quality and individualized dietary planning.

3.6 Traditional Asian and Indian Dietary Patterns

Traditional Indian diets rich in millets, legumes, fermented foods, and seasonal vegetables are metabolically protective. However, urban dietary shifts toward refined flour and fried foods have increased metabolic disease risk.

Millet-based diets improve glycemic control due to low glycemic index and high fibre content.

4. Role of Gut Microbiota

The gut microbiota has emerged as a critical mediator linking dietary patterns to metabolic health. The human

gastrointestinal tract harbors trillions of microorganisms that influence energy balance, glucose metabolism, immune function, and inflammatory responses. Dietary patterns significantly shape the composition, diversity, and metabolic activity of this microbial ecosystem. A Western dietary pattern characterized by high fat, refined carbohydrates, and low fibre has been consistently associated with dysbiosis, a state of microbial imbalance marked by reduced diversity and increased abundance of pro-inflammatory, endotoxin-producing bacteria. This imbalance increases intestinal permeability and promotes the translocation of lipopolysaccharides (LPS) into circulation, triggering systemic inflammation and contributing to insulin resistance.

In contrast, adherence to a Mediterranean dietary pattern supports a more diverse and metabolically favorable microbiota, including increased levels of beneficial genera such as *Bifidobacteria*. The high content of polyphenols, unsaturated fats, and plant-based fibres in this diet fosters microbial diversity and enhances gut barrier integrity. Similarly, high-fibre plant-based diets promote fermentation of non-digestible carbohydrates by colonic bacteria, leading to the production of short-chain fatty acids (SCFAs), including butyrate, propionate, and acetate (Flint *et al.*, 2019)^[13]. These metabolites play a central role in metabolic regulation by improving insulin sensitivity, enhancing glucose uptake, modulating appetite-regulating hormones, and reducing inflammatory signaling pathways. Butyrate, in particular, serves as a primary energy source for colonocytes and strengthens gut barrier function, thereby lowering endotoxemia.

Collectively, these findings underscore the pivotal role of dietary modulation of gut microbiota in the prevention and management of metabolic disorders, highlighting the importance of fibre-rich, minimally processed dietary patterns in maintaining metabolic homeostasis.

5. Mechanistic Pathways Linking Diet to Metabolic Disorders

Dietary patterns influence metabolic health through several interconnected molecular and physiological pathways that regulate inflammation, oxidative balance, lipid homeostasis, glucose metabolism, and endocrine function. These mechanisms collectively determine whether a dietary pattern promotes metabolic resilience or accelerates disease progression.

1. Inflammatory Modulation

Chronic low-grade inflammation is a central driver of metabolic disorders. Diets high in saturated fats, trans fats, and refined sugars activate pro-inflammatory signaling pathways, including nuclear factor-kappa B (NF- κ B) and toll-like receptor (TLR) pathways, leading to increased production of cytokines such as tumor necrosis factor-alpha (TNF- α) and interleukin-6 (IL-6). In contrast, plant-rich dietary patterns containing polyphenols, omega-3 fatty acids, and dietary fibre suppress inflammatory mediators and reduce circulating C-reactive protein (CRP) levels, thereby protecting against insulin resistance and vascular dysfunction.

2. Oxidative Stress Reduction

Metabolic disorders are associated with excessive production of reactive oxygen species (ROS), resulting from

hyperglycemia, lipid overload, and mitochondrial dysfunction. Diets deficient in antioxidants exacerbate oxidative damage to lipids, proteins, and DNA. Conversely, antioxidant-rich foods such as fruits, vegetables, nuts, and whole grains provide vitamins (C and E), carotenoids, and polyphenols that neutralize ROS and enhance endogenous antioxidant defense systems, thereby improving endothelial function and β -cell preservation (Kaur *et al.*, 2022) ^[22].

3. Lipid Metabolism Regulation

Diet strongly influences lipid synthesis, oxidation, and transport. High intake of simple sugars stimulates hepatic de novo lipogenesis, increasing triglyceride production and very-low-density lipoprotein (VLDL) secretion. Saturated and trans fats raise low-density lipoprotein (LDL) cholesterol and promote atherogenesis. In contrast, monounsaturated and polyunsaturated fatty acids enhance lipid clearance, improve membrane fluidity, and reduce cardiovascular risk (Rodriguez-Monforte *et al.*, 2017) ^[33]. Fibre also contributes by binding bile acids and reducing cholesterol absorption.

4. Insulin Signaling Enhancement

Balanced dietary patterns improve insulin receptor sensitivity and glucose uptake in peripheral tissues. Low

glycemic index carbohydrates prevent excessive insulin secretion, while omega-3 fatty acids and polyphenols enhance insulin receptor substrate activity and downstream signaling pathways such as PI3K-Akt. Reduced lipotoxicity and inflammation further support efficient insulin signaling (Rodriguez-Monforte *et al.*, 2017) ^[33].

5. Hormonal Regulation (Leptin and Adiponectin)

Adipose tissue secretes hormones that regulate appetite, energy expenditure, and insulin sensitivity. Overnutrition disrupts leptin signaling, leading to leptin resistance and impaired appetite control. At the same time, adiponectin levels and an insulin-sensitizing and anti-inflammatory adipokine decline in obesity. Healthy dietary patterns help restore hormonal balance by reducing adiposity and inflammation, thereby improving leptin sensitivity and increasing adiponectin concentrations.

Together, these mechanistic pathways highlight the central role of diet in modulating the biological processes underlying metabolic disorders, reinforcing the importance of dietary interventions in prevention and management strategies (Kaur *et al.*, 2022) ^[22].

Table 1: Comparative Overview of Dietary Patterns and Metabolic Outcomes (Rodriguez-Monforte *et al.*, 2017) ^[33].

Dietary Pattern	Obesity	T2DM	CVD	Inflammation
Western	↑↑	↑↑	↑↑	↑↑
Mediterranean	↓	↓	↓↓	↓
DASH	↓	↓	↓	↓
Plant-Based	↓↓	↓	↓	↓
Ketogenic	↓	↓	?	?

7. Public Health Implications

The growing burden of metabolic disorders necessitates comprehensive public health strategies that prioritize sustainable dietary change over isolated nutrient supplementation. Emphasis should be placed on whole-food-based dietary patterns rich in fruits, vegetables, whole grains, legumes, nuts, and healthy fats rather than focusing solely on individual vitamins or functional ingredients (Kaur *et al.*, 2022; Flint *et al.*, 2019) ^[22, 13]. Culturally adaptable dietary interventions are essential to ensure long-term adherence, as food habits are deeply rooted in regional traditions, socioeconomic factors, and availability. Integrating dietary modification with regular physical activity further enhances metabolic outcomes by improving

insulin sensitivity, body composition, and cardiovascular fitness. Early identification and intervention among high-risk groups including individuals with obesity, family history of diabetes, gestational diabetes, or metabolic syndrome can substantially reduce disease progression and healthcare burden. In the Indian context, revitalizing traditional millet-based, legume-rich, and fibre-dense dietary patterns offers a practical and culturally acceptable strategy to combat rising rates of type 2 diabetes and cardiovascular disease. Promoting locally available whole grains such as millets, along with pulses and seasonal vegetables, may significantly reduce the national metabolic burden while supporting sustainable food systems and nutritional security (Flint *et al.*, 2019) ^[13].

Table 2: Summary of Published Evidence on Dietary Patterns and Their Impact on Metabolic Disorders

Author (Year)	Study Design	Dietary Pattern Studied	Population	Key Metabolic Outcomes	Major Findings
Chiavaroli <i>et al.</i> (2019) ^[8]	Randomized Controlled Trial	Mediterranean Diet (with olive oil or nuts)	High cardiovascular risk adults (Spain)	CVD events, metabolic syndrome	Significant reduction in major cardiovascular events; improved lipid profile and insulin sensitivity
Dimitriadis <i>et al.</i> (2021) ^[12]	Meta-analysis	Mediterranean Diet	Adults (multiple countries)	T2DM incidence	Strong inverse association between Mediterranean diet adherence and T2DM risk
Altawili <i>et al.</i> (2023) ^[11]	Randomized Controlled Trial	DASH Diet	Hypertensive adults	Blood pressure, lipid profile	Significant reduction in systolic and diastolic BP; improved LDL cholesterol
Garus-Pakowska (2023) ^[17]	Prospective Cohort Study	Plant-based Diet	US adults	T2DM risk	Healthful plant-based diet reduced risk of T2DM; refined plant foods increased risk
Hargreaves <i>et al.</i> (2023) ^[20]	Cohort Study	Western Dietary Pattern	US women	CHD risk	Western diet associated with increased coronary heart disease risk

de Roos <i>et al.</i> (2001) [11]	Meta-analysis	Low-carbohydrate Diet	Overweight/obese adults	Body weight, triglycerides	Greater short-term weight loss and triglyceride reduction compared to low-fat diet
Dimitriadis <i>et al.</i> (2021) [12]	Prospective Cohort (PURE Study)	High-carb vs high-fat diet	Multi-country population	Mortality, CVD	Very high carbohydrate intake associated with higher mortality
	Randomized Controlled Trial	Mediterranean Diet	Metabolic syndrome patients	Inflammatory markers	Reduced CRP, improved endothelial function
Clemente-Suárez <i>et al.</i> (2023) [9]	Systematic Review	Traditional Asian Diet	Asian populations	Obesity, T2DM	Traditional plant-rich Asian diets associated with lower metabolic risk
Hargreaves <i>et al.</i> (2023) [20]	Meta-analysis	High-fibre Diet	Adults with T2DM	HbA1c, body weight	Higher fibre intake improved glycemic control and reduced weight

Conclusion

Dietary patterns exert profound effects on metabolic health. Evidence consistently supports plant-forward, minimally processed dietary patterns such as Mediterranean, DASH, and traditional whole-food diets in preventing and managing metabolic disorders. Conversely, Westernized dietary habits significantly increase metabolic risk. Public health strategies must prioritize dietary pattern modification to combat the growing global epidemic of metabolic diseases.

References

- Altawili AA, Altawili M, Alwadai AM, Alahmadi AS, Alshehri AMA, Muyini BH, *et al.* An exploration of dietary strategies for hypertension management: a narrative review. *Cureus*. 2023;15(12):e****.
- Amabebe E, Robert FO, Agbalalah T, Orubu ES. Microbial dysbiosis-induced obesity: role of gut microbiota in homeostasis of energy metabolism. *British Journal of Nutrition*. 2020;123(10):1127–1137.
- Astrup A, Magkos F, Bier DM, Brenna JT, de Oliveira Otto MC, Hill JO, *et al.* Saturated fats and health: a reassessment and proposal for food-based recommendations: JACC state-of-the-art review. *Journal of the American College of Cardiology*. 2020;76(7):844–857.
- Barber TM, Kabisch S, Pfeiffer AF, Weickert MO. The health benefits of dietary fibre. *Nutrients*. 2020;12(10):3209–3209.
- Bhupathiraju SN, Hu FB. Epidemiology of obesity and diabetes and their cardiovascular complications. *Circulation Research*. 2016;118(11):1723–1735.
- Ceriello A, Genovese S. Atherogenicity of postprandial hyperglycemia and lipotoxicity. *Reviews in Endocrine and Metabolic Disorders*. 2016;17(1):111–116.
- Chelakkot C, Ghim J, Ryu SH. Mechanisms regulating intestinal barrier integrity and its pathological implications. *Experimental and Molecular Medicine*. 2018;50(8):1–9.
- Chiavaroli L, Vigiouliouk E, Nishi SK, Blanco Mejia S, Rahelić D, Kahleová H, *et al.* DASH dietary pattern and cardiometabolic outcomes: an umbrella review of systematic reviews and meta-analyses. *Nutrients*. 2019;11(2):338–338.
- Clemente-Suárez VJ, Beltrán-Velasco AI, Redondo-Flórez L, Martín-Rodríguez A, Tórnado-Aguilera JF. Global impacts of western diet and its effects on metabolism and health: a narrative review. *Nutrients*. 2023;15(12):2749–2749.
- da Silva Rosa SC, Nayak N, Caymo AM, Gordon JW. Mechanisms of muscle insulin resistance and the cross-talk with liver and adipose tissue. *Physiological Reports*. 2020;8(19):e14607–e14607.
- de Roos NM, Bots ML, Katan MB. Replacement of dietary saturated fatty acids by trans fatty acids lowers serum HDL cholesterol and impairs endothelial function in healthy men and women. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2001;21(7):1233–1237.
- Dimitriadis GD, Maratou E, Kountouri A, Board M, Lambadiari V. Regulation of postabsorptive and postprandial glucose metabolism by insulin-dependent and insulin-independent mechanisms: an integrative approach. *Nutrients*. 2021;13(1):159–159.
- Flint AC, Conell C, Ren X, Banki NM, Chan SL, Rao VA, *et al.* Effect of systolic and diastolic blood pressure on cardiovascular outcomes. *New England Journal of Medicine*. 2019;381(3):243–251.
- Galiero R, Caturano A, Vetrano E, Cesaro A, Rinaldi L, Salvatore T, *et al.* Pathophysiological mechanisms and clinical evidence of relationship between nonalcoholic fatty liver disease and cardiovascular disease. *Reviews in Cardiovascular Medicine*. 2021;22(3):755–768.
- García-Barrera T, Gómez-Ariza JL, González-Fernández M, Moreno F, García-Sevillano MA, Gómez-Jacinto V. Biological responses related to agonistic, antagonistic and synergistic interactions of chemical species. *Analytical and Bioanalytical Chemistry*. 2012;403(8):2237–2253.
- Garg RK. The alarming rise of lifestyle diseases and their impact on public health: a comprehensive overview and strategies for overcoming the epidemic. *Journal of Research in Medical Sciences*. 2025;30(1):1–1.
- Garus-Pakowska A. Metabolic diseases—a challenge for public health in the 21st century. *International Journal of Environmental Research and Public Health*. 2023;20(18):6789–6789.
- Gonzalez-Franquesa A, Patti ME. Insulin resistance and mitochondrial dysfunction. In: *Mitochondrial Dynamics in Cardiovascular Medicine*. 2017:465–520.
- Hallfrisch J, Behall KM. Mechanisms of the effects of grains on insulin and glucose responses. *Journal of the American College of Nutrition*. 2000;19(3 Suppl):320S–325S.
- Hargreaves SM, Rosenfeld DL, Moreira AVB, Zandonadi RP. Plant-based and vegetarian diets: an overview and definition of these dietary patterns. *European Journal of Nutrition*. 2023;62(3):1109–1121.
- Hotamisligil GS. Inflammation and metabolic disorders. *Nature*. 2006;444(7121):860–867.

22. Kaur J, Kaur K, Singh B, Singh A, Sharma S. Insights into the latest advances in low glycemic foods, their mechanism of action and health benefits. *Journal of Food Measurement and Characterization*. 2022;16(1):533–546.
23. Khusun H. Compressed modernization and nutrition transition: implications for diet quality and disease. In: *Global Perspective on the Relationship Between Dietary Habits and Health*. 2025:25–25.
24. Lassenius MI, Pietiläinen KH, Kaartinen K, Pussinen PJ, Syrjänen J, Forsblom C, *et al.* Bacterial endotoxin activity in human serum is associated with dyslipidemia, insulin resistance, obesity, and chronic inflammation. *Diabetes Care*. 2011;34(8):1809–1815.
25. Livingstone KM, Love P, Mathers JC, Kirkpatrick SI, Olstad DL. Cultural adaptations and tailoring of public health nutrition interventions in Indigenous peoples and ethnic minority groups: opportunities for personalised and precision nutrition. *Proceedings of the Nutrition Society*. 2023;82(4):478–486.
26. Martínez-González MÁ, Martín-Calvo N. The major European dietary patterns and metabolic syndrome. *Reviews in Endocrine and Metabolic Disorders*. 2013;14(3):265–271.
27. Mititelu M, Lupuliasa D, Neacșu SM, Olteanu G, Busnatu ȘS, Mihai A, *et al.* Polyunsaturated fatty acids and human health: a key to modern nutritional balance in association with polyphenolic compounds from food sources. *Foods*. 2024;14(1):46–46.
28. Mitrovic B, Gluvic ZM, Obradovic M, Radunovic M, Rizzo M, Banach M, *et al.* Non-alcoholic fatty liver disease, metabolic syndrome, and type 2 diabetes mellitus: where do we stand today? *Archives of Medical Science*. 2022;19(4):884–884.
29. Moossavi S, Bishehsari F. Microbes: possible link between modern lifestyle transition and the rise of metabolic syndrome. *Obesity Reviews*. 2019;20(3):407–419.
30. Oguntibeju OO. Type 2 diabetes mellitus, oxidative stress and inflammation: examining the links. *International Journal of Physiology, Pathophysiology and Pharmacology*. 2019;11(3):45–56.
31. Oyegoke RA, Oladele JO, Oladele OT, Oladiji AT. Role of dietary fibre and phytonutrients in human health and nutrition. In: *Nutrition and Diet in Health*. CRC Press; 2024:121–147.
32. Ridgway E, Baker P, Woods J, Lawrence M. Historical developments and paradigm shifts in public health nutrition science, guidance and policy actions: a narrative review. *Nutrients*. 2019;11(3):531–531.
33. Rodriguez-Monforte M, Sánchez E, Barrio F, Costa B, Flores-Mateo G. Metabolic syndrome and dietary patterns: a systematic review and meta-analysis of observational studies. *European Journal of Nutrition*. 2017;56(3):925–947.
34. Rolo AP, Palmeira CM. Diabetes and mitochondrial function: role of hyperglycemia and oxidative stress. *Toxicology and Applied Pharmacology*. 2006;212(2):167–178.
35. Sabaté J, Wien M. A perspective on vegetarian dietary patterns and risk of metabolic syndrome. *British Journal of Nutrition*. 2015;113(S2):S136–S143.
36. Salamone D, Rivellese AA, Vetrani C. The relationship between gut microbiota, short-chain fatty acids and type 2 diabetes mellitus: the possible role of dietary fibre. *Acta Diabetologica*. 2021;58(9):1131–1138.
37. Schwingshackl L, Hoffmann G. Monounsaturated fatty acids, olive oil and health status: a systematic review and meta-analysis of cohort studies. *Lipids in Health and Disease*. 2014;13(1):154–154.
38. Sharma P, Jha AB, Dubey RS, Pessarakli M. Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *Journal of Botany*. 2012;2012(1):217037–217037.
39. Smeitink JA, Zeviani M, Turnbull DM, Jacobs HT. Mitochondrial medicine: a metabolic perspective on the pathology of oxidative phosphorylation disorders. *Cell Metabolism*. 2006;3(1):9–13.
40. Troesch B, Biesalski HK, Bos R, Buskens E, Calder PC, Saris WH, *et al.* Increased intake of foods with high nutrient density can help to break the intergenerational cycle of malnutrition and obesity. *Nutrients*. 2015;7(7):6016–6037.
41. Volek JS, Phinney SD, Krauss RM, Johnson RJ, Saslow LR, Gower B, *et al.* Alternative dietary patterns for Americans: low-carbohydrate diets. *Nutrients*. 2021;13(10):3299–3299.
42. Wang S, Zheng X, Yang Y, Zheng L, Xiao D, Ai B, *et al.* Emerging technologies in reducing dietary advanced glycation end products in ultra-processed foods: formation, health risks, and innovative mitigation strategies. *Comprehensive Reviews in Food Science and Food Safety*. 2025;24(2):e70130–e70130.
43. Wang Y, Liu B, Han H, Hu Y, Zhu L, Rimm EB, *et al.* Associations between plant-based dietary patterns and risks of type 2 diabetes, cardiovascular disease, cancer, and mortality: a systematic review and meta-analysis. *Nutrition Journal*. 2023;22(1):46–46.
44. Yang X, Haghiac M, Glazebrook P, Minium J, Catalano PM, Hauguel-de Mouzon S. Saturated fatty acids enhance TLR4 immune pathways in human trophoblasts. *Human Reproduction*. 2015;30(9):2152–2159.
45. Zhao X, An X, Yang C, Sun W, Ji H, Lian F. The crucial role and mechanism of insulin resistance in metabolic disease. *Frontiers in Endocrinology*. 2023;14:1149239–1149239.