



International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
 ISSN Online: 2617-4707
 NAAS Rating (2025): 5.29
 IJABR 2025; 9(2): 203-207
www.biochemjournal.com
 Received: 23-12-2024
 Accepted: 27-01-2025

Martín A Fernández
 Centro de Investigación en
 Biotecnología de Alimentos,
 Universidad Nacional del
 Litoral, Santa Fe, Argentina

Lucía M Rodríguez
 Centro de Investigación en
 Biotecnología de Alimentos,
 Universidad Nacional del
 Litoral, Santa Fe, Argentina

Pablo E Morales
 Departamento de Química
 Analítica, Universidad de
 Buenos Aires, Buenos Aires,
 Argentina

Carolina S Vega
 Departamento de Química
 Analítica, Universidad de
 Buenos Aires, Buenos Aires,
 Argentina

Corresponding Author:
 Martín A Fernández
 Centro de Investigación en
 Biotecnología de Alimentos,
 Universidad Nacional del
 Litoral, Santa Fe, Argentina

Development of kinetic models for predicting alpha-amylase activity during enzymatic hydrolysis of native and modified starch substrates

Martín A Fernández, Lucía M Rodríguez, Pablo E Morales and Carolina S Vega

DOI: <https://www.doi.org/10.33545/26174693.2025.v9.i2c.7537>

Abstract

Alpha-amylase enzymes play essential roles in starch processing industries for producing maltodextrins, glucose syrups, and fermentable sugars. This research developed predictive kinetic models describing alpha-amylase activity during enzymatic hydrolysis of native and chemically modified corn starch substrates. Michaelis-Menten kinetic parameters were determined for fungal, bacterial, and plant-derived alpha-amylases acting on native, cross-linked, waxy, and high-amylose starch varieties. The K_m values ranged from 6.2 to 15.8 milligrams per milliliter depending on substrate modification and enzyme source. Modified first-order kinetic models accurately predicted hydrolysis progress with correlation coefficients exceeding 0.96 across all substrate-enzyme combinations. Temperature and pH optima varied substantially among enzyme sources, informing process optimization strategies. The developed models provide practical tools for industrial starch hydrolysis process design and control.

Keywords: Alpha-amylase, starch hydrolysis, enzyme kinetics, Michaelis-Menten, modified starch, dextrose equivalent, maltodextrin, process modeling

Introduction

Starch represents the most abundant storage polysaccharide in plants and constitutes a major industrial raw material for producing diverse products including sweeteners, textiles, paper, and biofuels. The enzymatic modification of starch using alpha-amylase enzymes offers advantages over acid-catalyzed hydrolysis including milder processing conditions, improved product specificity, and reduced environmental impact. Alpha-amylase catalyzes the hydrolysis of alpha-1, 4-glycosidic linkages in starch polymers, generating dextrans and oligosaccharides of varying chain lengths depending on reaction conditions and enzyme characteristics.

Industrial starch processing employs alpha-amylases from diverse microbial and plant sources exhibiting distinct catalytic properties optimized for specific applications. Bacterial alpha-amylases from *Bacillus* species demonstrate thermostability suitable for high-temperature liquefaction processes, whereas fungal enzymes from *Aspergillus* typically operate at lower temperatures during saccharification stages. Plant-derived alpha-amylases from malted cereals find application in brewing and baking industries where specific flavor development accompanies starch conversion.

Native starch substrates present heterogeneous granular structures comprising linear amylose and branched amylopectin polymers in varying proportions depending on botanical source. Chemical modifications including cross-linking, oxidation, and esterification alter starch granule architecture and susceptibility to enzymatic attack. Understanding how these structural modifications affect alpha-amylase kinetics enables optimization of industrial processes utilizing modified starch feedstocks for specialty product manufacturing.

Kinetic modeling of enzymatic reactions provides quantitative frameworks for process design, optimization, and control. The Michaelis-Menten equation describes steady-state enzyme kinetics through maximum velocity and substrate affinity parameters that can be experimentally determined and correlated with reaction conditions. Extension to modified kinetic models incorporating substrate inhibition, product inhibition, and enzyme deactivation

phenomena enables more accurate prediction of industrial-scale hydrolysis processes.

Material and Methods

Material

Commercial alpha-amylase preparations were obtained from established enzyme suppliers for comparative kinetic evaluation. Fungal alpha-amylase from *Aspergillus oryzae* was supplied by Sigma-Aldrich with declared activity of 30 units per milligram solid. Bacterial thermostable alpha-amylase from *Bacillus licheniformis* was obtained from Novozymes with activity specification of 120 KNU per gram. Plant alpha-amylase was extracted from malted barley obtained from a regional maltster using standard aqueous extraction methodology.

Native corn starch was obtained from Ingredion Argentina as food-grade material with approximately 27 percent amylose content. Cross-linked starch was prepared through phosphorylation treatment using sodium trimetaphosphate according to established procedures. Waxy corn starch containing less than 1 percent amylose was obtained from the same supplier. High-amylose corn starch with approximately 70 percent amylose content was sourced as a specialty product for comparative analysis.

Methods

Enzyme activity assays employed the dinitrosalicylic acid method for reducing sugar quantification. Starch substrate suspensions were prepared at concentrations ranging from 0.5 to 50 milligrams per milliliter in appropriate buffer systems. Reaction mixtures containing enzyme and substrate were incubated at controlled temperature with aliquots withdrawn at timed intervals for analysis. Initial velocity measurements were obtained from linear portions of progress curves during the first five to ten percent of substrate conversion.

Michaelis-Menten kinetic parameters were determined through nonlinear regression analysis of initial velocity versus substrate concentration data using GraphPad Prism software. Lineweaver-Burk double reciprocal plots provided alternative parameter estimation and identification of inhibition patterns. Temperature and pH profiles were constructed by measuring activity under standardized conditions across relevant ranges with appropriate buffering systems maintaining ionic strength.

Hydrolysis progress experiments monitored starch conversion and dextrose equivalent development over extended reaction periods. Starch conversion was determined iodometrically through loss of blue-black starch-iodine complex. Dextrose equivalent was calculated from reducing sugar concentration relative to total carbohydrate content expressed as glucose equivalents. Modified first-order kinetic models were fitted to conversion data to extract apparent rate constants for predictive modeling applications.

Quality control

Enzyme preparation quality was verified through activity assays using standardized protocols prior to kinetic experiments. Activity values were required to fall within ten percent of manufacturer specifications for acceptance. Protein concentration was determined by Bradford assay to

enable specific activity calculations and ensure consistent enzyme dosing across experiments. Storage stability was monitored through periodic activity measurements with preparations demonstrating greater than five percent activity loss excluded from subsequent analyses.

Starch substrate characterization included moisture content determination by oven drying, ash content by muffle furnace incineration, and protein content by Kjeldahl nitrogen analysis. Amylose content was quantified using the iodine colorimetric method with potato amylose standards. Particle size distribution was assessed by laser diffraction to ensure consistency across substrate batches. Modified starch degree of substitution was verified through phosphorus content analysis for cross-linked samples.

Analytical method validation encompassed determination of linearity, precision, accuracy, and detection limits for reducing sugar and total carbohydrate quantification methods. Calibration curves demonstrated correlation coefficients exceeding 0.999 across working concentration ranges. Within-day precision expressed as coefficient of variation remained below three percent for replicate analyses. Recovery experiments with spiked samples confirmed accuracy within 98 to 102 percent of expected values.

Instrumentation

Spectrophotometric measurements utilized the Shimadzu UV-1800 double-beam spectrophotometer equipped with temperature-controlled cuvette holder. Wavelength accuracy was verified monthly using holmium oxide filter standards. Absorbance accuracy was confirmed using potassium dichromate solutions at certified concentrations. The instrument demonstrated photometric linearity across the full absorbance range employed in reducing sugar and iodometric assays.

Temperature control during enzymatic reactions employed the Julabo F25-ME circulating water bath maintaining setpoint within plus or minus 0.1 degrees Celsius. Calibration against NIST-traceable thermometers confirmed accuracy at multiple temperature levels. pH measurements utilized the Mettler Toledo SevenCompact meter with combined glass electrode calibrated daily using NIST-traceable buffer solutions at pH 4.00, 7.00, and 10.00.

Particle size analysis employed the Malvern Mastersizer 3000 laser diffraction system with aqueous dispersion unit. Optical model parameters were optimized for starch granule analysis with refractive index values from published literature. Replicate measurements demonstrated coefficient of variation below two percent for median particle diameter determination. Size standards verified instrument calibration across the relevant measurement range.

Results

Kinetic parameter determination revealed substantial differences among substrate types and enzyme sources. Native corn starch demonstrated intermediate susceptibility to alpha-amylase hydrolysis with Michaelis constant values of 8.5 milligrams per milliliter for fungal enzyme and 7.2 milligrams per milliliter for bacterial enzyme. Maximum velocity values reached 85 and 92 micromoles per minute per milligram enzyme protein, respectively, under optimal conditions.

Table 1: Michaelis-Menten kinetic parameters for different substrate-enzyme combinations

Substrate	Enzyme Source	K _m (mg/mL)	V _{max} (μmol/min/mg)	k _{cat} /K _m
Native Corn	Fungal	8.5±0.6	85±4	10.0
Native Corn	Bacterial	7.2±0.5	92±5	12.8
Cross-linked	Fungal	12.3±0.9	62±3	5.0
Waxy	Fungal	6.2±0.4	98±5	15.8
High-amylose	Fungal	15.8±1.2	72±4	4.6

Values represent mean ± standard deviation from triplicate determinations. Catalytic efficiency (k_{cat}/K_m) expressed as mL/mg/min

Cross-linked starch exhibited reduced susceptibility to enzymatic hydrolysis compared to native substrate, manifesting as elevated K_m values of 12.3 milligrams per milliliter and decreased maximum velocity of 62 micromoles per minute per milligram. The phosphate cross-

links restrict granule swelling and limit enzyme accessibility to glycosidic bonds within the modified polymer matrix. Catalytic efficiency expressed as k_{cat}/K_m ratio decreased by approximately fifty percent compared to native starch hydrolysis.




Fig 1: Michaelis-Menten kinetics of alpha-amylase with different starch substrates. Panel A shows velocity versus substrate concentration curves. Panel B presents Lineweaver-Burk double reciprocal plots for kinetic parameter determination

Waxy starch demonstrated enhanced hydrolysis susceptibility with lowest K_m value of 6.2 milligrams per milliliter and highest maximum velocity of 98 micromoles per minute per milligram. The absence of amylose and corresponding reduction in crystalline regions facilitates

enzyme access to the highly branched amylopectin Substrate. Conversely, high-amylose starch presented the greatest resistance to hydrolysis with K_m of 15.8 milligrams per milliliter, reflecting the crystalline nature of linear amylose chains impeding enzyme penetration.




Fig 2: Effect of temperature (A) and pH (B) on alpha-amylase activity from different sources. Fungal enzyme shows optimal activity at 55°C and pH 5.0; bacterial enzyme at 70°C and pH 6.5; plant enzyme at 45°C and pH 5.5

Temperature optimization experiments confirmed distinct thermal profiles among enzyme sources. Bacterial alpha-amylase exhibited optimal activity at 70 degrees Celsius with retention of greater than eighty percent activity at 80 degrees Celsius, consistent with thermostable characteristics

Suitable for high-temperature liquefaction. Fungal enzyme demonstrated optimal activity at 55 degrees Celsius with rapid inactivation above 65 degrees Celsius. Plant alpha-amylase from malted barley showed lowest optimal temperature of 45 degrees Celsius.

Table 2: First-order rate constants for starch hydrolysis

Substrate	k (min ⁻¹)	Max conversion (%)	R ²
Native corn	0.025±0.002	92±2	0.985
Cross-linked	0.015±0.001	78±3	0.978
Waxy	0.035±0.003	98±1	0.992
High-amylose	0.012±0.001	68±4	0.968

Rate constants determined using fungal alpha-amylase at 55°C and pH 5.0. R² indicates goodness of fit for first-order kinetic model




Fig 3: Enzymatic hydrolysis progress showing starch conversion (A) and dextrose equivalent development (B) over time for different starch substrates. Shaded regions in Panel B indicate product classification by DE range

Hydrolysis progress experiments demonstrated varying conversion kinetics among substrate types. Modified first-order kinetic models accurately described conversion progress with correlation coefficients exceeding 0.96 for all substrate-enzyme combinations. Apparent rate constants ranged from 0.012 per minute for high-amylose starch to 0.035 per minute for waxy starch. Maximum achievable conversion decreased from 98 percent for waxy starch to 68 percent for high-amylose substrate, reflecting incomplete accessibility of crystalline amylose regions.

Discussion

The kinetic parameters determined in this research provide quantitative foundations for industrial starch hydrolysis process design and optimization. The substantial variation in Michaelis constant values among substrate types directly impacts required enzyme dosages for achieving target conversion levels within specified timeframes. Cross-linked and high-amylose starches require elevated enzyme concentrations to compensate for reduced catalytic efficiency, affecting process economics.

The correlation between starch structural characteristics and alpha-amylase kinetics reflects fundamental relationships between substrate accessibility and enzymatic catalysis. Amylopectin-rich waxy starch presents numerous surface-accessible alpha-1, 4-glycosidic bonds within the branched polymer architecture. Conversely, linear amylose chains

form crystalline double helices resistant to enzymatic penetration, explaining the elevated K_m and reduced maximum conversion observed for high-amylose substrates. Temperature and pH optima differences among enzyme sources enable process flexibility through strategic enzyme selection. Bacterial thermostable alpha-amylase permits high-temperature operations reducing microbial contamination risks and decreasing starch paste viscosity. However, lower-temperature fungal enzyme processes may preserve heat-sensitive product attributes and reduce energy requirements. The developed kinetic models facilitate rational selection based on specific process requirements.

Limitations of the current research include focus on batch reaction conditions that may not fully represent continuous industrial processes. Scale-up effects including mixing efficiency, heat transfer, and enzyme distribution heterogeneity require additional investigation. Future research should address enzyme immobilization strategies enabling continuous processes and extended enzyme utilization, potentially improving process economics for modified starch hydrolysis applications.

Conclusion

Comprehensive kinetic models describing alpha-amylase activity during starch hydrolysis have been developed and validated across diverse substrate-enzyme combinations. Michaelis-Menten parameters quantify the substantial

impact of starch structural modifications on enzymatic susceptibility, with catalytic efficiency varying by factors exceeding three-fold among tested substrates. Modified first-order kinetic models accurately predict hydrolysis progress with practical utility for industrial process design.

The findings demonstrate that waxy starch provides optimal substrate characteristics for efficient α -amylase hydrolysis, while cross-linked and high-amylose substrates require process modifications including elevated enzyme dosages or extended reaction times. Temperature and pH optimization based on enzyme source characteristics enables further process enhancement. The developed kinetic framework provides quantitative tools supporting rational design and control of starch hydrolysis processes across diverse industrial applications.

Future applications of this research include integration of kinetic models into process simulation software for predictive process optimization, development of enzyme cocktail formulations optimized for specific substrate characteristics, and extension to additional modified starch varieties utilized in specialty product manufacturing.

References

1. Van der Maarel MJEC, van der Veen B, Uitdehaag JCM, *et al.* Properties and applications of starch-converting enzymes of the α -amylase family. *J Biotechnol.* 2002;94(2):137-155.
2. Gupta R, Gigras P, Mohapatra H, *et al.* Microbial α -amylases: a biotechnological perspective. *Process Biochem.* 2003;38(11):1599-1616.
3. Bertolini AC. *Starches: Characterization, Properties, and Applications.* Boca Raton (FL): CRC Press; 2010.
4. Tester RF, Karkalas J, Qi X. Starch composition, fine structure and architecture. *J Cereal Sci.* 2004;39(2):151-165.
5. Singh J, Kaur L, McCarthy OJ. Factors influencing the physico-chemical, morphological, thermal and rheological properties of some chemically modified starches for food applications. *Food Hydrocoll.* 2007;21(1):1-22.
6. Miller GL. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal Chem.* 1959;31(3):426-428.
7. Whitaker JR. *Principles of Enzymology for the Food Sciences.* 2nd ed. New York: Marcel Dekker; 1994.
8. Cornish-Bowden A. *Fundamentals of Enzyme Kinetics.* 4th ed. Weinheim: Wiley-VCH; 2012.
9. Goesaert H, Brijs K, Veraverbeke WS, *et al.* Wheat flour constituents: how they impact bread quality, and how to impact their functionality. *Trends Food Sci Technol.* 2005;16(1-3):12-30.
10. Synowiecki J. The use of starch processing enzymes in the food industry. In: Polaina J, MacCabe AP, editors. *Industrial Enzymes.* Dordrecht: Springer; c2007. p. 19-34.
11. Dona AC, Pages G, Gilbert RG, *et al.* Digestion of starch: in vivo and in vitro kinetic models used to characterise oligosaccharide or glucose release. *Carbohydr Polym.* 2010;80(3):599-617.
12. Baks T, Janssen AEM, Boom RM. The effect of carbohydrates on α -amylase activity measurements. *Enzyme Microb Technol.* 2006;39(1):114-119.
13. Wang JP, Quiroga JA, Rosentrater KA. Processing effects on corn protein concentrates as fish feed ingredients. *Trans ASABE.* 2015;58(4):1163-1173.
14. Lehmann U, Robin F. Slowly digestible starch its structure and health implications: A review. *Trends Food Sci Technol.* 2007;18(7):346-355.