



ISSN Print: 2664-9926
 ISSN Online: 2664-9934
 NAAS Rating (2025): 4.82
 IJBS 2025; 7(11): 155-157
www.biologyjournal.net
 Received: 15-09-2025
 Accepted: 18-10-2025

Vimal Balasubramanian
 Department of Structural
 Biochemistry, Cauvery College
 of Biological Sciences, Mysore,
 Karnataka, India

Bhoomika Lakhani
 Department of Structural
 Biochemistry, Cauvery College
 of Biological Sciences, Mysore,
 Karnataka, India

Akshay Khatri
 Department of Structural
 Biochemistry, Cauvery College
 of Biological Sciences, Mysore,
 Karnataka, India

Corresponding Author:
Vimal Balasubramanian
 Department of Structural
 Biochemistry, Cauvery College
 of Biological Sciences, Mysore,
 Karnataka, India

Redox enzyme systems and reactive oxygen species metabolism in plant chloroplasts

Vimal Balasubramanian, Bhoomika Lakhani and Akshay Khatri

DOI: <https://www.doi.org/10.33545/26649926.2025.v7.i11c.727>

Abstract

A veteran plant physiologist once described chloroplasts as factories that run on sunlight but risk burning down from their own exhaust. That exhaust—reactive oxygen species (ROS)—is an inevitable byproduct of photosynthetic electron transport, and chloroplasts have evolved an elaborate suite of redox enzymes to manage it. This research measured the activities of six antioxidant enzymes in chloroplasts isolated from spinach (*Spinacia oleracea*) leaves exposed to control and high-light conditions at the Cauvery College of Biological Sciences, Mysore, India, from April 2023 to July 2024. Superoxide dismutase (SOD), ascorbate peroxidase (APX), catalase (CAT), glutathione reductase (GR), dehydroascorbate reductase (DHAR), and monodehydroascorbate reductase (MDHAR) were assayed spectrophotometrically in purified chloroplast fractions. Under high light ($1,500 \mu\text{mol m}^{-2} \text{s}^{-1}$ for 4 hours), all six enzymes showed significant activity increases ranging from 76% (CAT) to 89% (MDHAR). Superoxide and hydrogen peroxide levels were elevated 2.3-fold and 1.8-fold respectively. Despite the enzymatic response, a net ROS surplus of approximately 8% remained under high light. These results quantify the antioxidant capacity of chloroplast redox systems and identify the residual ROS burden that may contribute to photoinhibition.

Keywords: Redox enzymes, reactive oxygen species, chloroplasts, plant biochemistry, oxidative metabolism, ascorbate-glutathione cycle, photoinhibition, antioxidant defense

Introduction

An old lab anecdote holds that a graduate student, puzzled by why plants bother producing antioxidants, was told to stare directly at the sun for five seconds—the pain in his eyes was essentially what chloroplasts deal with every clear afternoon. Photosynthetic electron transport inevitably generates superoxide (O_2^-) at photosystem I and singlet oxygen ($^1\text{O}_2$) at photosystem II when absorbed light energy exceeds what carbon fixation can consume [1]. Without efficient scavenging, these ROS would damage lipids, proteins, and DNA within seconds [2]. Chloroplasts rely on the ascorbate-glutathione (Asada-Halliwell) cycle as their primary ROS detoxification pathway, involving the sequential action of SOD, APX, MDHAR, DHAR, and GR [3]. Catalase provides an additional route for H_2O_2 removal, though its chloroplastic contribution is debated [4]. While the individual enzymes in this system are well characterized, quantitative measurements of the entire pathway operating in intact chloroplasts under defined stress conditions—allowing a mass balance of ROS production versus scavenging—remain limited [5]. This research aimed to measure the activities of all six major antioxidant enzymes in isolated chloroplasts under control and high-light stress and to construct a ROS budget estimating the net oxidative burden.

Materials and Methods

Materials

This research was conducted at the Department of Structural Biochemistry, Cauvery College of Biological Sciences, Mysore, Karnataka, India, from April 2023 to July 2024. Institutional approval was obtained (CCBS-RC/2023-041). Spinach (*Spinacia oleracea* cv. Bloomsdale) was grown in a controlled environment chamber at 22°C , 60% relative humidity, under moderate light ($400 \mu\text{mol m}^{-2} \text{s}^{-1}$, 14 h photoperiod) for 6 weeks. Fully expanded leaves from 6-week-old plants were harvested at the start of the light period. Intact chloroplasts were isolated by differential centrifugation through Percoll gradients following the method of

Lilley *et al.* [6]. Chloroplast integrity was confirmed by ferricyanide-dependent oxygen evolution (>85% intactness required).

Methods

Isolated chloroplasts ($0.5 \text{ mg chlorophyll mL}^{-1}$) were illuminated at either $400 \mu\text{mol m}^{-2} \text{ s}^{-1}$ (control) or $1,500 \mu\text{mol m}^{-2} \text{ s}^{-1}$ (high light) for 4 hours in reaction buffer containing 50 mM HEPES (pH 7.8), 330 mM sorbitol, and 2 mM MgCl_2 . Enzyme activities were measured spectrophotometrically: SOD by the nitro blue tetrazolium (NBT) reduction method [7], APX by monitoring ascorbate oxidation at 290 nm, CAT by H_2O_2 decomposition at 240 nm, GR by NADPH oxidation at 340 nm, DHAR by ascorbate formation at 265 nm, and MDHAR by NADH oxidation at 340 nm [3]. Superoxide was quantified by the SOD-inhibitable cytochrome c reduction assay, and H_2O_2 by the Amplex Red fluorometric method. Each measurement was performed on six independent chloroplast preparations. Comparisons between light treatments used paired t-tests. The ROS budget was calculated as: net ROS = total production – (SOD scavenging + APX scavenging + CAT scavenging).

scavenging).

Results

Table 1: Antioxidant Enzyme Activities in Chloroplasts Under Control and High Light

Enzyme	Control ($\mu\text{mol min}^{-1} \text{ mg}^{-1}$)	High Light ($\mu\text{mol min}^{-1} \text{ mg}^{-1}$)	Change (%)	p-value
SOD	42.3±4.8	78.6±7.2	+ 85.8	<0.001
APX	38.7±4.1	71.2±6.8	+ 84.0	<0.001
CAT	27.4±3.6	48.3±5.4	+ 76.3	<0.001
GR	31.8±3.9	54.7±5.7	+ 72.0	<0.001
DHAR	18.9±2.4	34.2±3.8	+ 81.0	<0.001
MDHAR	22.1±2.8	41.8±4.6	+ 89.1	<0.001

All six enzymes showed significant increases under high light ($p < 0.001$ for each). MDHAR showed the largest proportional increase (+ 89.1%), while CAT showed the smallest (+ 76.3%). Under high light, superoxide levels increased 2.3-fold and H_2O_2 increased 1.8-fold compared to controls despite the elevated enzyme activities.




Fig 1: Antioxidant enzyme activities in isolated spinach chloroplasts under control and high-light conditions. All enzymes were significantly upregulated under high light stress.




Fig 2: Chloroplast ROS budget under high-light conditions. Despite substantial enzymatic scavenging, a net surplus of approximately 8 relative units remained, representing the residual oxidative burden that may contribute to photoinhibition.

Discussion

The coordinated upregulation of all six antioxidant enzymes under high light reflects a system-level response rather than activation of any single pathway ^[1, 3]. The largest proportional increase in MDHAR activity is consistent with its position as a rate-limiting step in ascorbate regeneration—if MDHAR doesn't keep up, monodehydroascorbate spontaneously disproportionates to dehydroascorbate, diverting flux through the slower DHAR pathway ^[5]. The 8% net ROS surplus under high light, despite maximal enzymatic scavenging, quantifies the oxidative burden that drives photoinhibition through D1 protein damage in PSII ^[2]. This surplus may also serve a signaling function—moderate ROS levels are now recognized as retrograde signals from chloroplast to nucleus that trigger acclimation gene expression ^[8]. The relatively modest CAT response (+ 76%) may reflect the lower catalase content in chloroplasts compared to peroxisomes, where CAT is the dominant H₂O₂ scavenger ^[4]. These data provide a quantitative framework for modeling chloroplast redox homeostasis and its failure under photooxidative stress.

Conclusion

This research quantified the complete antioxidant enzyme response in isolated chloroplasts under high-light stress. All six enzymes of the ascorbate-glutathione cycle and associated pathways were upregulated by 72-89%, yet a net ROS surplus of approximately 8% persisted. This residual burden represents the threshold beyond which photoprotective mechanisms are overwhelmed, leading to photoinhibition. The data provide enzyme-level resolution for understanding chloroplast redox balance and may inform strategies to engineer improved stress tolerance in crop plants.

Acknowledgements

Funding Sources

Internal research funds from the Cauvery College of Biological Sciences supported this work.

Institutional Support

The authors thank the Cauvery College of Biological Sciences for spectrophotometry and growth chamber facilities.

Contributions Not Qualifying for Authorship

Greenhouse staff who maintained the spinach cultures are acknowledged.

References

1. Asada K. Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiology*. 2006;141(2):391-396.
2. Nishiyama Y, Allakhverdiev SI, Murata N. A new paradigm for the action of reactive oxygen species in the photoinhibition of photosystem II. *Biochimica et Biophysica Acta*. 2006;1757(7):742-749.
3. Foyer CH, Noctor G. Ascorbate and glutathione: the heart of the redox hub. *Plant Physiology*. 2011;155(1):2-18.
4. Mhamdi A, Queval G, Chaouch S, Vanderauwera S, Van Breusegem F, Noctor G. Catalase function in plants: a focus on Arabidopsis mutants as stress-mimic

models. *Journal of Experimental Botany*. 2010;61(15):4197-4220.

5. Miyake C, Asada K. Thylakoid-bound ascorbate peroxidase in spinach chloroplasts and photoreduction of its primary oxidation product monodehydroascorbate radicals in thylakoids. *Plant and Cell Physiology*. 1992;33(5):541-553.
6. Lilley RM, Fitzgerald MP, Rienits KG, Walker DA. Criteria of intactness and the photosynthetic activity of spinach chloroplast preparations. *New Phytologist*. 1975;75(1):1-10.
7. Beauchamp C, Fridovich I. Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Analytical Biochemistry*. 1971;44(1):276-287.
8. Dietz KJ, Turber N, Krieger-Liszkay A. ROS generation in peroxisomes and its role in cell signaling. *Plant Physiology*. 2016;171(3):1625-1631.