

International Journal of Statistics and Applied Mathematics

ISSN: 2456-1452
Maths 2021; 6(6): 91-96
© 2021 Stats & Maths
www.mathsjournal.com
Received: 16-09-2021
Accepted: 19-10-2021

Dr. Alok Kumar Srivastava
Department of Mathematics
Government Degree College
Barakhal, Sant Kabir Nagar,
Uttar Pradesh, India

Curvature analysis of minimal surfaces in higher-dimensional Riemannian manifolds

Alok Kumar Srivastava

DOI: <https://www.doi.org/10.22271/math.2021.v6.i6a.2303>

Abstract

The curvature theory of minimal surfaces in higher-dimensional ambient manifolds occupies a central place in modern differential geometry. This research revisits the principal curvature identities governing a minimal isometric immersion $\varphi: M^n \rightarrow N^{n+p}$ of an n -dimensional Riemannian manifold into a Riemannian manifold of codimension p , and analyses how the vanishing of the mean curvature vector constrains the geometry of the immersion. Starting from the equations of Gauss, Codazzi and Ricci for an arbitrary isometric immersion, the associated Laplacian identity of Simons for the squared norm of the second fundamental form is presented and specialised to the minimal case, producing the celebrated Simons inequality and, as a further specialisation, the pinching theorem of Chern, do Carmo and Kobayashi for minimal submanifolds of the sphere. The cases of the catenoid, helicoid and Clifford minimal hypersurfaces are examined in detail as touchstones of the general theory, and the Bernstein problem the rigidity of entire minimal graphs in Euclidean space is placed in its higher-dimensional context through the work of Bombieri, De Giorgi and Giusti, together with the stability results of Fischer-Colbrie and Schoen and of do Carmo and Peng. The main computational sections derive the relevant curvature expressions in ambient manifolds of constant curvature and extract the geometric content of the Simons inequality. The exposition closes with a discussion of the modern structure theory of Colding and Minicozzi for embedded minimal surfaces of fixed genus in three-manifolds, explaining how intrinsic curvature bounds translate into compactness and classification statements. The results gathered here provide a coherent curvature-theoretic framework for the research of minimal submanifolds in arbitrary dimension and codimension.

Keywords: Minimal submanifolds, second fundamental form, sectional curvature, Simons inequality, mean curvature vector, Chern-do Carmo-Kobayashi pinching, Gauss-Codazzi-Ricci equations, Bernstein theorem, totally geodesic immersions, higher codimension

1. Introduction

Minimal submanifolds are among the most extensively investigated objects in global differential geometry. An isometric immersion $\varphi: M^n \rightarrow N^{n+p}$ is said to be minimal when its mean curvature vector H vanishes identically, a condition that is equivalent, in the compact oriented case, to the immersion being a critical point of the volume functional among all variations that fix the boundary. This variational characterisation, going back to the work of Lagrange and Plateau on soap films, has motivated more than two centuries of research. In the classical setting of surfaces in Euclidean three-space the subject is by now a highly developed chapter of geometric analysis [4, 7, 8], while in higher dimensions and higher codimensions, and in more general ambient Riemannian manifolds, the curvature theory becomes both subtler and more rewarding.

The turning point for the higher-dimensional theory is Simons' 1968 paper on minimal varieties in Riemannian manifolds [1]. Simons derived an elegant Laplacian identity for the squared length $|\sigma|^2$ of the second fundamental form of a minimal submanifold and used it to prove integral and pointwise inequalities constraining the possible values of this quantity. A refinement of this argument due to Chern, do Carmo and Kobayashi [2] established that, for a compact minimal submanifold of the unit sphere S^{n+p} , the inequality $|\sigma|^2 \leq n/(2 - 1/p)$ forces either $|\sigma|^2 = 0$ (totally geodesic) or $|\sigma|^2 = n/(2 - 1/p)$, the latter case being realised precisely by

Corresponding Author:
Dr. Alok Kumar Srivastava
Department of Mathematics
Government Degree College
Barakhal, Sant Kabir Nagar,
Uttar Pradesh, India

Clifford minimal hypersurfaces and the Veronese surface in S^4 . This pinching theorem is the prototype of a long line of analogous results, a number of which are surveyed by Chen in [15, 17].

Alongside the curvature-pinching programme, the global theory of complete minimal surfaces in Euclidean space was developed through independent and parallel lines of research. The Bernstein problem whether every entire minimal graph in $\mathbb{R}^{n+1}$ is an affine hyperplane was settled affirmatively for $n \leq 7$ through the successive contributions of Bernstein, De Giorgi, Almgren and Simons, and shown to fail for $n \geq 8$ by Bombieri, De Giorgi and Giusti, who constructed an entire minimal graph over $\mathbb{R}^8$ that is not a hyperplane [18]. Stability and index-type theorems followed rapidly. Do Carmo and Peng showed that a complete stable minimal surface in $\mathbb{R}^3$ is a plane [17], while Fischer-Colbrie and Schoen proved the analogous result in three-manifolds of non-negative scalar curvature [10]. In the spherical setting, Lawson's rigidity theorems [3] and Reilly's extrinsic rigidity results [13] constrain the geometry of compact minimal hypersurfaces very sharply. The purpose of this research is twofold. First, the curvature identities for an arbitrary isometric immersion the equations of Gauss, Codazzi and Ricci are assembled in a form that is convenient for the subsequent analysis, and specialised to the minimal case. Second, the Simons identity is derived and its consequences examined, including the Chern-do Carmo-Kobayashi pinching theorem and its modern refinements. Along the way, explicit computations are carried out for the catenoid, the helicoid, the Clifford minimal hypersurface in S^{2m+1} , and the Veronese surface in S^4 , which serve as recurring examples throughout the theory.

2. Preliminaries and Methods

2.1 Material: Isometric immersions and the second fundamental form

All manifolds considered in this research are smooth, connected, Hausdorff and second-countable, and the metrics are of class C^∞ . Let (M^n, g) and $(N^{n+p}, \tilde{g})$ be Riemannian manifolds, and let $\varphi: M \rightarrow N$ be a smooth isometric immersion, so that $\varphi^*\tilde{g} = g$. At every point $p \in M$ the tangent space of the ambient manifold splits orthogonally,

$$T_{\varphi(p)}N = d\varphi(T_pM) \oplus T_pM^\perp,$$

defining the normal bundle $T M^\perp \rightarrow M$. Denote the Levi-Civita connections of M and N by ∇ and $\tilde{\nabla}$ respectively. The second fundamental form σ of the immersion is the symmetric bilinear mapping

$$\sigma: \Gamma(TM) \times \Gamma(TM) \rightarrow \Gamma(TM^\perp), \quad \sigma(X, Y) = \tilde{\nabla}_X Y^\perp,$$

where $(\cdot)^\perp$ denotes orthogonal projection onto $T M^\perp$. For a normal vector field ξ along M the shape operator A_ξ is the symmetric (1,1)-tensor on M characterised by the identity $g(A_\xi X, Y) = \tilde{g}(\sigma(X, Y), \xi)$. The normal connection on $T M^\perp$ is denoted by $\nabla^\perp$.

The mean curvature vector of the immersion at p is

$$H(p) = (1/n) \sum_{i=1}^n \sigma(e_i, e_i),$$

where $\{e_1, \dots, e_n\}$ is any orthonormal basis of T_pM . The immersion is called minimal if H vanishes identically, and totally geodesic if the second fundamental form σ vanishes identically. The squared norm of the second fundamental form is

$$|\sigma|^2 = \sum_{i,j=1}^n \tilde{g}(\sigma(e_i, e_j), \sigma(e_i, e_j)) = \sum_\alpha \text{tr}(A_{\xi_\alpha}^2),$$

where $\{\xi_\alpha\}$ is any orthonormal frame of $T M^\perp$.

2.2 Methods: The equations of Gauss, Codazzi and Ricci

Let R and $\tilde{R}$ denote the Riemann curvature tensors of (M, g) and $(N, \tilde{g})$ respectively, and let $R^\perp$ denote the curvature of the normal connection $\nabla^\perp$. For tangent vector fields X, Y, Z, W and normal vector fields ξ, η , the following identities hold [5, 6, 7]:

Gauss equation: $g(R(X, Y)Z, W) = \tilde{g}(\tilde{R}(X, Y)Z, W) + \tilde{g}(\sigma(X, W), \sigma(Y, Z)) - \tilde{g}(\sigma(X, Z), \sigma(Y, W))$.

Codazzi equation: $\tilde{R}(X, Y)Z^\perp = (\nabla_X^\perp \sigma)(Y, Z) - (\nabla_Y^\perp \sigma)(X, Z)$.

Ricci equation: $R^\perp(X, Y)\xi - \tilde{R}(X, Y)\xi^\perp = \sigma(X, A_\xi Y) - \sigma(Y, A_\xi X)$.

When the ambient space has constant sectional curvature c , the ambient Riemann tensor takes the model form $\tilde{R}(X, Y)Z = c(g(Y, Z)X - g(X, Z)Y)$, and the Gauss equation reduces to

$$g(R(X, Y)Z, W) = c(g(X, W)g(Y, Z) - g(X, Z)g(Y, W)) + \tilde{g}(\sigma(X, W), \sigma(Y, Z)) - \tilde{g}(\sigma(X, Z), \sigma(Y, W)).$$

Tracing this identity twice in the minimal case $H \equiv 0$ gives, at each point of M ,

$$\text{Ric}(X, X) = (n-1)c - \sum_\alpha \tilde{g}(A_{\xi_\alpha}^2 X, X), \quad \text{Scal} = n(n-1)c - |\sigma|^2.$$

The last identity makes explicit how the squared norm of the second fundamental form encodes the defect of the intrinsic scalar curvature relative to the ambient one in the minimal case.

2.3 Methods: The Simons identity

For an immersion into an ambient manifold of constant sectional curvature, Simons' computation [1] yields the following identity for the rough Laplacian of $|\sigma|^2$:

$$(1/2) \Delta |\sigma|^2 = |\nabla \sigma|^2 + nc |\sigma|^2 - \langle Q(\sigma), \sigma \rangle,$$

in the minimal case, where $Q(\sigma)$ is the quadratic expression in σ obtained by contracting pairs of shape operators. A pointwise estimate of $Q(\sigma)$ by $|\sigma|^4$ gives, for minimal submanifolds of the sphere S^{n+p} (i.e. $c = 1$),

$$(1/2) \Delta |\sigma|^2 \geq |\nabla \sigma|^2 + |\sigma|^2 (n - (2 - 1/p) |\sigma|^2).$$

Integration over a compact minimal submanifold and application of the divergence theorem then yields the pinching result of Chern, do Carmo and Kobayashi: if $0 \leq |\sigma|^2 \leq n/(2 - 1/p)$ pointwise, then either $|\sigma|^2 \equiv 0$ (and M is a totally geodesic submanifold of S^{n+p}), or $|\sigma|^2 \equiv n/(2 - 1/p)$ [1, 2].

3. Main Results: Curvature Structure of Minimal Submanifolds

3.1 Gauss-Bonnet consequence and intrinsic curvature

Combining the traced Gauss equation with the minimality condition produces a clean expression for the scalar curvature S of a minimal submanifold of an ambient space of constant curvature c :

$$S = n(n-1)c - |\sigma|^2.$$

Thus the scalar curvature of a minimal submanifold of the sphere is bounded above by $n(n-1)$, with equality precisely when M is totally geodesic. For compact minimal surfaces in S^3 , this inequality, combined with the Gauss-Bonnet theorem, gives sharp topological consequences studied in detail by Lawson^[3] and Reilly^[13].

3.2 Pinching theorem of Chern-do Carmo-Kobayashi

Theorem 1 (Chern, do Carmo & Kobayashi, 1970). Let M^n be a compact minimal submanifold of the unit sphere S^{n+p} with codimension p . If the squared norm of the second fundamental form satisfies

$$|\sigma|^2 \leq n/(2-1/p)$$

pointwise on M , then either (a) $|\sigma|^2 \equiv 0$ and M is totally geodesic (an equatorial sphere $S^n \subset S^{n+p}$), or (b) $|\sigma|^2 \equiv n/(2-1/p)$.

In case (b) the possibilities are precisely the Clifford minimal hypersurfaces $M_{k,n-k}$ when $p=1$, and the Veronese surface in S^4 when $n=2$ and $p=2$. The proof proceeds by the integrated Simons inequality reproduced in Section 2.3 and a careful analysis of the equality case using the algebraic Lemma of Chern, do Carmo and Kobayashi on pairs of symmetric matrices.

3.3 Explicit examples

Catenoid: The surface of revolution generated by rotating the catenary $y = \cosh(z)$ about the z -axis, parametrised by $\mathbf{x}(u, v)$

$= (\cosh v \cos u, \cosh v \sin u, v)$, is the unique minimal surface of revolution in $\mathbb{R}^3$ other than the plane. A direct computation gives principal curvatures $\pm \operatorname{sech}^2 v$, so that the Gaussian curvature is $K = -\operatorname{sech}^4 v$ and the squared length of the second fundamental form is $|\sigma|^2 = 2 \operatorname{sech}^4 v$. The catenoid is a complete, embedded minimal surface with finite total curvature $\int K = -4\pi$, and is characterised by this property among complete embedded minimal surfaces of revolution^[4, 8].

Helicoid: Parametrised by $\mathbf{x}(u, v) = (v \cos u, v \sin u, cu)$, the helicoid is a complete minimal surface in $\mathbb{R}^3$ with principal curvatures $\pm c/(c^2 + v^2)$ and $K = -c^2/(c^2 + v^2)^2$. By a classical theorem of Catalan, the helicoid is the only ruled minimal surface in $\mathbb{R}^3$ other than the plane^[4].

Clifford minimal hypersurfaces: For integers $k, n-k \geq 1$, the product immersion

$$S^k(\sqrt{k/n}) \times S^{n-k}(\sqrt{(n-k)/n}) \hookrightarrow S^{n+1}$$

is a minimal hypersurface of the unit sphere with constant $|\sigma|^2 = n$. These hypersurfaces, together with the equatorial sphere, realise the equality cases of Theorem 1 when $p=1$.

Veronese surface: The Veronese embedding of the real projective plane $\mathbb{RP}^2$ into S^4 via quadratic spherical harmonics produces a minimal surface of constant Gaussian curvature $1/3$ and constant $|\sigma|^2 = 4/3 = 2/(2-1/2)$, realising the equality case of Theorem 1 for $n=p=2$ ^[1, 2].

4. Results




Fig 1: The catenoid and the helicoid, two classical complete minimal surfaces in $\mathbb{R}^3$. Both surfaces have vanishing mean curvature $H \equiv 0$ at every point; they are locally isometric through a one-parameter family of minimal surfaces joining them^[4]

The curvature theory of minimal submanifolds displays a rich interplay between local invariants (the second fundamental form, the shape operator, the mean curvature vector) and global invariants (the scalar curvature, the index, the total curvature). Tables 1 and 2 collect the key analytical relations

required for this research and the associated geometric quantities on the two classical surfaces of Figure 1. Figure 2 then displays the geometric setting schematically, and Figure 3 summarises the pinching classification that emerges from the Simons inequality of Section 2.3.

Table 1: Structural curvature identities for an isometric immersion $M^n \rightarrow N^{n+p}$ (minimal specialisation in the third column)

Identity	General form	Minimal case + const. curvature c
Gauss	$R = R + \langle \sigma, \sigma \rangle$ term	$S = n(n-1)c - \sigma ^2$
Codazzi	$\tilde{R}(X, Y)Z^\perp = (\nabla^\perp \sigma)(Y, Z) - (\nabla^\perp \sigma)(X, Z)$	$(\nabla^\perp \sigma)$ totally symmetric
Ricci	$R^\perp(X, Y)\xi^\perp - \tilde{R}(X, Y)\xi^\perp = \sigma(X, A_\xi Y) - \sigma(Y, A_\xi X)$	Flat normal bundle iff $[A_\xi, A_\eta] = 0$
Mean curvature	$H = (1/n) \operatorname{tr} \sigma$	$H \equiv 0$ (minimality)
Simons (Laplacian)	$(1/2) \Delta \sigma ^2 = \nabla \sigma ^2 + \langle \sigma, \nabla^2 H + \tilde{\sigma} \text{ terms} \rangle$	$(1/2) \Delta \sigma ^2 = \nabla \sigma ^2 + nc \sigma ^2 - \langle Q(\sigma), \sigma \rangle$
Pinching	Pointwise inequality on $ \sigma ^2$	$ \sigma ^2 \in \{0, n/(2-1/p)\}$ (Thm 1)

The third column of Table 1 highlights how the minimality condition decouples the Simons identity into a form amenable to integration by parts on compact manifolds, which is the

starting point of every pinching theorem since Simons ^[1] and Chern-do Carmo-Kobayashi ^[2].

Table 2: Curvature data for four fundamental examples of minimal submanifolds

Submanifold	Ambient	Dim / codim	$ \sigma ^2$	Compact?
Catenoid	$\mathbb{R}^3$	$n=2, p=1$	$2 \operatorname{sech}^4 v$	No
Helicoid	$\mathbb{R}^3$	$n=2, p=1$	$2c^2/(c^2+v^2)^2$	No
Clifford $S^k \times S^{n-k}$	S^{n+1}	$\dim n, p=1$	n (constant)	Yes
Veronese $\mathbb{R}P^2$	S^4	$n=2, p=2$	$4/3$ (constant)	Yes

The last two rows are the equality cases of Theorem 1. Note that for the Clifford hypersurface $n/(2 - 1/p) = n$ since $p = 1$,

and for the Veronese surface $n/(2 - 1/p) = 2/(3/2) = 4/3$, consistently with the values in the $|\sigma|^2$ column.




Fig 2: Schematic of the immersion $M^n \hookrightarrow N^{n+p}$ at a point p . The tangent vector X is horizontal with respect to the immersion; ξ is a normal vector; the mean curvature vector vanishes because the immersion is minimal. The second fundamental form σ and the Gauss equation are recalled in the boxed insets




Fig 3: Pinching classification of compact minimal submanifolds of the sphere S^{n+p} in the Chern-do Carmo-Kobayashi sense ^[1, 2]. The critical value of $|\sigma|^2$ is $n/(2 - 1/p)$, achieved precisely by the Clifford minimal hypersurfaces (when $p = 1$) and by the Veronese surface (when $n = p = 2$)

Comprehensive interpretation

The combined picture provided by Tables 1 and 2 and Figures 1-3 can be read as follows. Table 1 assembles the curvature identities that any isometric immersion must satisfy, together with their minimal-case specialisations in an ambient manifold of constant sectional curvature. These identities are

the algebraic backbone of every quantitative result about minimal submanifolds: The Gauss equation relates intrinsic and extrinsic curvature; the Codazzi equation controls the differentiability structure of the second fundamental form; and the Ricci equation measures the failure of the normal connection to be flat. The Simons identity, stated in the last

two rows, packages these into a single elliptic inequality for the scalar quantity $|\sigma|^2$ whose global consequences are extracted via integration by parts.

Table 2 shows how the abstract pinching threshold $n/(2 - 1/p)$ translates into concrete numerical values on the fundamental examples. For the Clifford hypersurface the threshold equals n , and the squared length of the second fundamental form is exactly n . For the Veronese surface the threshold is $4/3$, and the surface realises this value as well. The non-compact examples the catenoid and the helicoid have non-constant $|\sigma|^2$ and do not fit directly into the pinching paradigm of Theorem 1 (which requires compactness), but they do illustrate how minimality constrains the pointwise geometry of a surface in Euclidean space.

Figures 1 and 2 visualise the local geometric setting: the catenoid and helicoid display the classical surface-theoretic picture, and the schematic in Figure 2 clarifies the tangent/normal decomposition on which the Gauss, Codazzi and Ricci equations are built. Figure 3 then distils the upshot of the Simons-Chern-do Carmo-Kobayashi chain into a one-dimensional picture: on the real axis of $|\sigma|^2$ values, the gap between 0 and $n/(2 - 1/p)$ is a forbidden zone for compact minimal submanifolds of the sphere.

5. Discussion

The results surveyed here have been refined in several directions since their original appearance. Chen introduced his now widely used δ -invariants and the associated fundamental inequalities ^[5, 15], which provide sharper pointwise bounds on mean and scalar curvatures in terms of sectional curvatures of the ambient space. The relationship between the mean curvature vector, the shape operator and the ambient curvature in space forms was made explicit by Chen in ^[12] and has formed the basis of a sizeable body of subsequent work. The stability and index theory of minimal submanifolds, initiated by Simons ^[1], developed by Spruck ^[9] and by Fischer-Colbrie and Schoen ^[10], and refined by do Carmo and Peng ^[17], gives a second, complementary perspective, placing the second variation of the volume functional at the centre of the analysis.

On the global side, the Bernstein problem the rigidity of entire minimal graphs over $\mathbb{R}^n$ in $\mathbb{R}^{n+1}$ was resolved in its essential aspects over a sequence of papers between 1915 and 1969. Bernstein settled the classical case $n = 2$; De Giorgi handled $n = 3$; Almgren treated $n = 4$; and Simons ^[1], in the same 1968 paper that introduced the Laplacian identity discussed above, settled the cases $n = 5, 6$ and 7 . The sharpness of these results was then established in 1969 by Bombieri, De Giorgi and Giusti ^[18], who constructed an entire minimal graph over $\mathbb{R}^8$ that is not a hyperplane, thereby showing that Bernstein rigidity fails from dimension 8 onward. The counterexample relies on a careful analysis of the Simons cone $\{x \in \mathbb{R}^8: x_1^2 + \dots + x_4^2 = x_5^2 + \dots + x_8^2\}$ and introduced the use of minimal cones in high dimensions as a central analytical tool.

In the ambient class of three-manifolds of non-negative scalar curvature, Fischer-Colbrie and Schoen's rigidity theorem ^[10] states that a complete stable minimal surface is either a plane or a flat totally geodesic torus, and in the latter case the ambient manifold must split isometrically as a product near the surface. This is a prototype of the interaction between ambient curvature, intrinsic curvature, and extrinsic stability, which has been developed further by Meeks, Pérez and their collaborators ^[8, 16]. Colding and Minicozzi's four-paper series on the space of embedded minimal surfaces of fixed genus in a three-manifold, of which ^[16] is the final instalment,

establishes powerful compactness and structure theorems under intrinsic curvature bounds a development closely tied to the Poincaré and geometrisation programmes.

Several reasons support the continued interest in the extrinsic curvature analysis of minimal submanifolds. First, minimality is a natural relaxation of the totally geodesic condition and allows the construction of rich families of examples the Clifford and Veronese examples are only the simplest. Second, the curvature identities reproduced in Section 2 remain the analytical engine behind most pointwise and integral estimates. And third, higher-dimensional and higher-codimensional theorems often disclose phenomena invisible from the classical surface-theoretic viewpoint, as the Bombieri-De Giorgi-Giusti counterexample and the Chern-do Carmo-Kobayashi pinching dichotomy both demonstrate. The framework presented in this research is intended to serve as a compact reference from which these more specialised directions can be approached.

6. Conclusion

The main contribution of this research is a consolidated curvature-theoretic treatment of minimal isometric immersions into higher-dimensional Riemannian manifolds, organised so as to expose the logical links between local curvature identities and global classification statements. The development proceeds from the equations of Gauss, Codazzi and Ricci, through the Simons Laplacian identity for the squared norm of the second fundamental form, to the Chern-do Carmo-Kobayashi pinching theorem and its sharp realisations on Clifford and Veronese minimal submanifolds of the sphere. The logical chain is short and transparent: traced Gauss gives the scalar-curvature identity, Simons gives the elliptic inequality for $|\sigma|^2$, and integration on a compact minimal submanifold then forces the dichotomy of Theorem 1.

The supporting evidence assembled here draws from six decades of sustained research on the topic. The Simons identity ^[1], the Chern-do Carmo-Kobayashi theorem ^[2], Lawson's and Reilly's rigidity results ^[3, 13], the Bernstein-type results in arbitrary dimension settled by Bombieri, De Giorgi and Giusti ^[18], the stability and structure theory of Fischer-Colbrie and Schoen ^[10], of do Carmo and Peng ^[17], and of Colding and Minicozzi ^[7, 16], and the δ -invariant framework of Chen ^[5, 12, 15] collectively demonstrate how fertile and interconnected the curvature analysis of minimal submanifolds has become.

For the field as a whole, the significance of the results collected here is that they provide a concise entry point into both the pinching and the rigidity programmes. Several open questions remain vigorous. Sharp versions of the Chern-do Carmo-Kobayashi theorem in codimension greater than two are still actively investigated. The classification of complete embedded minimal surfaces of finite total curvature in arbitrary three-manifolds is not complete. And the index theory of unstable minimal submanifolds in higher codimension, especially in ambient manifolds without constant curvature, remains largely open. The framework assembled in this research is intended to make these directions more accessible to readers approaching the subject for the first time.

Acknowledgements

The author is sincerely thankful to the concerned authorities for their kind support and encouragement throughout the preparation of this manuscript.

References

1. Simons J. Minimal varieties in Riemannian manifolds. *Annals of Mathematics*. 1968;88(1):62-105.
2. Chern SS, do Carmo M, Kobayashi S. Minimal submanifolds of a sphere with second fundamental form of constant length. In: Browder FE, editor. *Functional Analysis and Related Fields*. Berlin: Springer-Verlag; 1970. p. 59-75.
3. Lawson HB Jr. Local rigidity theorems for minimal hypersurfaces. *Annals of Mathematics*. 1969;89(1):187-197.
4. Osserman R. A survey of *minimal surfaces*. 2nd ed. New York: Dover Publications; 1986.
5. Chen B-Y. Some pinching and classification theorems for *minimal submanifolds*. *Archiv der Mathematik*. 1993;60(6):568-578.
6. do Carmo MP. *Riemannian geometry*. Boston: Birkhäuser; 1992.
7. Colding TH, Minicozzi WP II. A course in *minimal surfaces*. Graduate Studies in Mathematics; vol. 121. Providence: American Mathematical Society; 2011.
8. Meeks WH III, Pérez J. A survey on classical *minimal surface theory*. University Lecture Series; vol. 60. Providence: American Mathematical Society; 2012.
9. Spruck J. Remarks on the stability of *minimal submanifolds* of $\mathbb{R}^n$. *Mathematische Zeitschrift*. 1975;144(2):169-174.
10. Fischer-Colbrie D, Schoen R. The structure of complete stable *minimal surfaces* in 3-manifolds of non-negative scalar curvature. *Communications on Pure and Applied Mathematics*. 1980;33(2):199-211.
11. Calabi E. *Minimal immersions* of surfaces in Euclidean spheres. *Journal of Differential Geometry*. 1967;1(1-2):111-125.
12. Chen B-Y. Mean curvature and shape operator of isometric immersions in real-space-forms. *Glasgow Mathematical Journal*. 1996;38(1):87-97.
13. Reilly RC. Extrinsic rigidity theorems for compact submanifolds of the sphere. *Journal of Differential Geometry*. 1970;4(4):487-497.
14. Yau ST. Submanifolds with constant mean curvature I. *American Journal of Mathematics*. 1974;96(2):346-366.
15. Chen B-Y. *Pseudo-Riemannian geometry, δ -invariants and applications*. Singapore: World Scientific; 2011.
16. Colding TH, Minicozzi WP II. The space of embedded *minimal surfaces* of fixed genus in a 3-manifold IV; locally simply connected. *Annals of Mathematics*. 2004;160(2):573-615.
17. do Carmo MP, Peng CK. Stable complete *minimal surfaces* in $\mathbb{R}^3$ are planes. *Bulletin of the American Mathematical Society (New Series)*. 1979;1(6):903-906.
18. Bombieri E, De Giorgi E, Giusti E. Minimal cones and the Bernstein problem. *Inventiones Mathematicae*. 1969;7(3):243-268.
19. Montiel S. An integral inequality for compact spacelike hypersurfaces in de Sitter space and applications to the case of constant mean curvature. *Indiana University Mathematics Journal*. 1988;37(4):909-917.