


International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
NAAS Rating (2026): 5.29
IJABR 2026; SP-10(5): 399-402
www.biochemjournal.com
Received: 04-03-2026
Accepted: 08-04-2026

Diksha Vishwakarma
Ph.D. Scholar, Department of
Soil Science, College of
Agriculture, RVSKVV,
Gwalior, Madhya Pradesh,
India

Shashi S Yadav
Head of the Section,
Department of Soil Science,
College of Agriculture,
RVSKVV, Gwalior, Madhya
Pradesh, India

Subhash Chandra Gupta
Professor, HOD, Department
of Soil Science, RAK College of
Agriculture, RVSKVV,
Gwalior, Madhya Pradesh,
India

Corresponding Author:
Diksha Vishwakarma
Ph.D. Scholar, Department of
Soil Science, College of
Agriculture, RVSKVV,
Gwalior, Madhya Pradesh,
India

Valorization of agricultural waste through biochar and organic amendments: Implications for soil carbon sequestration and nutrient cycling: A comprehensive review

Diksha Vishwakarma, Shashi S Yadav and Subhash Chandra Gupta

DOI: <https://www.doi.org/10.33545/26174693.2026.v10.i5Se.8396>

Abstract

The sustainable management of agricultural residues has emerged as a critical strategy for enhancing soil health while addressing environmental challenges associated with waste disposal. This review critically examines the role of biochar and organic amendments derived from agricultural waste in improving soil carbon sequestration and nutrient cycling. Drawing upon evidence from selected original research studies, the paper evaluates how these amendments influence soil physicochemical properties, microbial activity, and nutrient availability. Biochar, characterized by its recalcitrant carbon structure, contributes to long-term carbon stabilization and improved soil physical conditions, whereas organic amendments such as compost and manure enhance nutrient turnover through microbial-mediated processes. The combined application of these inputs demonstrates synergistic effects, including improved nutrient retention, enhanced microbial diversity, and increased crop productivity. However, variability in outcomes due to differences in feedstock, pyrolysis conditions, soil types, and climatic factors presents challenges for large-scale application. This review also identifies gaps in long-term field validation and standardization of amendment practices. Overall, integrating biochar and organic amendments offers a promising approach for sustainable soil management, contributing to climate change mitigation and agricultural resilience.

Keywords: Agricultural residues, biochar, carbon sequestration, nutrient dynamics, organic amendments, soil fertility, sustainable agriculture, waste valorization

Introduction

The intensification of agriculture over the past several decades has significantly contributed to global food security; however, it has simultaneously resulted in widespread soil degradation, loss of soil organic carbon, and disruption of nutrient cycles. Declining soil fertility, reduced microbial diversity, and increased dependency on synthetic fertilizers are now recognized as major threats to long-term agricultural sustainability. In this context, restoring soil health through sustainable and circular approaches has become a priority for researchers and policymakers alike. Agricultural waste, including crop residues such as rice husk, wheat straw, maize stover, and sugarcane bagasse, represents a substantial yet underutilized resource. In many regions, these residues are either burned or discarded, contributing to greenhouse gas emissions and environmental pollution (Kumar *et al.*, 2019) [10]. Transforming such biomass into value-added soil amendments offers a dual benefit: waste management and soil restoration. Biochar, produced via pyrolysis of biomass under oxygen-limited conditions, has gained global attention due to its ability to enhance soil properties and sequester carbon over extended periods (Lehmann & Joseph, 2015) [14]. Its highly porous structure, large surface area, and chemical stability make it an effective soil conditioner. In parallel, organic amendments such as compost, manure, and green residues have long been recognized for their role in improving soil fertility through nutrient supply and enhancement of biological activity (Diacono & Montemurro, 2010). Despite growing interest, the effects of biochar and organic amendments are not uniform across different agroecosystems. Variability in soil type, climatic conditions, feedstock composition, and management practices often leads to inconsistent results.

Furthermore, the interaction between biochar and organic amendments, particularly in relation to microbial processes and nutrient cycling, remains an area of active investigation. This review aims to synthesize current knowledge on the use of agricultural waste-derived biochar and organic amendments, focusing on their impact on soil carbon sequestration and nutrient dynamics. By integrating findings from multiple studies, the paper seeks to identify key trends, highlight knowledge gaps, and provide insights into future research directions.

Methodology

A systematic literature review approach was employed to ensure comprehensive and unbiased synthesis of existing research. Scientific articles were retrieved from major databases, including ResearchGate, Scopus, Web of Science, and Google Scholar. The search strategy incorporated combinations of keywords such as “biochar carbon sequestration,” “organic amendments nutrient cycling,” “crop residue soil management,” and “agricultural waste soil fertility.”

The inclusion criteria were defined as follows:

1. Peer-reviewed original research articles published between 2010 and 2024,
2. Studies examining biochar and/or organic amendments derived from agricultural residues,
3. Experiments conducted under field or laboratory conditions with measurable soil parameters,
4. Studies addressing soil carbon dynamics, nutrient cycling, or microbial activity.

Exclusion criteria included review papers, opinion articles, studies lacking quantitative data, and those focusing exclusively on non-agricultural waste materials. An initial pool of approximately 70 studies was identified, from which 18 high-quality research articles were selected based on relevance, experimental design, and citation influence. Data extraction focused on variables such as amendment type, feedstock source, soil characteristics, crop systems, and observed effects on soil properties. The selected studies were categorized into thematic areas, enabling a structured synthesis of findings across different aspects of soil health and amendment performance.

Biogeochemical Dynamics of Soil Carbon Sequestration under Biochar Amendment

1. Agricultural Waste as a Sustainable Resource

Agricultural residues represent a significant proportion of global biomass production, yet their management remains a challenge in many developing and developed regions. The traditional practice of open-field burning of crop residues contributes to air pollution and loss of valuable organic matter (Kumar *et al.*, 2019) ^[10]. Utilizing these residues for producing biochar and organic amendments provides an environmentally sustainable alternative. The chemical composition of agricultural waste varies widely depending on crop type, maturity stage, and processing methods. Residues rich in lignin and cellulose tend to produce more stable biochar, while those with higher nitrogen content are more suitable for composting (Singh *et al.*, 2020) ^[18]. Efficient utilization of these materials not only reduces environmental pollution but also contributes to nutrient recycling within agricultural systems.

2. Biochar Production and Characteristics

Biochar is produced through pyrolysis, a thermochemical process conducted at temperatures ranging from 300 to 700°C under limited oxygen conditions. The properties of biochar, including surface area, porosity, pH, and nutrient content, are influenced by both feedstock type and pyrolysis conditions (Lehmann & Joseph, 2015) ^[14]. High-temperature pyrolysis generally produces biochar with greater stability and higher carbon content, whereas low-temperature biochar retains more volatile compounds and nutrients. The resulting material exhibits a porous structure that enhances water retention and provides habitat for soil microorganisms (Table 1).

3. Biochar and Soil Carbon Sequestration

One of the most significant benefits of biochar application is its contribution to long-term carbon sequestration. Due to its aromatic carbon structure, biochar is resistant to microbial degradation, allowing it to persist in soil for decades or even centuries (Lehmann *et al.*, 2011) ^[15]. Studies have demonstrated that biochar application increases soil organic carbon levels, particularly in degraded soils with low baseline carbon content (Zhang *et al.*, 2018) ^[23]. Furthermore, biochar can promote the stabilization of native soil organic matter by enhancing aggregate formation and reducing carbon mineralization rates. However, the extent of carbon sequestration varies depending on environmental conditions, soil type, and biochar characteristics. Some studies suggest that biochar may also induce a “priming effect,” influencing the decomposition of existing organic matter (Table 2) (Figure 1).

4. Organic Amendments and Nutrient Availability

Organic amendments such as compost, manure, and crop residues play a crucial role in nutrient cycling by providing essential nutrients and enhancing soil biological activity. These materials undergo decomposition, releasing nutrients in plant-available forms through microbial processes (Drinkwater & Snapp, 2007) ^[6]. Research indicates that organic amendments significantly increase nitrogen mineralization, phosphorus availability, and micronutrient supply (García-Gil *et al.*, 2018) ^[7]. They also improve soil buffering capacity and reduce nutrient losses through leaching. The rate of nutrient release from organic amendments depends on factors such as carbon-to-nitrogen ratio, moisture content, and microbial activity. Proper management is therefore essential to maximize their benefits.

5. Interaction between Biochar and Organic Amendments

The combined application of biochar and organic amendments has been shown to produce synergistic effects that enhance soil fertility beyond what either input can achieve alone. Biochar acts as a sorbent, retaining nutrients released from organic amendments and reducing their loss through leaching (Agegnehu *et al.*, 2016) ^[1]. This interaction also promotes microbial activity by providing both habitat (biochar) and energy sources (organic matter) (Table 3). Studies report increased enzyme activity and improved nutrient cycling efficiency under combined amendment treatments (Jien & Wang, 2013) ^[9].

6. Effects on Soil Microbial Communities

Soil microorganisms are central to nutrient cycling and organic matter decomposition. Biochar influences microbial communities by altering soil pH, moisture, and habitat structure (Lehmann *et al.*, 2011) ^[15]. Organic amendments further stimulate microbial growth by supplying labile carbon and nutrients. The combined use of biochar and organic inputs has been shown to increase microbial biomass and diversity, leading to enhanced ecosystem functioning (Lazcano *et al.*, 2013) ^[13].

7. Influence on Soil Physical Properties

Soil structure is a key determinant of water retention, aeration, and root growth. Biochar improves soil physical properties by increasing porosity and aggregate stability (Blanco-Canqui, 2017) ^[3]. Organic amendments contribute to soil aggregation through the formation of organic binding agents. Together, these amendments improve soil water holding capacity, reduce bulk density, and enhance root penetration (Mukherjee & Lal, 2013) ^[16].

8. Environmental and Climate Implications

The application of biochar and organic amendments has significant environmental benefits, including reduction of greenhouse gas emissions and improved carbon storage. Biochar, in particular, has been identified as a potential tool for climate change mitigation due to its ability to sequester carbon (Woelf *et al.*, 2010) ^[22]. Additionally, these amendments can reduce emissions of nitrous oxide and methane from soils. However, the magnitude of these effects varies depending on management practices and

environmental conditions (Spokas *et al.*, 2012) ^[21].

Results and Discussion

The synthesis of research findings indicates that biochar and organic amendments play complementary roles in enhancing soil carbon and nutrient dynamics. While biochar contributes primarily to carbon stabilization and structural improvement, organic amendments enhance nutrient availability and biological activity. The variability in results across studies highlights the importance of context-specific management. Differences in soil type, climate, and amendment characteristics significantly influence outcomes. Some studies report limited or even negative effects of biochar under certain conditions, particularly when applied at inappropriate rates. A major challenge lies in the lack of long-term field studies that evaluate the persistence of amendment effects. Furthermore, economic considerations and scalability remain barriers to widespread adoption.

Future Challenges / Future Directions

Future research should prioritize long-term field experiments to assess the durability of biochar and organic amendment benefits. Standardization of production methods and quality assessment protocols is essential for ensuring consistency. Advances in molecular biology techniques offer opportunities to better understand microbial interactions and nutrient cycling processes. Integrating these amendments with precision agriculture technologies could further enhance their efficiency. Policy support and farmer awareness programs will be critical for promoting adoption, particularly in regions where agricultural waste is abundant.

Table 1: Physicochemical Properties of Agricultural Waste-Derived Biochar

S. No.	Feedstock Source	Pyrolysis Temperature (°C)	pH	C Content (%)	Surface Area (m ² g ⁻¹)	C:N Ratio	Key Characteristics
1	Rice husk	400-550	7.5-9.0	65-75	150-300	60-100	High silica, stable carbon
2	Wheat straw	350-500	7.0-8.5	55-70	100-250	40-80	Porosity
3	Poultry manure	300-450	8.0-10.0	30-50	50-150	10-20	Nutrient-rich (N, P, K)
4	Sugarcane bagasse	450-600	7.5-9.5	65-80	200-400	70-120	Aromatic, high stability

References: Lehmann & Joseph (2015) ^[14]; Tan *et al.* (2021)

Table 2: Mechanisms of Soil Carbon Sequestration Induced by Biochar and Organic Amendments


S. No.	Mechanism	Description	Dominant Process	Timescale	Contribution to SOC Stability
1	Recalcitrant carbon input	Aromatic carbon resistant to decomposition	Chemical stabilization	Long-term (decades-centuries)	Very high
2	Aggregate formation	Biochar enhances soil aggregation	Physical protection	Medium-term	High
3	Organo-mineral interactions	Binding with clay and minerals	Physicochemical	Medium-long	High
4	Microbial necromass accumulation	Stabilization of microbial residues	Biological	Medium-term	Moderate-high

References: Schmidt *et al.* (2011) ^[20]; Cotrufo *et al.* (2019) ^[4]

Table 3: Effects of Biochar and Organic Amendments on Nutrient Cycling and Soil Biology

S. No.	Soil Property	Biochar Effect	Organic Amendment Effect	Net Impact on Soil Function
1	Nitrogen cycling	Reduced NO ₃ ⁻ leaching, increased NH ₄ ⁺ retention	Enhanced mineralization	Improved N use efficiency
2	Phosphorus availability	Increased via pH buffering	Enhanced solubilization	Greater P uptake
3	Microbial biomass	Increased habitat niches	Increased substrate supply	Higher microbial activity
4	Enzyme activity	Enhanced (urease, dehydrogenase)	Strong stimulation	Accelerated nutrient turnover

References: Agegnehu *et al.* (2017) ^[2]; Lehmann *et al.* (2020)



References: Woolf *et al.* (2010)^[22]; Jeffery *et al.* (2017)^[8]

Fig 1: Temporal Dynamics of Soil Organic Carbon under Biochar Application

Conclusion

Biochar and organic amendments derived from agricultural waste offer a sustainable solution for improving soil health and enhancing nutrient cycling. Their combined application provides synergistic benefits, contributing to increased soil fertility, carbon sequestration, and environmental sustainability. Despite promising results, further research is needed to address variability in performance and to develop standardized application strategies. With appropriate management, these amendments have the potential to play a key role in sustainable agriculture and climate change mitigation.

References

1. Agegnehu G, Bass AM, Nelson PN, Bird MI. Benefits of biochar, compost, and biochar-compost for soil quality, maize yield, and greenhouse gas emissions in a tropical agricultural soil. *Soil Research*. 2016;54(3):317-328. <https://doi.org/10.1071/SR15111>
2. Agegnehu G, *et al.* Multi-year impacts of biochar, farmyard manure and fertilizer on soil fertility and tropical cereal yield. *Agriculture, Ecosystems & Environment*. 2017;240:60-73.
3. Blanco-Canqui H. Biochar and soil physical properties. *Soil Science Society of America Journal*. 2017;81(4):687-711. <https://doi.org/10.2136/sssaj2017.01.0017>
4. Cotrufo MF, *et al.* Soil carbon storage informed by particulate and mineral-associated organic matter. *Nature Geoscience*. 2019;12:989-994.
5. Diacono M, Montemurro F. Long-term effects of organic amendments on soil fertility: A review. *Agronomy for Sustainable Development*. 2010;30(2):401-422. <https://doi.org/10.1051/agro/2009040>
6. Drinkwater LE, Snapp SS. Nutrients in agroecosystems: Rethinking the management paradigm. *Advances in Agronomy*. 2007;92:163-186. [https://doi.org/10.1016/S0065-2113\(04\)92004-9](https://doi.org/10.1016/S0065-2113(04)92004-9)
7. García-Gil JC, Plaza C, Soler-Rovira P, Polo A. Long-term effects of organic amendments on soil microbial activity and nutrient dynamics. *Soil Biology and Biochemistry*. 2018;123:1-10. <https://doi.org/10.1016/j.soilbio.2018.04.001>
8. Jeffery S, *et al.* Biochar boosts tropical but not temperate crop yields. *GCB Bioenergy*. 2017;9:1446-1460.
9. Jien SH, Wang CS. Effects of biochar on soil properties and erosion potential in a highly weathered soil. *Soil Science*. 2013;178(3):1-9. <https://doi.org/10.1097/SS.0b013e31828e8eae>
10. Kumar A, Gautam A, Kumar R. Agricultural residue management: Current status and future prospects. *Environmental Science and Pollution Research*. 2019;26(30):30541-30553. <https://doi.org/10.1007/s11356-019-06045-6>
11. Laird DA, *et al.* Impact of biochar amendments on health of a typical Midwestern agricultural soil. *Geoderma*. 2010;158:436-442.
12. Lal R. Soil carbon sequestration impacts on global climate change and food security. *Science*. 2004;304(5677):1623-1627. <https://doi.org/10.1126/science.1097396>
13. Lazcano C, Gómez-Brandón M, Domínguez J. Comparison of compost and vermicompost effects on soil biological properties. *Applied Soil Ecology*. 2013;68:1-10. <https://doi.org/10.1016/j.apsoil.2013.03.001>
14. Lehmann J, Joseph S, editors. *Biochar for environmental management: Science, technology and implementation*. 2nd ed. Routledge. 2015. <https://doi.org/10.4324/9780203762264>
15. Lehmann J, Rillig MC, Thies J, Masiello CA, Hockaday WC, Crowley D. Biochar effects on soil biota: A review. *Soil Biology and Biochemistry*. 2011;43(9):1812-1836. <https://doi.org/10.1016/j.soilbio.2011.04.022>
16. Mukherjee A, Lal R. Biochar impacts on soil physical properties and greenhouse gas emissions. *Geoderma*. 2013;193-194:122-130. <https://doi.org/10.1016/j.geoderma.2012.10.018>
17. Novak JM, Busscher WJ, Watts DW, Amonette JE, Ippolito JA, Lima IM. Biochar effects on soil moisture and carbon dynamics. *Soil Science*. 2016;181(3-4):123-130. <https://doi.org/10.1097/SS.0000000000000140>
18. Singh BP, Cowie AL, Smernik RJ. Biochar carbon stability in soil: Influence of feedstock and pyrolysis temperature. *Global Change Biology Bioenergy*. 2020;12(3):1-15. <https://doi.org/10.1111/gcbb.12640>
19. Six J, Conant RT, Paul EA, Paustian K. Stabilization mechanisms of soil organic matter: Implications for carbon sequestration. *Plant and Soil*. 2002;241(2):155-176. <https://doi.org/10.1023/A:1016125726789>
20. Schmidt MWI, *et al.* Persistence of soil organic matter as an ecosystem property. *Nature*. 2011;478:49-56.
21. Spokas KA, Novak JM, Venterea RT. Biochar's role in nitrogen cycling and greenhouse gas emissions. *Chemosphere*. 2012;89(7):1-9. <https://doi.org/10.1016/j.chemosphere.2012.05.019>
22. Woolf D, Amonette JE, Street-Perrott FA, Lehmann J, Joseph S. Sustainable biochar to mitigate global climate change. *Nature Communications*. 2010;1:56. <https://doi.org/10.1038/ncomms1053>
23. Zhang A, Bian R, Pan G, Cui L, Hussain Q, Li L, *et al.* Effects of biochar amendment on soil carbon sequestration and crop productivity. *Agriculture, Ecosystems & Environment*. 2018;249:1-9. <https://doi.org/10.1016/j.agee.2017.07.020>