

International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
NAAS Rating (2026): 5.29
IJABR 2026; SP-10(5): 1104-1116
www.biochemjournal.com
Received: 18-02-2026
Accepted: 21-03-2026

Pragya Singh Deo
Ph.D. Scholar, Department of Fruit Science, Mahatma Gandhi University of Horticulture and Forestry, (MGVUU), Chhattisgarh, India

Nandini Sarthi
Ph.D. Scholar, Department of Vegetable Science, Mahatma Gandhi University of Horticulture and Forestry, (MGVUU), Chhattisgarh, India

Shristi Yadav
M.Sc. Scholar, Department of Post-harvest Management, Mahatma Gandhi University of Horticulture and Forestry, (MGVUU), Chhattisgarh, India

Suhel Eraqui
Ph.D. Research Scholar, Horticulture (Vegetable Science), Pt.KLS College of Horticulture and Research Station, Rajnandgaon (MGUVV,Durg), Chhattisgarh, India

Divya Patel
M.Sc. Scholar, Department of Genetics and Plant Breeding, IGKV Raipur, Chhattisgarh, India

Rana Abhilash Singh
Chairman, Department of Plant Breeding Society for Advancement of Science & Social Environment (SASSE), Lucknow, Uttar Pradesh, India

Pradip Pandey
Ph.D. Scholar Department of Vegetable Improvement Integral University IIASST, Lucknow, Uttar Pradesh, India

Dr. Aditya Singh
Contractual Teacher, Department of Agricultural Economics, College of Agriculture, RVSKVV, Indore, Madhya Pradesh, India

Saroj Jangade
Ph.D. Scholar, Department of Vegetable Science, Pt. Parshuram Dwivedi College, Jagmmanpur, Jalaun, Uttar Pradesh, India

Corresponding Author:
Pragya Singh Deo
Ph.D. Scholar, Department of Fruit Science, Mahatma Gandhi University of Horticulture and Forestry, (MGVUU), Chhattisgarh, India

Expanding frontiers in vegetable cultivation: From traditional practices to modern greenhouse technologies

Pragya Singh Deo, Nandini Sarthi, Shristi Yadav, Suhel Eraqui, Divya Patel, Rana Abhilash Singh, Pradip Pandey, Aditya Singh and Saroj Jangade

DOI: <https://www.doi.org/10.33545/26174693.2026.v10.i5Sm.8518>

Abstract

The cultivation of vegetables is a cornerstone of global agriculture, evolving dramatically through technological innovation and changing societal needs. This review maps the trajectory of vegetable farming from its early roots in simple field agriculture to advanced controlled-environment systems. We begin by recounting historical practices and key figures who shaped horticulture (from early domesticators to modern breeders). Traditional farming methods (seed saving, hand sowing, manure fertilization, etc.) are described alongside their cultural contexts. We then outline transition phases of intensification in the 20th century (chemical fertilizers, irrigation and protected culture). The core sections cover modern greenhouse-based methods: soilless culture (hydroponics, aeroponics), vertical farming and comprehensive greenhouses equipped with climate controls, automated sensing and AI-driven optimization. Each technology is explained along with its advantages and limitations. Crop management topics are discussed in detail, including varietal selection, propagation techniques, nutrient regimes and integrated pest and disease control. The analysis also addresses economic aspects (investment, yield, market factors) and environmental considerations (resource use, emissions, waste). Policy frameworks and socioeconomic barriers to adoption (regulatory, financial and educational) are examined. Finally, we look forward to emerging trends and research needs, emphasizing the promise of digital technologies, breeding innovations and sustainable practices in shaping the future of vegetable production.

Keywords: Vegetable cultivation, greenhouse, hydroponics, aeroponics, vertical farming, controlled-environment agriculture, crop management, integrated pest management, sustainability, automation, AI, agricultural economics

Introduction

Vegetable cultivation has been central to human societies since the dawn of agriculture. At its core, growing vegetables involves nurturing plants from seed to harvest, but this seemingly simple task has undergone profound transformation. Historically, farmers relied on empirical knowledge and manual labor, shaped by local climate and traditions [88]. In recent decades, advances in plant science and engineering have radically altered how we grow vegetables, especially in controlled environments. The evolution spans from hand tended plots and organic manures to automated, sensor equipped greenhouses and vertical farms. Modern techniques emphasize precision and efficiency optimizing light, water, nutrients and protection to achieve higher yields and quality [18].

This review aims to capture the breadth of this transformation. We begin by tracing the history of vegetable cultivation, highlighting milestones and contributors from ancient farmers to pioneering scientists. Next, we describe traditional farming practices how crops were grown using age-old methods followed by a discussion of the transition phases that introduced mechanization and early protected cultivation. The heart of the paper examines modern greenhouse systems: soilless technologies like hydroponics and aeroponics, the rise of vertical farming and the sophisticated climate management in contemporary greenhouses. We then delve into crop management, covering plant breeding (varieties), propagation, nutrition and pest/disease control within these systems [36].

We further analyze the economics and scalability of high-tech farming versus conventional methods, along with sustainability and environmental impacts (water and energy use, carbon footprints). We address policy and adoption barriers factors that inhibit or slow the spread of advanced cultivation. Finally, we suggest future directions and research gaps, considering innovations like AI, robotics and advanced breeding. Through this detailed exploration, we illuminate how vegetable farming has advanced and what lies ahead for feeding a growing population sustainably ^[54].

History of Vegetable Cultivation

- The history of vegetable cultivation is a journey from simple garden plots to technological marvels. Early humans began domesticating plants around 8000-5000 BCE, initiating the Neolithic Revolution. The first farmers selected wild edible plants and gradually improved them through careful selection selecting seeds from the most vigorous individuals year after year. These prehistoric cultivators had an intuitive grasp of heredity, even without genetic science and effectively began the process of plant breeding ^[71].
- Over millennia, various cultures added innovations. Ancient Mesopotamians and Egyptians developed irrigation systems (canals and basins) to water fields and gardens. The Hanging Gardens of Babylon (circa 600 BCE) are often cited (though semi-mythical) as an early example of engineered cultivation, possibly resembling a hydroponic terrace for vegetables and fruits. In the Roman Empire, Emperor Tiberius (1st century AD) famously demanded cucumbers year-round. Roman gardeners responded with wheeled greenhouse-like structures (specularia) that could be moved indoors or outdoors to control temperature an early form of protected cultivation ^[37].
- Far East traditions also contributed: by the 15th century, the Korean Joseon dynasty had developed Ondol greenhouses glass-covered structures with underfloor heating to grow vegetables in winter. Medieval Europe had monastic gardens and later botanical gardens where exotic crops were grown. During the Renaissance and Enlightenment, naturalists and horticulturists systematically documented plant varieties. The scientific revolution brought quantitative approaches: in the 17th century, Jan Baptista van Helmont (Belgian chemist) showed plants derive mass largely from water (not soil alone) and John Woodward (English physician) found that water quality affects plant growth (leading to early fertilizer use ideas) ^[19].
- The 18th-19th centuries saw formal scientific contributions. Justus von Liebig (Germany) pioneered agricultural chemistry, identifying the role of minerals (nitrate, phosphate) in plant nutrition spurring artificial fertilizer development. Julius von Sachs developed nutrient solutions to grow plants in water, laying groundwork for hydroponics. Plant breeding advanced as well: Luther Burbank (USA, late 1800s-early 1900s) created thousands of new vegetable varieties (for example, new strains of spinach and potatoes) through hybridization and selection. Nikolai Vavilov (Russia, early 1900s) identified centers of crop origin and assembled seed collections, preserving the genetic diversity of vegetables and other crops. By the mid-20th century, controlled-environment agriculture took off: NASA and agronomists experimented with soilless culture and artificial lighting for space crops ^[55].

Table 1: Key Historical Milestones in Vegetable Cultivation ^[20, 38, 72]

Year/Period	Event/Milestone
~8000-5000 BCE	Neolithic Revolution: Domestication of edible plants (e.g. legumes, cereals, vegetables)
~600 BCE	Hanging Gardens of Babylon (legend): Ancient irrigated terraces for year-round crops
1st century AD	Roman "Greenhouses": Emperor Tiberius' mobile beds for winter cucumbers
1440s	Korean Ondol Greenhouses: Heated glasshouses for winter vegetable growth
1627	Francis Bacon: Publishes <i>Sylva sylvarum</i> , including early greenhouse experiments
1648	Jan van Helmont: Willow experiment showing plant mass comes from water
1699	John Woodward: Demonstrates better growth in nutrient-rich water vs distilled
1865	Julius von Sachs: Formulates a liquid nutrient solution (foundation of hydroponics)
1870s	Luther Burbank: Breeds thousands of new plant varieties (spinach, potato, etc.)
1914	Nikolai Vavilov: Identifies centers of origin for crops, establishes one of first seed banks
1937	William Gericke: Coins the term "hydroponics", grows 7-meter tomatoes in nutrient culture
1940	Hoagland Solution: Standard nutrient solution introduced for hydroponic research
1960s	NASA Plant Research: Development of Controlled Environment Life Support (MOWS unit)
1970s	Commercial Hydroponics: Emergence of large-scale soilless vegetable production (plastic greenhouses, drip irrigation)
1999	Vertical Farming Concept: Dr. Dickson Despommier popularizes multi-layer indoor farms
2000s	LED Grow Lights: Adoption of energy-efficient lighting transforms indoor cultivation
2010s	Urban Vertical Farms: Establishment of commercial vertical farms (multi-tier greenhouses, aeroponics)
2020s	AI & Robotics: Integration of sensors, AI and robotics in greenhouses and vertical farms

Traditional and Age-Old Practices

Traditional vegetable farming methods were shaped by the tools, climate and knowledge of their time. In many regions, gardens were small and labour-intensive. Farmers hand-tilled the soil with simple tools, broadcast or hand-planted seeds in rows or hills and used crop rotations and animal manures to maintain fertility. Seed selection was a key practice: farmers saved seed from the healthiest, most productive plants, a form of early genetic selection ensuring gradual improvement ^[39]. Key traditional practices included:

- **Soil Preparation:** Ploughing or digging to loosen the earth, addition of organic matter (farmyard manure, compost) to enrich the soil. Clods were broken and debris removed to create a fine tilts for sowing ^[56].
- **Sowing:** Seeds were either broadcast by hand, scattered in rows with a stick, or planted individually. Adequate spacing (based on crop type) prevented overcrowding and ensured each plant had enough light and nutrients ^[73].

- **Irrigation and Water Management:** Most irrigation was rudimentary. In wetter climates, reliance on rainfall was common, in semi-arid regions, farmers used basic canals, buckets or watering cans. They gauged soil moisture by feel and observation. Raised beds or ridges were used in flood-prone areas to improve drainage ^[89].
- **Nutrient Supply:** In the absence of synthetic fertilizers, fertility came from organic sources. Farmyard manure was the primary amendment, often composted in winter. In some cultures, ashes (lime), plant residues, or animal bone/guano were added. Folk practices like mulching with straw conserved moisture and added organic matter ^[90].
- **Weed and Pest Control:** Weed removal was manual (hand-pulling or hoeing). Pests and diseases were managed by cultural and botanical means. For example, certain plants (e.g., marigolds, onions, garlic) were

interplanted or their extracts used as insect repellents. Traditional biopesticides included extracts of neem, chili or tobacco. Mechanical traps and barriers (sticky traps, trapping boards) were also used. Crop rotation helped reduce soil pathogens and pests ^[21].

- **Crop Rotation and Polyculture:** Alternating crops (legumes, grains, vegetables) in fields prevented nutrient depletion and broke pest cycles. In some systems (e.g. the Three Sisters of the Americas), complementary species were grown together (maize, beans, squash) to improve soil and utilization ^[50].
- **Harvesting and Post-harvest:** Harvest was by hand with knives or sickles, selecting ripe produce carefully. Post-harvest, vegetables were often stored in cool cellars or preserved (drying, pickling, fermentation) due to lack of cold storage ^[1].

Table 2: Notable Contributors to Vegetable Cultivation ^[57, 40]

Contributor (Years)	Contribution/Field
Justus von Liebig (1803-1873)	Advanced plant nutrition theory, pioneered artificial fertilizers
Gregor Mendel (1822-1884)	Genetics, principles of heredity from pea experiments
Julius von Sachs (1832-1897)	Plant physiology, formulated nutrient solutions for plant growth
Charles Darwin (1809-1882)	Evolutionary theory, studied plant variation and breeding
Luther Burbank (1849-1926)	Developed thousands of crop varieties (spinach, potatoes, etc.) by hybridization
Nikolai I. Vavilov (1887-1943)	Identified centres of crop origin, established one of the first seed gene banks
Albert Howard (1873-1947)	Promoted organic farming and composting in horticulture
Norman Borlaug (1914-2009)	"Green Revolution" breeder, increased crop yields through improved varieties (e.g. dwarf grains)
William F. Gericke (1887-1979)	"Father of hydroponics", popularized soilless cultivation of vegetables
Dennis R. Hoagland (1886-1971)	Developed Hoagland nutrient solution, key figure in hydroponic research

Transition Phases in Vegetable Cultivation

- In the 1970s, many farmers worldwide still relied on knowledge passed down generations: hand-sowing in soil amended with compost and manure. Labor was intensive and pest control was mostly botanical or mechanical. However, even then, new tools like plastic seedling trays became popular. The introduction of drip irrigation revolutionized water use, offering precision hydration that saved water and increased yields. A notable breakthrough of the era was the spread of low-cost polyethylene greenhouse tunnels, making protected cultivation accessible beyond wealthy operations ^[74].
- The 1980s saw adoption of more modern inputs. Synthetic fertilizers and pesticides became commonplace, boosting short-term yields but raising environmental concerns. Greenhouses gained heating systems (often propane or wood-fired boilers) in temperate zones, enabling winter production of tomatoes and cucumbers in Europe and North America. The concept of raising seedlings in *plugs* (cells in trays) became standard, improving uniformity. Crucially, the notion of *vegetable quality* emerged, farmers began selecting for traits like shelf-life, taste and nutritional content, not just yield ^[22].
- By the 1990s, biotechnology and precision farming entered the picture. Seed technologies evolved: seed

priming (pre-soaking seeds to improve germination) and *pelleting* (coating seeds for easier handling) were developed. Tissue culture became viable for mass-producing disease-free seedlings, especially for high-value crops like potatoes and ornamentals. There was a surge in *biologicals*: biofertilizers (beneficial microbes) and biopesticides (e.g. *Bacillus thuringiensis* toxins) offered eco-friendly alternatives. GPS-guided equipment and GIS mapping started to be used on farms, although more in row crops, the tools foreshadowed precision applications in horticulture ^[2].

- With the turn of the millennium, digital and automation technologies began to influence vegetable production. Farmers and scientists combined sensors with satellite/GPS data to monitor fields and greenhouses. Drip and micro-spray irrigation became finely tunable by computer controllers. In greenhouses, climate computers regulated temperature, humidity and CO₂ levels to optimal setpoints. Robotics made early inroads: transplanters and harvest assistants appeared and automatic climate screens and fans responded to sensors. The grafting of tomatoes and cucurbits onto disease-resistant rootstocks became popular in regions plagued by soil pathogens, increasing resilience without genetic modification of the fruit ^[41].

Table 3: Comparative Overview of Vegetable Cultivation Systems ^[75, 58]

System/Method	Characteristics, Advantages, Limitations
Open-field (traditional)	Outdoor soil-based production, low capital costs, subject to weather and pests, limited season, large land required
Irrigated field (soil)	Adds controlled watering (drip/flood) to fields, higher yield than dryland, still climate-dependent, risk of nutrient leaching
Rainfed subsistence	Small-scale relying on natural rainfall, minimal inputs, low yield, very vulnerable to droughts and climate variability
Shade house	Partially shaded structure (netting), lowers sunlight intensity, protects from sunscald, lower yield than full sun
Polytunnel/High tunnel	Plastic-covered unheated or passively heated greenhouse, extends season, inexpensive, modest climate control, ventilation-dependent
Glasshouse (soil)	Glass/steel greenhouse with soil beds, year-round potential, higher yields, expensive structure and energy for heating
Greenhouse (hydroponic)	Glass/plastic structure with nutrient solution systems, very high yields, precise water use, high setup & energy costs
NFT Hydroponics	Shallow recirculating nutrient film over roots, water-efficient, easy to maintain, limited plant size buffer (pump risk)
Deep Water Culture (DWC)	Plants float roots in aerated solution, simple design, high oxygenation, risk of hypoxia if aeration fails
Aeroponics	Roots suspended in mist, ultra water-efficient, rapid growth, requires precise pump/atomizer, sensitive to disruptions
Aquaponics	Combines hydroponics with fish farming, recycles nutrients from fish waste, multi-use yields, complex balance, requires skills
Vertical Farming	Multi-layer indoor cultivation with artificial light, maximizes space use, year-round, very high energy and capital cost

These transition phases set the stage for the current era of high-tech cultivation, blending genetics, engineering and information technology. Traditional reliance on experience and broad tools gave way to precise, data-driven management. The next section examines the state of the art in greenhouse-based vegetable production ^[3].

Modern High-Tech Greenhouse Systems

Modern greenhouses and indoor farms represent the convergence of multiple technologies. Unlike a simple hothouse, these controlled-environment systems integrate hydroponics, advanced lighting, precise climate control, sensors, automation and sometimes robotics. Collectively, such approaches are often called Controlled Environment Agriculture (CEA). In this section we dissect the main components of high-tech greenhouse cultivation ^[42].

Soilless Cultivation: Hydroponics and Aeroponics

Hydroponics is the practice of growing plants without soil, using nutrient-enriched water solutions. It allows direct delivery of minerals, control of nutrient levels and elimination of soil-borne pests ^[91]. There are several hydroponic methods:

- **Nutrient Film Technique (NFT):** A thin film of nutrient solution continuously flows past plant roots in a sloped channel. Common for leafy greens and herbs. Its advantages are very high water-use efficiency and simplicity. Drawback: plants have little solution buffer, so pump failure or power loss can quickly stress plants ^[76].
- **Deep Water Culture (DWC):** Plant roots dangle in a reservoir of oxygenated solution. Often used for larger plants like tomatoes in greenhouses. Benefits include simplicity and high oxygenation (with air stones). Risk: any drop in dissolved oxygen (e.g. pump outage) can rapidly harm roots ^[23].
- **Drip Irrigation (closed-loop):** Nutrient solution is dripped onto plant roots (in a substrate like coco coir or rockwool) and excess drains back to the reservoir. Widely used for vegetables like tomatoes, cucumbers,

peppers. It supports larger plants and yields, but requires monitoring of EC (electrical conductivity) and pH to prevent salt build-up ^[4].

- **Wick Systems:** Passive, simpler for home or herb gardens. Plants in a growing medium wick solution from a reservoir. Low cost but not suitable for large or water-demanding crops ^[59].
- **Aeroponics:** A related technology where plant roots are suspended in air and periodically misted with nutrient solution. It is extremely water-efficient and can promote vigorous root growth. NASA has utilized aeroponics for potential space agriculture. However, it requires precise control of misting, nozzle or pump failure can quickly desiccate roots. It is used in some high-tech indoor farms ^[43].

Hydroponic greenhouses can dramatically increase yields. For example, lettuce grown hydroponically can cycle every 30 days year-round, compared to one or two outdoor crops. Studies report hydroponic systems using up to 90% less water than field agriculture (by recirculating water). Nutrient management is crucial: growers use formulations (like the Hoagland solution) tailored to crop needs, often adjusting for each growth stage (higher nitrogen during leaf growth, more potassium/boron during fruiting, etc.) ^[77]. Many modern greenhouses feature fertigation controllers automated injectors that mix and deliver precise nutrient and pH levels as plants require ^[60]. Aeroponic systems (roots in mist) share advantages with hydroponics but emphasize even greater aeration and nutrient uptake. Research indicates up to 95% water savings over soil culture. Aeroponics often requires higher initial engineering (mist nozzles, timers, filtration) and is sensitive to electrical disruptions, but it yields fast growth cycles, especially for leafy greens and some vegetables. For example, experiments have shown tomato plants growing 50% faster in aeroponic conditions than in traditional media ^[5]. Regardless of the specific method, soilless systems offer uniformity and cleanliness. Without soil pathogens, many diseases (*Fusarium*, *Pythium*) are minimized. Greenhouses can be sterile zones: before

planting, the substrate and benches are sanitized and then no external soil is introduced. This hygiene level improves seedling survival and reduces pesticide needs ^[44].

Vertical Farming and Multi-Tier Cultivation

Vertical farming takes soilless cultivation to three dimensions. Crops are grown in stacked layers (shelves, towers, or re-purposed warehouses), enabling much higher yields per ground footprint. For example, one analysis suggests a vertical farm can produce the same yield in one acre that would require up to 40 acres of traditional farmland. Typically, vertical farms use hydroponic or aeroponic methods combined with artificial lighting (usually LEDs) ^[24]. This approach is particularly suited for high-value, fast-growing vegetables and herbs (lettuce, spinach, microgreens, basil, strawberries) ^[61]. Challenges include high energy consumption for lights and climate control. However, vertical farms can exploit waste heat and CO₂ from adjacent industries, use renewable energy and place production closer to consumers (e.g. in cities) to reduce food miles. The yield benefits are enormous: some reports note vertical yields 50-100 times greater per square foot than conventional farming ^[6].

Controlled Environment Agriculture (CEA) and Greenhouses

Controlled Environment Agriculture (CEA) encompasses high-tech greenhouses and indoor farms with tight climate regulation. Key features include:

- **Structure and Envelope:** Modern greenhouses use glass or polycarbonate panels for high light transmission with insulation. Designs may be arched, gabled, or ridge-and-furrow, often with roof vents and side curtains ^[62].
- **Climate Control:** Integrated systems monitor and adjust temperature, humidity and CO₂. Heating (boilers, geothermal, biogas) is used in cold climates, cooling can involve fans, evaporative pads, or adiabatic walls in hot regions. Automated vents and shade screens respond to weather sensors. CO₂ generators or enrichment tanks boost CO₂ levels to 800-1200 ppm, elevating photosynthesis and yields ^[78].
- **Lighting:** Supplemental lighting extends day length and intensity. The shift to LED technology has been transformative. LEDs consume far less power than older high-pressure sodium lamps and can emit targeted spectra (red, blue, far-red, etc.). Growers tune the light spectrum for specific crops for instance, extra red light to boost tomato fruiting, or blue light to produce compact lettuce. Adjusting photoperiod (night duration) can accelerate flowering in day-neutral crops. Light intensity and duration are automated by sensors and controllers ^[92].
- **Irrigation and Nutrition:** Drip and ebb-and-flow systems precisely deliver water and dissolved nutrients. Recycled systems capture drain water, filtering and re-dosing it. The nutrient composition is continuously monitored via electrical conductivity and pH sensors, allowing fine-tuning. Some advanced setups even adjust nutrient mixes automatically based on plant stage and sensor feedback ^[25].
- **Sensors and Automation:** The heart of a smart greenhouse is its sensor network. Sensors measure air temperature, humidity, CO₂, leaf wetness, root-zone

moisture, nutrient levels and even plant stress indicators. This data feeds into controllers or cloud-based platforms. Automated actuators then adjust heaters, vents, lights, pumps and misting nozzles. In many facilities, climate computers integrate all sub-systems: for example, turning off CO₂ injection if ventilation is open or dimming lights if natural sun is sufficient ^[7]. Modern greenhouses often include cameras and imaging sensors. These can do remote crop monitoring: cameras detect leaf colour changes, growth rate, or pest outbreaks. Artificial intelligence algorithms analyse these images to forecast issues (e.g. nitrogen deficiency signs) before visible symptoms appear ^[93].

- **Automation and Robotics:** High-tech operations are increasingly using robotics. In some large greenhouses, robotic arms handle tedious tasks like sorting and packaging tomatoes. Autonomous vehicles are being developed to scout and act on plant health issues. Pollination (essential for tomatoes, peppers) may be performed by robotic pollinators or simply by distributing bumblebee hives under controlled conditions ^[45].

Advantages and challenges

The payoff of high-tech greenhouses is improved yield, quality and resource efficiency. Yields per hectare can be an order of magnitude higher than open field. Crops can be grown year-round, free from seasonal limitations. Protection from weather (frost, hail, heavy rain) and pests reduces losses. However, these systems are capital- and skill-intensive. The initial investment for a full controlled-environment greenhouse (structure, heating/cooling, lighting, controls) is high. Energy costs (especially for heating and lighting) are a major factor. Economically, such systems are typically justified for high-value vegetables (e.g. tomatoes, peppers, leafy greens) where markets can bear the added cost ^[8].

Advances in Crop Cultivation within CEA

Beyond structural and environmental tech, modern cultivation has innovated in plant materials and propagation:

- **Variety selection:** Breeding has created cultivars suited for greenhouse conditions: compact growth, high fruit set in low wind, disease resistance (e.g. *Fusarium* wilt-resistant cucumbers). Hybrid seeds dominate, giving uniform and predictable crops. Some varieties are tailored to vertical farming (e.g. red-leaf lettuce responds well to certain LED spectra). Genetically modified varieties (e.g. virus-resistant tomatoes or Bt brinjal engineered to resist borers) exist in some regions, though their use in CEA is mixed due to regulations and market acceptance ^[79].
- **Propagation:** Seedling production has become an industry in itself. Many growers buy greenhouse-grown plug seedlings rather than seeding directly. Tissue culture (micropropagation) is used for crops like potatoes and strawberries, producing disease-free clones. Some vegetables (e.g. tomatoes, peppers) are grafted onto vigorous rootstocks to improve nutrient/water uptake and disease tolerance, grafting stations have automated much of this once manual process ^[63].
- **Microbial technology:** There is growing use of plant growth-promoting rhizobacteria (PGPR) and

mycorrhiza inoculants even in hydroponics, aiming to mimic beneficial soil interactions. These can enhance nutrient use efficiency and stress tolerance. Research is also exploring the greenhouse microbiome to naturally suppress pathogens, reducing need for chemical fumigation ^[46].

In summary, modern high-tech greenhouses combine advanced materials (glazing, plastics), engineering (climate systems, lighting) and information technology (sensors, AI) with sophisticated horticulture (hydroponics, IPM, optimized genetics). This multi-disciplinary integration allows unprecedented control over vegetable production ^[9].

Crop Management in Vegetable Cultivation

Intensive production in either open fields or greenhouses depends on optimized crop management. Key elements include genetics (varieties), propagation methods, nutrition management and pest/disease control ^[80]. Below, we outline each component.

Varieties and Breeding

Modern agriculture uses improved vegetable varieties to maximize yield, quality and stress tolerance ^[94]. Breeding programs (public and private) continuously develop hybrids and pure lines for various traits:

- **Hybrid vs Open-Pollinated:** Most commercial production relies on F1 hybrids. These are crossbred for uniformity and vigour. Heirloom and open-pollinated varieties are still valued by small farmers for heritage qualities. Hybrids offer advantages like higher yield, disease resistance and consistent maturation ^[64].
- **Genetic Modification and Editing:** A few genetically modified vegetables exist (e.g. Flavr Savr tomato, Bt eggplant), though many GM traits are found in commodities. Gene editing (CRISPR/Cas9) is being explored for traits like shelf-life and disease resistance with potentially fewer regulatory hurdles ^[26].
- **Traits for Controlled Environments:** Greenhouse-specific traits include shorter internodes, better fruit setting at high CO₂ and tolerance to low light. There is also emphasis on nutritional content and flavour as market demands rise ^[81].
- **Disease Resistance:** Breeders incorporate resistance genes to major pathogens (e.g. Fusarium wilt in cucumbers, tomato mosaic virus), reducing reliance on chemicals and improving consistency ^[47].

Propagation and Seedling Production

Vegetables propagate either by seed or vegetative means:

- **Seeds:** Most vegetables (tomatoes, beans, cabbages, etc.) are grown from seed. Seed selection (size, vigour) is key. Seed treatments (soaking, priming, fungicide/insecticide coatings) enhance germination. Many greenhouse operations pre-germinate or raise seedlings in trays, then transplant robust seedlings into final beds or systems ^[95].
- **Cuttings and Transplants:** Some crops (e.g. basil, strawberries, sweet potatoes) can be started from cuttings or runners. Transplanting methods are advanced: machines can set young plants (tomato, pepper, eggplant) into beds rapidly. Tissue culture laboratories produce disease-free transplants for potato, vanilla, bananas and other crops ^[62].

- **Tissue Culture (Micropropagation):** This *in vitro* technique clones plants in sterile medium. It's widely used for crops like strawberries, lilies and seedless cucumbers, ensuring large numbers of uniform, pathogen-free plants. Specialized labs control the process to avoid somaclonal variation ^[27].
- **Grafting:** Common in tomato and cucumber production, grafting combines a vigorous rootstock (resistant to soil diseases, tolerant of high EC) with a high-yielding scion. Mechanized grafting lines can produce thousands of grafts per day ^[10].

Nutrition and Soil Management

Fertilization strategies have shifted from broad generalizations to precision management:

- **Soil vs Soilless:** In soil-based systems, fertilization still relies on a combination of organic amendments and synthetic fertilizers. Greenhouses often use soilless media (coco coir, rockwool, perlite) or none (hydroponics). Nutrient delivery is then via water-soluble formulations ^[65].
- **Fertigation and Precision Feeding:** In high-tech systems, fertilizers are delivered through irrigation (fertigation) based on real-time needs. EC (electrical conductivity) and pH sensors in the nutrient solution ensure levels stay within target ranges. Automated dosing pumps adjust nutrient supply daily or weekly as plants grow ^[82].
- **Nutrient Balancing:** Different vegetables and growth stages require specific N-P-K ratios. For example, leafy crops demand higher nitrogen, whereas fruiting vegetables need more phosphorus and potassium for flower and fruit development. Micronutrients (Fe, Mg, B, Zn, etc.) are also routinely supplied ^[28].
- **Soil Health and Substrates:** In greenhouses using soil or compost, crop rotation and cover cropping (in cover bed areas) help maintain fertility. Sterilization (steam or chemical fumigation) is often practiced between crops to kill soil pathogens. In hydroponics, substrates like coco coir are occasionally refreshed or recharged to prevent pH drift and salt build-up ^[96].
- **Advanced Amendments:** Technologies such as controlled-release fertilizers (CRFs) release nutrients slowly in soil systems, while organic coatings minimize nutrient leaching. Microbial inoculants (mycorrhizae, N-fixers for legumes) can be applied to promote root health ^[48].

Pest and Disease Management

Pests (insects, mites) and diseases (fungal, bacterial, viral) are major constraints. Management strategies combine preventive, cultural, biological and chemical methods:

- **Prevention and Monitoring:** Greenhouses implement strict hygiene: workers may wear protective clothing, tools are sterilized and incoming plant material is quarantined. Yellow and blue sticky traps monitor flying pests (whiteflies, thrips). Regular scouting (visual or via cameras) detects early signs of infestation or disease ^[11].
- **Integrated Pest Management (IPM):** This approach, now standard in modern cultivation, integrates multiple tactics:

- **Physical/Mechanical:** Insect screens on vents exclude pests, sticky traps catch adults, nets or row covers physically protect crops [83].
- **Cultural Controls:** Crop rotation, debris removal and disinfection (e.g. using hydrogen peroxide or UV-light devices) reduce pathogen build-up. Adjusting humidity and temperature can make conditions unfavourable for specific diseases (e.g. lowering humidity to prevent powdery mildew) [97].
- **Biological Controls:** Beneficial insects (ladybugs, predatory mites, parasitic wasps) are released to suppress pests. Biopesticides derived from entomopathogenic fungi or bacteria are used (e.g. *Beauveria bassiana* against whiteflies).
- **Resistant Varieties:** Using disease-resistant cultivars (e.g. powdery mildew-resistant cucumber) is a key strategy [29].
- **Chemical Controls:** Modern pesticides are used as a last resort, often tailored precisely (spot treatments, minimal-risk compounds). Application may be automated via sprayers or foggers. Even organic-approved chemicals (Bt toxins, neem oil) are used sparingly under IPM guidelines [84].
- **Disease Management:** Fungal diseases (e.g. Fusarium, Phytophthora, powdery mildew) are controlled by sanitation, resistant cultivars and targeted fungicides. Viruses are often managed by controlling insect vectors (e.g. controlling thrips to prevent tomato spotted wilt virus). Bacterial diseases require exclusion and biocontrol (some beneficial bacteria compete with pathogens) [98].
- **Sanitation of Equipment:** Trays, tools and conveyances are sanitized regularly. In hydroponics, ultraviolet (UV) lights or ozone may treat recirculating nutrient water to kill microbes [30].

Table 4: Crop Management Attributes for Common Vegetable Crops [12, 49, 66]

Vegetable	Propagation	Typical Cultivation (Environment)	Nutrient Focus
Tomato	Seeds (hybrids), grafting	Greenhouse or field, often hydroponics	High potassium, balanced N
Lettuce	Seeds	Hydroponics/Greenhouse/Vertical	High nitrogen (leafy growth)
Cucumber	Seeds, cuttings (greenhouse)	Greenhouse (often hydroponic)	Balanced NPK, extra K for fruits
Onion	Bulbs, sets, or seeds	Field	High sulfur and potassium
Carrot	Seeds	Field (deep, loose soil)	Adequate potassium and boron
Broccoli	Transplants	Field or unheated greenhouse	High nitrogen early, then balanced
Kale	Seeds/transplants	Field or greenhouse	High nitrogen
Beans	Seeds (legumes)	Field or greenhouse (warmer climates)	Phosphorus for root development, fix own N
Leafy Greens (e.g. arugula, etc.)	Seeds	Hydroponics/greenhouse	High nitrogen
Herbs (basil, mint)	Seeds/cuttings	Hydroponics or pots	Moderate N, prefer steady moderate feeding
Strawberry (fruit)	Runners or tissue-cultured plants	Greenhouse/vertical	High potassium and calcium

Economics and Scalability

Economic viability is a crucial factor driving adoption of cultivation methods. High-tech systems require substantial capital investment but promise higher returns through increased yield and value-added products.

- **Capital Investment:** Building a high-tech greenhouse (with glass panels, heating/cooling systems, automation, lighting) can cost hundreds to thousands of dollars per square meter. In contrast, open-field farming requires minimal upfront infrastructure (beyond land and basic tools). For vertical farms, the cost is even higher due to multi-story construction and heavy equipment. Large-scale studies show payback periods for greenhouses can range from 5-15 years, depending on crop prices and technology used [67].
- **Production Costs:** Operating expenses include energy, labour, nutrients and maintenance. High-energy use (heating in winter, lighting year-round) makes electricity and fuel major costs. However, automation and mechanization can reduce labour requirements: e.g., one worker can manage far more area in a computerized greenhouse than in fields. Fertilizer and water costs are lower per unit yield in hydroponics due to efficiency, but the specialized nutrient formulations can be expensive [13].
- **Yields and Output Value:** High-tech cultivation can multiply yields several-fold. For example, greenhouse tomatoes often produce 5-10 kg/m² annually, versus 3-5 kg/m² outdoors. Leafy greens in vertical farms can produce 30-40 harvests per year per shelf space. Combined with out-of-season production, growers capture premium prices (winter tomatoes or fresh lettuce in desert regions fetch higher prices). A report noted vertical farming can produce up to 60 times the yield per acre compared to field farming [50].
- **Market and Demand:** Urban markets can support high-tech farms because of consumer demand for local, fresh and pesticide-free produce. Direct-to-consumer models (farmers' markets, CSA, or grocery partnerships) often yield better prices. In addition, specialized products (seedlings, microgreens, organic-certified produce) open niche markets. However, growers must be careful of market saturation, if many compete in a small area with similar crops, prices can fall [31].
- **Scalability:** Large-scale adoption of high-tech farming is most advanced in places like the Netherlands, Japan and now China and the UAE, where land is scarce or climate harsh. Government incentives (subsidies, research funding) have spurred growth in these regions. Nevertheless, even small farms can adopt elements: for example, a family can start a greenhouse with plastic film and basic hydroponics to increase productivity on limited land. Scalable technologies include modular hydroponic kits and "smart" controllers that even smallholders can use via smartphone apps [14].

Table 5: Comparative Economics of Cultivation ^[51, 85]

System	Capital Cost	Yield per Area	Water Use Efficiency	Energy Use	Typical Crops
Open-field (soil)	Low	Baseline	Low (high usage)	Low	Most field crops
Greenhouse (soil)	Medium	5-12× field	Medium (drip/recirc)	Medium (heating in winter)	Tomatoes, peppers
Hydroponic GH	High	10-20× field	Very high (90% saved)	High (lights, pumps)	Lettuce, herbs, cucumbers
Vertical Farm	Very High	50-100× field	Extremely high	Very High (lights, climate)	Lettuce, microgreens
Aquaponics	High	5-10× field	High (recycled)	High	Leafy greens, herbs

Economic and Social Considerations

The high cost of technology is a major barrier for many. Small farmers in developing countries may not afford greenhouses or hydroponic setups. Microfinance, cooperatives, or public-private partnerships can help. For example, community greenhouses have been implemented in some regions to teach sustainable horticulture ^[68]. From a workforce perspective, these systems require skilled operators familiar with sensors and software, not just traditional farming knowledge. Training and extension services are crucial. On the other hand, they create new jobs in engineering and IT related to farming. In aggregate, global investment in advanced agriculture is rising, reflecting the economic potential. The market for hydroponic systems, greenhouse structures and vertical farms is growing annually. Companies supplying LED lights, climate computers and robotics see booming sales. This suggests a strong belief in the profitability of high-tech farming where conditions are right ^[32].

Sustainability and Environmental Impacts

One of the main motivations for high-tech vegetable cultivation is sustainability producing more food with fewer resources and lower environmental impact. We examine key dimensions: water use, land use, energy and emissions and broader ecological effects.

- **Water Use:** Controlled-environment systems excel at water conservation. Recirculating hydroponics can use up to 90% less water than comparable field irrigation because runoff is collected, filtered and reused. Even vertical farms (often cited as “10×” savings) dramatically reduce water footprints, critical in arid regions. This efficiency helps protect freshwater resources. However, water quality must be monitored to prevent salt accumulation, necessitating occasional reservoir flushing or replenishment with clean water ^[15].
- **Land Use:** Vertical and greenhouse farming drastically reduce land requirements. A vertical farm can cut land use by over 95% compared to field farming for the same output. This allows food production near or within cities, reducing transportation emissions and land conversion pressures. It also means less deforestation and habitat loss for agriculture. On the other hand, these systems may rely on non-renewable materials (plastics, metals) and generate waste (spent media, bulbs) ^[52].
- **Energy and Carbon Footprint:** Energy is the Achilles’ heel of indoor farming. Heating greenhouses in cold climates and powering LED lights year-round can generate significant CO₂ emissions unless renewables are used. Life-cycle studies show vertical farming lettuce, for example, can have a higher carbon footprint per kg than field-grown, due largely to electricity use. However, efficiency is improving: LED efficacy (μmol/J) doubles roughly every few years and integration with solar/wind power can offset carbon. In addition, thermal insulation and combined heat-and-

power (CHP) systems in greenhouses can reuse waste heat and CO₂, improving overall sustainability ^[33].

- **Chemical Use:** High-tech systems can reduce pesticide and fertilizer pollution. Since greenhouses exclude many pests, pesticide applications are often lower or replaced by biological controls. Hydroponics prevents nutrient runoff, a common source of waterway pollution in traditional farms. Any waste solution is contained rather than discharged. However, reliance on synthetic fertilizers (manufactured by energy-intensive processes) remains an issue, bio-based nutrient sources or closed-loop recycling of waste could improve this in the future ^[69].
- **Biodiversity and Local Impact:** Urban farming can re-green cities and provide jobs, but extensive greenhouse complexes may change local microclimates (though much smaller footprint than open fields). Discarded nutrient solutions and growing media must be managed properly. Some facilities compost crop residues or use them for bioenergy, creating a circular system ^[86]. The net environmental impact thus depends on context and technology integration. Ideally, renewable energy powers these farms, waste streams are minimized and the benefits of high yields and local production offset any energy costs. Policymakers and researchers are studying these trade-offs to guide sustainable deployment ^[16].

Future Directions and Research Gaps

Looking ahead, vegetable cultivation will continue evolving. Key trends and unanswered questions include:

- **Digital Agriculture and AI:** The role of machine learning and data analytics is still expanding. Future greenhouses may use predictive models to anticipate plant needs (watering, nutrients, pest risk) and actuate systems in real time. Robots may do planting, thinning and even harvesting delicate crops in lieu of manual labour. Drones inside large greenhouses could monitor plants or even assist pollination. Development of user-friendly decision-support tools for farmers will be critical ^[34].
- **Genetic and Biotechnologies:** Gene editing could produce varieties specifically tailored for hydroponics or low-light environments (e.g. algae-based photosynthetic boost). Microbiome engineering (introducing beneficial microbial consortia to nutrient solutions) is a promising area to enhance resilience and yield. There is also research into edible vaccines and functional foods via plant biotechnology which might be grown in contained systems for biosecurity ^[53].
- **Energy and Climate Resilience:** Innovations are needed to cut the energy footprint. This includes more efficient LEDs, greenhouse-integrated solar panels (agrivoltaics), geothermal heating and novel insulation materials (e.g. phase-change materials for thermal buffering). In terms of climate change, controlled-

environment systems themselves become refuges against extreme weather for food production ^[87].

- **Integration and Urban Farming:** Plans to integrate high-tech farms into urban planning (rooftop greenhouses, vertical farms in buildings) are underway. Research on how to scale city farms sustainably including wastewater reuse and integrating with municipal infrastructure is a gap. Social research is also needed on how communities benefit (or not) from local high-tech farms ^[99].
- **Resource Recycling and Circularity:** Closed-loop systems (e.g. recycling all water and nutrients, composting plant residues) are the ideal. However, practical models for on-site waste processing, or creating biogas from residues, need development. Similarly, strategies to dispose of or repurpose spent plastics and equipment sustainably are needed ^[70].
- **Economic Models:** More economic analyses on long-term viability under different scenarios (e.g. changing energy prices, carbon taxes, crop prices) will help investors. Research should target the viability of small-scale systems (for developing countries) versus industrial mega-farms ^[35].
- **Ecological Studies:** Long-term ecological comparisons (e.g. full life-cycle assessments of various crops in different systems) will clarify true sustainability. Effects on nutrition (e.g. differences in micronutrient content when grown hydroponically vs soil) are still being studied ^[17].

Conclusion

From the earliest kitchen gardens to towering vertical farms, the journey of vegetable cultivation spans millennia of innovation. This review charted that journey, highlighting how age-old principles (like careful seed selection and organic nutrition) now merge with sensors, automation and biotechnology. Traditional farmers relied on seasons and nature, modern growers augment nature with exacting control over every growth factor. Key scientists and pioneers laid the foundation (e.g. Liebig's fertilizers, Mendel's genetics, Gericke's hydroponics) that enabled the current leap forward. In controlled environments, vegetables can be produced more abundantly and with greater consistency than ever before. At the same time, we are learning how to do this efficiently: conserving water, recycling nutrients and using clean energy to minimize environmental costs. The economic case for high-tech cultivation is compelling for high-value vegetables, especially where land or climate is a constraint. However, high-tech does not automatically equal sustainability. Integrating renewable energy and circular practices is essential. Smallholders and developing regions face hurdles in financing and knowledge, inclusive solutions are needed to ensure these advances benefit all. Looking forward, advances in AI, robotics, genomics and materials science will continue to reshape vegetable farming. The challenges of feeding a growing population under climate change demand that we refine and adapt these approaches. Research must address current gaps (energy use, microbial management, socio-economics) to fully realize the promise of high-tech agriculture. Ultimately, the goal remains the same as in traditional farming: to produce safe, nutritious and abundant vegetables. What has changed is the scale and precision with which we achieve that goal. As this review

has shown, the horizon of vegetable cultivation continues advancing toward ever greater efficiency, sustainability and resilience. Each technological step builds on the last, forging a continuum from ancient gardens to futuristic farms. The collaboration of farmers, scientists, engineers and policymakers will determine how fully these horizons are reached in practice.

References

1. Abd-Elgawad MM. Plant-parasitic nematodes of strawberry in egypt: A review. Bull Natl Res Cent. 2019;43:1-13. doi: 10.1186/s42269-019-0049-2
2. Abdullah HM, Mohana NT, Khan BM, Ahmed SM, Hossain M, Islam KS, *et al.* Present and future scopes and challenges of plant pest and disease (P&D) monitoring: Remote sensing, image processing and artificial intelligence perspectives. Remote Sens Appl. 2023;32:100996. doi: 10.1016/j.rsase.2023.100996
3. Aini N, Yamika WD, Ulum B. Effect of nutrient concentration, PGPR and AMF on plant growth, yield and nutrient uptake of hydroponic lettuce. Int J Agric Biol. 2019;21:175-183. doi: 10.17957/IJAB/15.0879
4. Akrami M, Salah AH, Dibaj M, Porcheron M, Javadi AA, Farmani R, *et al.* A zero-liquid discharge model for a transient solar-powered desalination system for greenhouse. Water (Switzerland). 2020;12:1440. doi: 10.3390/w12051440
5. Alipio MI, Dela Cruz AE, Doria JD, Fruto RM. On the design of Nutrient Film Technique hydroponics farm for smart agriculture. Eng Agricult Environ Food. 2019;12:315-324. doi: 10.1016/j.eaef.2019.02.008
6. Alreshidi E. Smart Sustainable Agriculture (SSA) solution underpinned by Internet of Things (IoT) and Artificial Intelligence (AI). Int J Adv Comput Sci Appl. 2019;10:93-102. doi: 10.14569/IJACSA.2019.0100513
7. Ampatzidis Y, Partel V. UAV-based high throughput phenotyping in citrus utilizing multispectral imaging and artificial intelligence. Remote Sens (Basel). 2019;11:410. doi: 10.3390/rs11040410
8. Ampim PA, Sloan JJ, Cabrera RI, Harp DA, Jaber FH. Green roof growing substrates: types, ingredients, composition and properties. J Environ Hortic. 2010;28:244-252. doi: 10.24266/0738-2898-28.4.244
9. Antonious GF, Chiluwal A, Nepal A. Chitin, biochar and animal manures impact on eggplant and green pepper yield and quality. Agric Sci. 2023;14:368-383. doi: 10.4236/as.2023.143024
10. Anushi B, Sachan K. Bioformulation: A new frontier in horticulture for eco-friendly crop management. J Plant Biota. 2024;28.
11. Anushi M, Jain S, Sharma R, Thapliyal V. The Horticulture Encyclopedia.
12. Anushi RM, Deshmukh RN, Sharma R. From DNA to Deliciousness: A Journey into Molecular Markers in Fruits.
13. Anushi SS, Krishnamoorthi A, Kumar S, Pareta P, Kalaiselvi P, Sinha G, *et al.* Biotech bounty on verge: gm (genetically modified) crops and the science of sustainable agriculture and horticulture.
14. Anushi VK, Shukla P. Influence of biostimulants and organic mulch on soil microbial population in strawberry (*F. × ananassa* Dutch.).
15. Asghar S, Ghori N, Hyat F, Li Y, Chen C. Use of auxin and cytokinin for somatic embryogenesis in plant: a

- story from competence towards completion. *Plant Growth Regul.* 2023;99:413-428. doi: 10.1007/s10725-022-00923-9
16. Aslan MF, Durdu A, Sabanci K, Ropelewska E, Gültekin SS. A comprehensive survey of the recent studies with UAV for precision agriculture in open fields and greenhouses. *Appl Sci (Switzerland)*. 2022;12:1047. doi: 10.3390/app12031047
 17. Baidoo PK, Mochiah MB. Comparing the Effectiveness of Garlic (*Allium sativum* L.) and Hot Pepper (*Capsicum frutescens* L.) in the Management of the Major Pests of Cabbage *Brassica oleracea* (L.). *Sustain Agric Res.* 2016;5:83-91. doi: 10.5539/sar.v5n2p83
 18. Barceló-Muñoz A, Barceló-Muñoz M, Gago-Calderon A. Effect of led lighting on physical environment and microenvironment on *in vitro* plant growth and morphogenesis: The need to standardize lighting conditions and their description. *Plants.* 2022;11:60. doi: 10.3390/plants11010060
 19. Bárcenas-Moreno G, Jiménez-Compán E, San Emeterio LM, Jiménez-Morillo NT, González-Pérez JA. Soil pH and soluble organic matter shifts exerted by heating affect microbial response. *Int J Environ Res Public Health.* 2022;19:15751. doi: 10.3390/ijerph192315751
 20. Bar-Yosef B. Fertilization under drip irrigation. In: Palgrave DA, editor. *Fluid fertilizer: Science and technology*. Boca Raton: CRC Press; 2020. p. 285-329. Available from: <https://www.taylorfrancis.com/chapters/edit/10.1201/9781003065746-14/fertilization-drip-irrigation-bnayahu-bar-yosef> [Accessed 26th September 2023].
 21. Benelli G, Canale A, Toniolo C, Higuchi A, Murugan K, Pavela R, *et al.* Neem (*Azadirachta indica*): towards the ideal insecticide? *Nat Prod Res.* 2017;31:369-386. doi: 10.1080/14786419.2016.1214834
 22. Bhandari J, Nayama S. A review on nursery management in horticultural crops: A beneficial way for enhancing income. *Int J Chem Stud.* 2020;8:410-413. doi: 10.22271/chemi.2020.v8.i4g.10179
 23. Bhowmik P, Konkin D, Polowick P, Hodgins CL, Subedi M, Xiang D, *et al.* CRISPR/Cas9 gene editing in legume crops: Opportunities and challenges. *Legume Sci.* 2021;3:e96. doi: 10.1002/leg3.96
 24. Bista D, Sapkota D, Paudel H, Adhikari G. Effect of foliar application of growth regulators on growth and yield of onion (*Allium cepa*). *Int J Hortic Sci Technol.* 2022;9:247-254. doi: 10.22059/ijhst.2021.321019.451
 25. Brown CS, Schuerger AC, Sager JC. Growth and photomorphogenesis of pepper plants under red light-emitting diodes with supplemental blue or far-red lighting. *J Am Soc Hortic Sci.* 1995;120:808-813. doi: 10.21273/JASHS.120.5.808
 26. Caradus JR. Intended and unintended consequences of genetically modified crops-myth, fact and/or manageable outcomes? *New Z J Agric Res.* 2022;66:519-619. doi: 10.1080/00288233.2022.2141273
 27. Chalker-Scott L, Scott JD. Elevated ultraviolet-B radiation induces cross-protection to cold in leaves of rhododendron under field conditions. *Photochem Photobiol.* 2004;79:199-204. doi: 10.1562/0031-8655(2004)079<0199:eurict>2.0.co;2
 28. Chandel AK, Khot LR, Sallato C. Apple powdery mildew infestation detection and mapping using high-resolution visible and multispectral aerial imaging technique. *Sci Hortic.* 2021;287:110228. doi: 10.1016/j.scienta.2021.110228
 29. Chen C, Li Y, Li N, Wei S, Yang F, Ling H, *et al.* A computational model to determine the optimal orientation for solar greenhouses located at different latitudes in China. *Solar Energy.* 2018;165:19-26. doi: 10.1016/j.solener.2018.02.022
 30. Chen K, Li GJ, Bressan RA, Song CP, Zhu JK, Zhao Y. Absciscic acid dynamics, signaling and functions in plants. *J Integr Plant Biol.* 2020;62:25-54. doi: 10.1111/jipb.12899
 31. Deshmukh PK, Mehrete PS, Kokane SC, Joshi PE. Multi-tasking agricultural robot. *Int J Res Engineering Sci Manage.* 2020;3:336-338.
 32. Desurmont GA, Guiguet A, Turlings TC. Invasive insect herbivores as disrupters of chemically-mediated tritrophic interactions: effects of herbivore density and parasitoid learning. *Biol Invasions.* 2018;20:195-206. doi: 10.1007/s10530-017-1526-x
 33. Dhen N, Ben Ammar I, Bayoudh C, Al Mohandes Dridi B. Optimization of a novel vegetable nursery substrate using date palm wastes peat and indigenous arbuscular mycorrhizal fungi. *Commun Soil Sci Plant Anal.* 2019;50:959-973. doi: 10.1080/00103624.2019.1594879
 34. Ding H, Fu TJ, Smith MA. Microbial contamination in sprouts: how effective is seed disinfection treatment? *J Food Sci.* 2013;78:R495-R501. doi: 10.1111/1750-3841.12064
 35. Doan TN. An efficient system for real-time mobile smart device-based insect detection. *Int J Adv Comput Sci Appl.* 2022;13:30-36. doi: 10.14569/IJACSA.2022.0130605
 36. Galletti PA, Carvalho ME, Hirai WY, Brancaglioni VA, Arthur V, Barboza da Silva C. Integrating Optical Imaging Tools for Rapid and Non-invasive Characterization of Seed Quality: Tomato (*Solanum lycopersicum* L.) and Carrot (*Daucus carota* L.) as Study Cases. *Front Plant Sci.* 2020;11:577851. doi: 10.3389/fpls.2020.577851
 37. González-Guerrero M, Navarro-Gómez C, Rosa-Núñez E, Echávarri-Erasun C, Imperial J, Escudero V. Forging a symbiosis: transition metal delivery in symbiotic nitrogen fixation. *New Phytol.* 2023;239:2113-2125. doi: 10.1111/nph.19098
 38. Gorai T, Yadav PK, Choudhary GL, Kumar A. Site-specific crop nutrient management for precision agriculture-A review. *Curr J Appl Sci Technol.* 2021;40:37-52. doi: 10.9734/cjast/2021/v40i1031357
 39. Grant F, Sheline C, Sokol J, Amrose S, Brownell E, Nangia V, *et al.* Creating a Solar-Powered Drip Irrigation Optimal Performance model (SDrOP) to lower the cost of drip irrigation systems for smallholder farmers. *Appl Energy.* 2022;323:119563. doi: 10.1016/j.apenergy.2022.119563
 40. Gulzar B, Mujib A, Qadir Malik M, Mamgain J, Syeed R, Zafar N. Plant tissue culture: Agriculture and industrial applications. In: Kiran U, Abidin MZ, Kamaluddin, editors. *Transgenic technology based value addition in plant biotechnology*. Cambridge, Massachusetts: Academic Press; 2020. p. 25-49.

- doi: 10.1016/B978-0-12-818632-9.00002-2
41. Gupta A, Kumar R. Management of seed-borne diseases: An integrated approach. In: Kumar R, Gupta A, editors. Seed-borne diseases of agricultural crops: Detection, diagnosis & management. Singapore: Springer; 2020. p. 717-745. doi: 10.1007/978-981-32-9046-4_25
 42. Hewlett JA, Szczepaniec A, Eubanks MD. The effects of sugarcane aphid density in sorghum on predation by lady beetles and lacewings. Biol Control. 2019;129:171-177. doi: 10.1016/j.biocontrol.2018.10.015
 43. Huber BM, Louws FJ, Hernández R. Impact of different daily light integrals and carbon dioxide concentrations on the growth, morphology and production efficiency of tomato seedlings. Front Plant Sci. 2021;12:615853. doi: 10.3389/fpls.2021.615853
 44. Hussain J, Pan Y, Ali G, Xiaofang Y. Pricing behavior of monopoly market with the implementation of green technology decision under emission reduction subsidy policy. Sci Total Environ. 2020;709:136110. doi: 10.1016/j.scitotenv.2019.136110
 45. Husson O. Redox potential (Eh) and pH as drivers of soil/plant/microorganism systems: A transdisciplinary overview pointing to integrative opportunities for agronomy. Plant Soil. 2013;362:389-417. doi: 10.1007/s11104-012-1429-7
 46. Indergard MO, Bellantuono A, Rodriguez-Lanetty M, Heng F, Gilg MR. Acclimation to elevated temperatures in *Acropora cervicornis*: effects of host genotype and symbiont shuffling. Mar Ecol Prog Ser. 2022;701:41-65. doi: 10.3354/meps14192
 47. Jain S, Ranganna G, Jadhav PS, Shrivastava U, Sood A, Prasad L, Luthra S. Cultivating Tomorrow: Harnessing Gene Editing and Silencing for Precision Horticulture. J Adv Biol Biotechnol. 2024;27(1):147-160.
 48. Kacheyo OC, de Vries ME, van Dijk LC, Schneider HM, Struik PC. Agronomic consequences of growing field-transplanted hybrid potato seedlings. Crop Sci. 2023;1-19. doi: 10.1002/csc2.20997
 49. Kong Y, Kamath D, Zheng Y. Blue versus red light can promote elongation growth independent of photoperiod: A study in four brassica microgreens species. HortScience. 2019;54. doi: 10.21273/HORTSCI14286-19
 50. Koukounaras A. Advanced greenhouse horticulture: New technologies and cultivation practices. Horticulturae. 2021;7:1. doi: 10.3390/horticulturae7010001
 51. Krishnamoorthi A, Chaurasia J, Baidya BK, Singh A, Kumar V, Singh A. A comprehensive review on evolution, challenges of models, opportunities in urban farming practices in the world. J Sci Res Rep. 2024 Apr 25;30(6):136-156.
 52. Li R, Li Q, Pan L. Review of organic mulching effects on soil and water loss. Arch Agron Soil Sci. 2021;67:136-151. doi: 10.1080/03650340.2020.1718111
 53. Liu G, Simonne EH, Morgan KT, Hochmuth GJ, Agehara S, Mylavarapu R, et al. Chapter 2. fertilizer management for vegetable production in Florida. EDIS. 2022. doi: 10.32473/edis-cv296-2022
 54. Mahmud MS, Zahid A, Das AK. Sensing and automation technologies for ornamental nursery crop production: current status and future prospects. Sensors. 2023;23:1818. doi: 10.3390/s23041818
 55. Mantzoukas S, Eliopoulos PA. Endophytic entomopathogenic fungi: A valuable biological control tool against plant pests. Appl Sci (Switzerland). 2020;10:360. doi: 10.3390/app10010360
 56. Mariotti B, Martini S, Raddi S, Tani A, Jacobs DF, Olié JA, et al. Coconut coir as a sustainable nursery growing media for seedling production of the ecologically diverse quercus species. Forests. 2020;11:522. doi: 10.3390/f11050522
 57. Mohanta S. Nursery Raising for Vegetables and Flowers in Greenhouse. In: Protected Cultivation and Smart Agriculture. New Delhi, India: New Delhi Publishers; 2020. p. 87-95. doi: 10.30954/ndp-pcsa.2020.9
 58. Moriya EA, Imai NN, Tommaselli AM, Berveglieri A, Honkavaara E, Soares MA, et al. Detecting Citrus Huanglongbing in Brazilian Orchards using Hyperspectral Aerial images. In: The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences. 2019;XLII-2-W13:1881-1886. doi: 10.5194/ISPRS-ARCHIVES-XLII-2-W13-1881-2019
 59. Morrow RC. LED lighting in horticulture. HortScience. 2008;43:1947-1950. doi: 10.21273/HORTSCI.43.7.1947
 60. Mustapha S, Musa AK, Apalowo OA, Lawal AA, Olayiwola OI, Bamidele HO, et al. Open vertical farms: a plausible system in increasing tomato yield and encouraging natural suppression of whiteflies. Acta Agric Slov. 2022;118:1-9. doi: 10.14720/aas.2022.118.2.2272
 61. Nagasubramanian K, Jones S, Singh AK, Sarkar S, Singh A, Ganapathysubramanian B. Plant disease identification using explainable 3D deep learning on hyperspectral images. Plant Methods. 2019;15:1-10. doi: 10.1186/s13007-019-0479-8
 62. Naik MS, Desai S, Sairam KV, Chaitra SN. IoT-based nursery management system. In: Advances in Artificial Intelligence and Data Engineering. AIDE 2019. Advances in Intelligent Systems and Computing. Singapore: Springer; 2021. p. 1335-1344. doi: 10.1007/978-981-15-3514-7_100
 63. Nursyahid A, Setyawan TA, Sa'diyah K, Wardihani ED, Helmy H, Hasan A. Analysis of Deep Water Culture (DWC) hydroponic nutrient solution level control systems. IOP Conf Ser Mater Sci Eng. 2021;1108:012032. doi: 10.1088/1757-899X/1108/1/012032
 64. Patil JA, Yadav S, Kumar A. Management of root-knot nematode, *Meloidogyne incognita* and soil borne fungus, *Fusarium oxysporum* in cucumber using three bioagents under polyhouse conditions. Saudi J Biol Sci. 2021;28:7006-7011. doi: 10.1016/j.sjbs.2021.07.081
 65. Paucek I, Durante E, Pennisi G, Quaini S, Gianquinto G, Orsini F. A methodological tool for sustainability and feasibility assessment of indoor vertical farming with artificial lighting in Africa. Sci Rep. 2023;13:2109. doi: 10.3389/s41598-023-29027-8
 66. Paudel S, Sah LP, Devkota M, Poudyal V, Prasad PV, Reyes MR. Conservation agriculture and integrated pest management practices improve yield and income while

- reducing labor, pests, diseases and chemical pesticide use in smallholder vegetable farms in Nepal. Sustainability (Switzerland). 2020;12:6418. doi: 10.3390/su12166418
67. Peerzada YY, Iqbal M. Leaf Senescence and Ethylene Signaling. In: Aftab T, Hakeem KR, editors. Plant Growth Regulators: Signalling under Stress Conditions. Cham: Springer International Publishing; 2021. p. 153-171. doi: 10.1007/978-3-030-61153-8_7
 68. Pezhman H, Saeidi K. Effectiveness of various solar light traps with and without sex pheromone for mass trapping of tomato leaf miner (*Tuta absoluta*) in a tomato field. Not Sci Biol. 2018;10:475-484. doi: 10.15835/nsb10410303
 69. Podar D, Maathuis FJ. Primary nutrient sensors in plants. iScience. 2022;25:104029. doi: 10.1016/j.isci.2022.104029
 70. Polder G, Blok PM, de Villiers HA, van der Wolf JM, Kamp J. Potato virus Y detection in seed potatoes using deep learning on hyperspectral images. Front Plant Sci. 2019;10:434052. doi: 10.3389/fpls.2019.00209
 71. Prajval V, Krishnamoorthi A, Thakur R, Ruchitha T, Kapoor M, Singh A, Chittibomma K. From code to crop: how bioinformatics is transforming crop genomics in modern agriculture and bettering environment. J Adv Biol Biotechnol. 2024 Mar 16;27(4):27-49.
 72. Reddy GV. Comparative effect of integrated pest management and farmers' standard pest control practice for managing insect pests on cabbage (*Brassica* spp.). Pest Manag Sci. 2011;67:980-985. doi: 10.1002/ps.2142
 73. Redekar NR, Bourret TB, Eberhart JL, Johnson GE, Pitton BJ, Haver DL, et al. The population of oomycetes in a recycled irrigation water system at a horticultural nursery in southern California. Water Res. 2020;183:116050. doi: 10.1016/j.watres.2020.116050
 74. Reed RC, Bradford KJ, Khanday I. Seed germination and vigor: ensuring crop sustainability in a changing climate. Heredity (Edinb). 2022;128:450-459. doi: 10.1038/s41437-022-00497-2
 75. Rimpika, Anushi, Manasa S, Anusha KN, Sharma S, Thakur A, Shilpa, Sood A. An Overview of Precision Farming. Int J Environ Clim Change. 2023;13(12):441-456.
 76. Shi L, He H, Yang G, Huang H, Vasseur L, You M. Are yellow sticky cards and light traps effective on tea green leafhoppers and their predators in Chinese tea plantations? Insects. 2021;12:14. doi: 10.3390/insects12010014
 77. Shimoda M, Honda K. Insect reactions to light and its applications to pest management. Appl Entomol Zool. 2013;48:413-421. doi: 10.1007/s13355-013-0219-x
 78. Shoa P, Hemmat A, Amirfattahi R, Gheysari M. Automatic extraction of canopy and artificial reference temperatures for determination of crop water stress indices by using thermal imaging technique and a fuzzy-based image-processing algorithm. Quant Infrared Thermogr J. 2022;19:85-96. doi: 10.1080/17686733.2020.1819707
 79. Singh B. New Systems of Vegetable Production: Protected Cultivation, Hydroponics, Aeroponics, Vertical, Organic, Microgreens. In: Singh B, Kalia P, editors. Vegetables for Nutrition and Entrepreneurship. Singapore: Springer; 2023. p. 31-54. doi: 10.1007/978-981-19-9016-8_2
 80. Tahiri AI, Meddich A, Raklami A, Alahmad A, Bechtaoui N, Anli M, et al. Assessing the potential role of compost, PGPR and AMF in improving tomato plant growth, yield, fruit quality and water stress tolerance. J Soil Sci Plant Nutr. 2022;22:1-22. doi: 10.1007/s42729-021-00684-w
 81. Talukder B, Blay-Palmer A, vanLoon GW, Hipel KW. Towards complexity of agricultural sustainability assessment: Main issues and concerns. Environ Sustainability Indicators. 2020;6:100038. doi: 10.1016/j.indic.2020.100038
 82. Tzounis A, Katsoulas N, Bartzanas T, Kittas C. Internet of Things in agriculture, recent advances and future challenges. Biosyst Eng. 2017;164:31-48. doi: 10.1016/j.biosystemseng.2017.09.007
 83. Valcke R. Can chlorophyll fluorescence imaging make the invisible visible? Photosynthetica. 2021;59:381-398. doi: 10.32615/ps.2021.017
 84. Valdes VM, Bradford KJ. Effects of seed coating and osmotic priming on the germination of lettuce seeds. J Am Soc Hortic Sci. 2022;112:153-156. doi: 10.21273/jashs.112.1.153
 85. Valero CI, Pla EI, Vaño R, Garro E, Boronat F, Palau CE. Design and development of an aiot architecture for introducing a vessel eta cognitive service in a legacy port management solution. Sensors. 2021;21:8133. doi: 10.3390/s21238133
 86. Vaughn SF, Byars JA, Jackson MA, Peterson SC, Eller FJ. Tomato seed germination and transplant growth in a commercial potting substrate amended with nutrient-preconditioned Eastern red cedar (*Juniperus virginiana* L.) wood biochar. Sci Hortic. 2021;280:109947. doi: 10.1016/j.scienta.2021.109947
 87. Waiba KM, Sharma P, Sharma A, Chadha S, Kaur M. Soil-less vegetable cultivation: A review. J Pharmacogn Phytochem. 2020;9:631-636.
 88. Wang C, Lv J, Coulter JA, Xie J, Yu J, Li J, et al. Slow-release fertilizer improves the growth, quality and nutrient utilization of wintering Chinese chives (*Allium tuberosum* Rottler ex Spreng.). Agronomy. 2020;10:381. doi: 10.3390/agronomy10030381
 89. Wang X, Dou Z, Shi X, Zou C, Liu D, Wang Z, et al. Innovative management programme reduces environmental impacts in Chinese vegetable production. Nat Food. 2021;2:47-53. doi: 10.1038/s43016-020-00199-0
 90. Warner LA, Lamm AJ, Beattie P, White SA, Fisher PR. Identifying opportunities to promote water conservation practices among nursery and greenhouse growers. HortScience. 2018;53:958-962. doi: 10.21273/HORTSCI12906-18
 91. Warner L, Rihn A, Fulcher A, Schexnayder S, LeBude A. Relating grower perceptions and adoption of automated nursery technologies to address labor needs. J Agric Educ. 2022;63:150-168. doi: 10.5032/jae.2022.02150
 92. Yang EM, Kang HM, Kim CS, Pak CH. Dependence of the color appearance of some flowers on illumination. Color Res Appl. 2014;39:28-36. doi: 10.1002/col.21766
 93. Yang P, Wu L, Cheng M, Fan J, Li S, Wang H, et al. Review on drip irrigation: impact on crop yield, quality

- and water productivity in China. *Water* (Switzerland). 2023;15:1733. doi: 10.3390/w15091733
94. Yan T, Xu W, Lin J, Duan L, Gao P, Zhang C, *et al.* Combining multi-dimensional convolutional neural network (CNN) with visualization method for detection of *Aphis gossypii* Glover infection in cotton leaves using hyperspectral imaging. *Front Plant Sci.* 2021;12:604510. doi: 10.3389/fpls.2021.604510
95. Zhang H, Asutosh A, Hu W. Implementing vertical farming at university scale to promote sustainable communities: A feasibility analysis. *Sustainability* (Switzerland). 2018;10:4429. doi: 10.3390/su10124429
96. Zhang K, Khan Z, Wu H, Khan MN, Hu L. Gibberellic acid priming improved rapeseed drought tolerance by modulating root morphology, ROS homeostasis and chloroplast autophagy. *J Plant Growth Regul.* 2022;42:5977-5990. doi: 10.1007/s00344-022-10718-x
97. Zhang R, Xu C, Bao Z, Xiao R, Chen X, Xiao W, *et al.* Auxin alters sodium ion accumulation and nutrient accumulation by playing protective role in salinity challenged strawberry. *Plant Physiol Biochem.* 2021;164:1-9. doi: 10.1016/j.plaphy.2021.04.008
98. Zhao L, Gong L, Lv C, Li Q, Li B, Yang M, *et al.* Effects of different artificial light sources on mechanical properties of closed stereoscopic rice seedlings. *Adv Eng Technol Res.* 2023;4:171-171. doi: 10.56028/aetr.4.1.171.2023
99. Zhao M, Liao H, Molokeev MS, Zhou Y, Zhang Q, Liu Q, *et al.* Emerging ultra-narrow-band cyan-emitting phosphor for white LEDs with enhanced color rendition. *Light: Sci Appl.* 2019;8:1-9. doi: 10.1038/s41377-019-0148-8