



International Journal of Research in Agronomy

E-ISSN: 2618-0618
P-ISSN: 2618-060X
© Agronomy
NAAS Rating (2026): 5.20
www.agronomyjournals.com
2026; SP-9(5): 505-511
Received: 14-03-2026
Accepted: 18-04-2026

Sandeep Yadav
Department of Physics, CCSHAU
College of Agriculture Bawal,
Haryana, India

Mohinder Singh
Department of Agronomy,
CCSHAU College of Agriculture
Bawal, India

Babita Rani
Department of Biochemistry,
CCSHAU College of Agriculture
Bawal, Haryana, India

Jitender Kumar
Department of Agronomy,
CCSHAU Regional Research
Station Bawal, Haryana, India

Renu Bala
Lecturer Physics, Government
Girls Sr. Sec. School, Bawani
Khera, Bhiwani, Haryana, India

Black carbon to green prosperity: Biochar innovation for climate mitigation, soil health, and sustainable agriculture

Sandeep Yadav, Mohinder Singh, Babita Rani, Jitender Kumar and Renu Bala

DOI: <https://www.doi.org/10.33545/2618060X.2026.v9.i5Sf.5660>

Abstract

The availability of agricultural residues in India is large which is more than 600 million tonnes annually having immense potential for biomass derived carbon materials production. The application of this carbon enriched material in the soil is considered highly attractive as well as sustainable strategy that enhances carbon sequestration in soil and to mitigate climate change. It is one of ancient practice whose traces are found in terra-preta soils in Amazon. Our comprehensive literature analysis shows a historical perspective to modern initiative taken by Google and VARAHA to tackle the emissions of poisonous gases, stubble burning as well as soil degradation that engaged 15, 000+ farmers, yielding 10–25% crop boosts and 30–50% N₂O cuts. Integrating into 2026 carbon markets could sequester 0.1 Gt CO₂ yearly, creating rural jobs via decentralized kilns in states like Rajasthan and Madhya Pradesh.

Keywords: Biochar, pyrolysis, sustainable agriculture

Introduction

The agricultural waste, forest residues, or even household organic scraps are heated in a low-oxygen environment instead of rot or burn. The resulting black color porous material buried deep into the soil. The material emerges is not just a charcoal but it is biochar. It is a modern twist on an ancient practice that could help turn the tide against climate change. For centuries, Indigenous peoples in the Amazon created fertile "terra preta" soils by intentionally incorporating charcoal into the ground, transforming nutrient-poor tropical earth into remarkably productive land that still holds rich carbon stores today. Now, scientists, farmers, and climate experts are rediscovering this technique on a global scale [1-14]. Biochar isn't just about enriching soil—it's one of the most promising, scalable tools we have for pulling carbon dioxide out of the atmosphere and locking it away for hundreds to thousands of years [15, 16].

Biochar stands out as a very feasible answer in current scenario when every gigaton of CO₂ reduced counts toward preventing calamitous warming. It is produced by the process by heats biomass without oxygen known as pyrolysis. It converts temporary, cycle-bound carbon into a stable, recalcitrant form that resists breakdown. Researchers reports a widespread adoption may sequester anywhere from 0.1-2 gigatons of CO₂ equivalent per year. This has wide range of benefits such as healthier soils that retain more water and nutrients, higher crop yields in degraded lands, reduced greenhouse gas emissions from fields, and even improved resilience against drought and flooding [8-14, 17-21]. Despite of its long history and innovative potential, biochar is quite neglected. The concept of burying carbon to save planet has evolved from fringe thought to real competitor in global climatic toolbox whereas one knows factors such as wildfire explode, soil deterioration and continuously elevation of CO₂ levels.

It is specially engineered form of charcoal which is designed not for burning of biomass but for burying in the soil to fight climate change and boost soil health. So biochar is a carbon-rich, stable, porous material synthesized by heating organic biomass such as crop residues (straw, husks), wood scraps, forest thinning, animal manure and food waste in the absence of oxygen through a process known by pyrolysis. During pyrolysis biomass is heated up to temperatures

Corresponding Author:
Sandeep Yadav
Department of Physics, CCSHAU
College of Agriculture Bawal,
Haryana, India

300–700 °C in a low-oxygen or oxygen-free environment. The volatile compounds (gases and oils) escape, leaving behind a solid charcoal-like residue which is rich in carbon. The biochar is highly resistant to decay which locks the carbon which plants

originally takeout from the atmosphere during the process of photosynthesis. The key difference in Biochar and regular charcoal are in table-1.

Table 1: The key difference in biochar and regular charcoal

Aspect	Biochar	Charcoal
Primary Use	Soil amendment, carbon sequestration, agriculture ^[3-6]	Fuel for cooking, heating, manufacturing ^[3-6]
Production Temperature	Higher (450-650°C or 840-1200°F) for stability ^[3-6]	Lower (~400 °C or 750°F) ^[3-6]
Feedstock	Diverse biomass (plants, waste, manure) ^[3-6]	Mainly wood (e.g., hardwood like oak) ^[3-6]
Porosity/Surface Area	Higher porosity, larger surface area for nutrient/water retention ^[3-6]	Lower porosity, less stable in soil ^[3-6, 22, 23]
Stability/pH	More stable (centuries in soil); pH 4.6-9.3 ^[3-6]	Less stable, breaks down faster; pH 8-11 ^[3-6]
Environmental Impact	Improves soil health, sequesters carbon ^[3-6]	Can release pollutants if poorly produced ^[3-6]




Fig 1: The Pyrolysis Process: A Quick Overview

1. Feedstock preparation — Biomass (e.g., crop waste, wood chips) is dried and in some cases, it is grounded.
2. Heating in pyrolysis reactor — In this process material enters into a sealed chamber/kiln heated to 300–700 °C with some amount of oxygen to no oxygen.
3. Decomposition — Volatiles release as syngas (for energy) and bio-oil (potential fuel/chemicals); a solid carbon remains in the end as a residue.
4. Cooling and collection — Biochar is quenched or cooled. Sometimes it is "activated" or charged with some nutrients for better soil health performance.
5. Application — The final product obtained is well mixed into the soil.

This whole process as shown in fig 1 turns the temporary plant carbon into a durable form along with some by-products that provides renewable energy. This is suitable way to both climate and waste management.

Special properties of biochar: Biochar is special due to following properties such as:

1. **High surface area:** Its surface area is usually 100–500 m²/g or more which is equivalent to several football fields in a single teaspoon. This also shows huge internal porosity that can be visualized under scanning electron microscopes (SEM). The structure seen look like a honeycomb as shown in fig. 2 which may be suitable habitat for beneficial microbes as well as it acts like a sponge for both water and nutrients.




Fig 2: Honeycomb Structure of biochar (carbongold.com)

2. **Exceptional stability:** In biochar carbon is highly volatile and hardly bounded and, resisting to microbial breakdown. Amazonian terra preta as an ancient example shows that it can persist in soil for centuries to millennia^[24]. Which provide long-term carbon sequestration in environment.
3. **pH effects:** The nature of biochar is mostly alkaline (pH 7.5–11) in nature that acts as a natural amendment agent to improve soil pH in acidic conditions. This improves phosphorus and calcium availability and reduces aluminum toxicity in low-pH soils. However, pH varies by parent material and temperature regimes eg., lower-temperature biochar can be near-neutral or slightly acidic^[25].

These multifaceted traits make biochar far more material for soil stability than charcoal. This make biochar as a carbon sink for carbon sequestration, soil conditioner, water retainer in capillary tubes, and nutrient mineralization from organic wastes that otherwise be lost by volatilization, leaching or burning. In the current scenario of land degradation and climate change the incorporation of biochar is the simplest and most powerful way to harvest atmospheric CO₂ and also plays an important role in improving flora and fauna in degraded lands.

Path From Ancient Wisdom to Modern Science

Deep in the Amazon rainforest, patches of extraordinarily dark, fertile soil have puzzled observers for generations. Known locally as terra preta—"black earth" in Portuguese—these soils stand in stark contrast to the typically nutrient-poor, highly weathered tropical earth surrounding them. While adjacent lands quickly lose fertility under heavy rains and intense sun, terra preta plots remain productive, often yielding double the crops even centuries after abandonment. Covering an estimated 6, 000 to 18, 000 square kilometers (and possibly more) across the Amazon Basin, these enigmatic soils contain up to three times more phosphorus and nitrogen, and dramatically higher organic carbon which is usually 9% as compared to 1% in nearby soils^[24]

Archaeological evidence points to Indigenous Amazonian communities as the creators. Between roughly 450 BCE and 950 CE (In some sites carbon dating back over 2, 500 years or more), pre-Columbian farmers practiced sophisticated land management. They deliberately incorporated charcoal which was produced from burning organic materials such as wood, crop residues and household waste under low-oxygen conditions. They also added broken pottery shards, animal bones, manure, fish remains and other debris into garden plots ranging from small household plots to larger areas in several hectares. This "slash-and-char" approach (as opposed to slash-and-burn) transformed temporary biomass carbon into a stable, porous form that resisted rapid decomposition. As a result of

these soils enriched with nutrients, a better water retention, increase in microbial communities and extraordinary resilience against leaching and erosion. It was fully intentional or a byproduct of settlement waste but these practices supported dense, settled populations in what Europeans once dismissed as inhospitable wilderness. European settlers come upon terra preta in the 19th century, as per explorers like Charles Hartt in 1885 who revealed the unusually dark, fertile patches of land. But irony is that a systematic scientific interest was not there until the mid of 20th century. Wim Sombroek who was a Dutch soil scientist documented these soils in detail in his influential 1966 book *Amazon Soils* [19, 26, 27]. His work claimed the anthropogenic origins and sparked curiosity about the "black carbon" responsible for their long life.

A bridge between ancient practice and modern climate science was appeared in the late 1990s and early 2000s. Johannes Lehmann at Cornell University who was soil biogeochemist known as known figure after his work on study of nutrient and carbon dynamics in the central Amazon. With inspiration from terra preta's stability, Lehmann and his collaborators like Bruno Glaser analyzed the pyrogenic carbon which can be said as charcoal-like material in these soils. Lehmann (2003) reported that charcoal reduce nutrient leaching in soil and boost crop growth and act as a major tool for carbon sequestration to mitigate climate change [19, 28]. In the mid-2000s the "biochar"

(short for biomass-derived charcoal used as a soil amendment) gained attraction for scientific studies. The controlled experiments started by several scientists worldwide by replicating terra preta principles. Their studies confirmed that properly produced biochar sequesters carbon long-term and 80–90% persists even after 100 years in models based on H/C ratios enhancing soil fertility in degraded lands as well as a efficient tool to combat climate change. They tailor biochar for specific soils and climates from different feedstocks. A blend of old practices with modern cutting-edge soil science biochar emerges as most powerful tools for our planet's future that may be buried in the past. Now it's time to adapt the new and improved form of that old practice.

The Production Puzzle: How It's Made Today Synthesis Methods of Biochar

Biochar is synthesized by the method of pyrolysis that involves heating of the biomass in an anaerobic or oxygen-limited environment. It leads to thermal decomposition of material. This technique is good for production of high-quality biochar for soil applications as well as carbon sequestration. Pyrolysis depends upon the on temperature, heating rate and residence time which are shown in table-2 below. These techniques transform temporary biomass carbon into durable, buried forms that makes biochar a practical tool in the fight against climate change.

Table-2: Comparison of different Pyrolysis temperature process [25, 29-37]

Parameters	Slow Pyrolysis	Fast Pyrolysis	Flash/Ultra-Fast Pyrolysis
Temperature	300–700 °C	400–600 °C	>800 °C
Heating rate	Very low (0.1–1 °C/s)	High (10–200 °C/s)	Extremely high (>1000 °C/s)
Residence time	Long (minutes to hours or days)	Very short (seconds)	Milliseconds
Biochar yield	High (25–50%, sometimes up to 80%)	Lower (10–25%)	Very low (<10–15%)
Advantages	Maximizes solid biochar production with good carbon stability; produces balanced amounts of syngas and bio-oil. Ideal for soil amendment and long-term carbon storage.	Optimized for bio-oil production (up to 60–75%); biochar is a by-product with finer particles and higher surface area.	Focuses on rapid vaporization for maximum liquid yields.
Common reactors	Kilns, retorts, rotary kilns, or simple earth mounds (traditional methods)	Fluidized beds, ablative reactors	

Now a days biochar production method is inspired from ancient kilns to cutting-edge engineering. This turns organic solid waste into environment-friendly product. This is a thermochemical process in which biomass is heated without (or with very limited) oxygen to take out volatile compounds and carbon-rich solids remains as residue. It is observed that slow pyrolysis produces high-quality biochar. It is very much suitable for durable carbon sequestration and soil applications. In the slow pyrolysis process the given biomass is heated gradually at 300–700 °C with low heating rates (0.1–1 °C/s). The residence times is kept long which may be from minutes to hours. In the slow pyrolysis biochar yield is higher which is usually 25–50%. It creates a stable, porous structure that is ideal for locking carbon away for centuries. The method produces balanced by-products: syngas for energy and bio-oil for fuels or chemicals. This is very much suitable for dry to moist feedstocks. But for the wet biomass such as food waste, sewage sludge, algae or fresh manure hydrothermal carbonization (HTC) is most suitable. This method is also known as hydro-char production or wet pyrolysis. This method requires lower temperatures (180–250 °C) and no pre-drying is needed which makes it energy-efficient and cost-effective method. HTC's product is more acidic and less aromatic than pyrolytic biochar but excels for nutrient-rich or contaminated wastes.

For the farmers in Bharat on-site crop waste conversion to biochar a low-tech side, traditional and semi-traditional methods

needed for small-scale operations. These must require minimal infrastructure, affordable and produce biochar from local agricultural residues. A decent quality biochar is produced with minimize emissions. These are low-tech kilns such as earth mounds, flame-curtain designs, Kon-Tiki cone kilns or brick retorts which rely on manual or semi-manual operation. They're ideal. For the fast scale production there are fluidized beds, continuous pyrolysis units and rotary kilns process tons per hour. The common sources for biochar include agricultural residues such as rice husks, wheat straw, maize stalks, mustard stalks, cotton stalks, forestry slash, pruned branches, municipal green waste, animal manures and even industrial by-products from paper mill sludge and spent coffee grounds [34]. Using these avoids land-use competition and turns disposal problems into resources that reduces methane from landfills or open burning while creating value.

Scalability of biochar remains the perplex process. Low-tech approaches validate scattered community-level production, while industrial systems enable mass potential efficiency and volume. Innovation in microwave, solar pyrolysis, including better emissions technology helps in reducing barriers. As per projected market needs to expand scaling process upto 2030s and the production scenario is shifting from "how" to "how fast quality produce" with waste-to-biochar production pathways offering a versatile and practical route to cope up climate management.

How Biochar Fights Climate Change

The biochar mitigates climate change by reversing the carbon cycle. We know that plants naturally capture carbon dioxide (CO₂) from the atmosphere by photosynthesis process. This incorporates carbon into their tissues. When plants die, they decomposed or burned then they release most of that carbon back as CO₂ or methane (CH₄) into atmosphere. Here the role of biochar comes and it interrupts this short, rapid loop. During pyrolysis of biomass in low-oxygen conditions approximately 50% or more of the original carbon is locked. This locked form is highly stable, aromatic structure which resists microbial breakdown for centuries to millennia when mixed in to the soil suggesting multi-century stability based on ancient terra preta examples and lab aging tests. It transforms short lived biomass carbon into a durable sink by removing CO₂ from the atmosphere for the long haul as a direct carbon sequestration. In recent independent assessments of carbon sequestration are around 2.6–2.7 gigatons (Gt) of CO₂-equivalent per year when using waste biomass and residues, avoiding competition with food production or deforestation [37]. Technically in broader prospects that includes saved emissions as well as bioenergy can reach up to 10.3 Gt CO₂/yr but a realistic sustainable deployment may be near to 0.5–2.5 Gt CO₂/yr range for focused carbon dioxide removal (CDR). Whereas the global fossil fuel emissions are around 35–40 Gt CO₂ annually. If scaled thoughtfully then even in the lower end could counterbalances a meaningful fraction which is 1–7%.

Along with the carbon sequestration biochar also delivers powerful indirect mitigation to soil greenhouse gas emissions. As agricultural soils are major sources of nitrous oxide (N₂O) which is a potential factor in Green House Gases (GHG). It creates 265 times more warming than CO₂ over 100 years.

Investigation reveals that biochar steadily cuts N₂O emissions ranges 18–54% on average, with reductions up to 33–39% in many field trials. Mechanisms for this reduction includes:

- Adsorption of ammonium and nitrate that limits substrate for nitrification and denitrification.
- An improved soil aeration as well as soil pH shifts that favor complete denitrification to harmless N₂ over N₂O.
- Suppresses some of key microbial enzymes which are involved in GHG production.

The reduction in CH₄ is variable but in CO₂ it is interesting to note that net effects are typically neutral to positive when combined with other amendments. During biochar production the generation of syngas and bio-oil which can displace use of fossil fuels for heat, power, or chemicals that counterbalances the emissions from these elsewhere. Although biochar by-products added up to overall negative emissions.

Biochar settles CO₂ in gigaton-scale that offers CDR solution. It is currently sequestering 0.2–0.6 Mt CO₂/yr approximately. To meet out a 1.5°C goals by 2050 requires 5–16 Gt/yr for the scaling up the carbon pricing along with policy support [17, 38]. Along with the enhancement of agricultural resilience it serves as a pragmatic, multi-solving tool for reforestation efforts and reduction in emission.

Comparison of Biochar Impact by Soil Type

A comparative representation is shown in the table-3 for biochar impact on different type of soil. This will help to visualize how biochar performs across different environments and where biochar can it acts as a "hero" and where it might be a "hindrance."

Table 3: Comparison of Biochar Impact by Soil Type

Soil Type	Primary Benefit	Potential Risk/Roadblock	Recommendation
Acidic/Tropical	Neutralizes pH; improves cation exchange capacity (CEC).	High transport costs to remote regions.	Use biochar with high liming potential.
Sandy/Arid	Increases water retention; prevents nutrient leaching.	Can be blown away by wind if not incorporated quickly.	Apply "charged" biochar (pre-soaked in liquid fertilizer).
Heavy Clay	Improves aeration and drainage (porosity).	May have negligible yield impact in the short term.	Use coarse-grained biochar to break up compaction.
Fertile/Alkaline	Long-term carbon sequestration.	Can raise pH too high, causing "nutrient lockout."	Use acidified biochar or lower application rates.

Biochar and Agriculture Future

Biochar has a wide range of benefits in agriculture such as soil amendment, contributes to sustainable, climate-smart, and resource-efficient farming systems. When biochar on incorporation in fields it interacts with physical, chemical, and biological processes, often leading to long-term improvements in productivity and environmental quality.

Soil physical improvements

Biochar improves the soil structure and tilth by increasing the pore space and stabilising the soil aggregates. The carbon rich and porous matrix of biochar reduces compaction in soil that improves aeration which is very important for heavy clay or degraded soils. A better physical environment encourages stronger root development as well as deeper root penetration in soil. This helps in efficient water uptake and nutrient absorption in plants. Micropores of biochar behaves like a sponge which increase the ability of soil to retain the moisture and release slowly to the roots. This slow release of water reduces the frequency of irrigation and water loss through evaporation as

well as drainage. It becomes useful for rain-fed agriculture in arid region and in the periods of stress to erratic rainfall or drought [39–41].

Chemical buffering and nutrient management

Biochar is alkaline in nature most of time that can help buffer acidic soil and help to reduce aluminum toxicity in tropical and subtropical soils. After the consecutive use this can lower the need of large lime applications as well as improve the availability of pH-sensitive nutrients like phosphorus and molybdenum. Biochar has strong cation exchange capacity (CEC) due to its high porous structure, high surface area and the presence of oxygen-containing functional groups viz. carboxyl, hydroxyl, and phenolic groups which are developed during pyrolysis. High CEC makes it to adsorb and retain essential nutrients like magnesium, potassium, calcium and ammonium. It reduces leaching losses to groundwater that allows the nutrients to be released slowly as plants needed. As a result, this reduces the nutrient runoff and improves efficiency of fertilization when combined with compost [34, 41–43].

Microbial habitat and soil biology

The porous structure of biochar creates a network of micro-habitats that shelter beneficial soil microorganisms, including bacteria, fungi, and beneficial microbes involved in nitrogen fixation and phosphorus solubilization. Enhanced microbial activity accelerates the decomposition of organic matter, improves nutrient cycling, and produces plant-growth-promoting compounds such as hormones and enzymes [15, 41, 44].

In some cases, biochar-rich environments can favor the growth of mycorrhizal fungi, which form symbiotic associations with plant roots and greatly extend the plant's effective root surface area. This leads to improved uptake of water, phosphorus, and other poorly mobile nutrients, often resulting in stronger, more vigorous plants and better yields, especially in nutrient-poor or degraded soils.

Contamination mitigation and stress resilience

Biochar can adsorb and immobilize certain heavy metals (such as Cd, Cu, Pb and some forms of arsenic) and organic pollutants by reduction in their bioavailability to plants. This lowers the risk of food-chain contamination. This can be part of a remediation strategy in area where soil is heavily contaminated due to industrialisation. As biochar improves water retention, soil structure and nutrient availability so biochar also enhances plant resilience to abiotic stresses. These includes drought, salinity, and temperature extremes. In saline soils, for example, biochar can help condition the rhizosphere so that plants experience less osmotic stress and ion toxicity, supporting better growth under marginal conditions [25, 37].

Climate-change mitigation and carbon sequestration

Because biochar is composed largely of stable aromatic carbon, it can persist in soils for decades to centuries, effectively locking away carbon that would otherwise return to the atmosphere through decomposition or burning of biomass. This long-term carbon storage contributes to climate-change mitigation by reducing net greenhouse-gas emissions from agricultural systems [17]. In addition to sequestering carbon, biochar can suppress emissions of nitrous oxide (N_2O) from soils high in nitrogen, likely by altering microbial activity and nitrogen cycling pathways. Reduced N_2O emissions, combined with lower fossil-fuel inputs in some cases (due to improved fertilizer efficiency), make biochar an important tool for climate-smart agriculture [38].

Waste valorization and circular-agriculture opportunities

Biochar can be produced from agricultural residues (rice husk, straw, sugarcane bagasse, crop pruning waste), forestry residues, municipal organic waste, and even certain manures. Instead of burning or dumping these materials, converting them to biochar turns low-value biomass into a high-value soil enhancer.

This approach supports circular-agriculture models by closing nutrient loops, reducing open-field burning (and associated air-quality problems), and generating a useful co-product from biomass-energy processes. In many small-holder and rural settings, on-farm biochar systems can be low-cost, renewable, and compatible with traditional farming practices, especially when integrated with compost or manure [2, 3, 8].

Crop productivity and yield stability

Numerous field trials and meta-analyses show that biochar application can increase crop yields, particularly in infertile, sandy, or highly weathered soils where nutrient retention and

water-holding capacity are weak. Effects are often more pronounced in tropical and subtropical regions, but benefits are also observed in temperate cropping systems, especially when biochar is combined with good agronomic practices [32, 42].

Over time, biochar can contribute to more stable yields by buffering soil against extreme fluctuations in moisture and nutrient availability. This stability is valuable for food-security programs and for farmers who face increasing climate variability and input-cost pressures. In some systems, biochar-based amendments are being explored as part of conservation-agriculture or regenerative-agriculture packages that emphasize minimal soil disturbance, permanent soil cover, and diverse cropping [4].

Customization and integration with other technologies

Biochar properties—such as surface area, pH, nutrient content, and porosity—depend on the feedstock (rice husk, wood, crop residues, etc.) and production conditions (temperature, heating rate, residence time). This allows for tailored biochar products suited to specific soil types, crops, and management goals [10]. Biochar can be combined with other sustainable inputs, such as compost, microbial inoculants, and slow-release fertilizers, to create “designer” soil amendments that optimize nutrient delivery and biological activity. In precision-agriculture frameworks, biochar-based formulations can be applied in targeted zones (e.g., planting rows, seed beds) to maximize plant-root interactions while minimizing material use and environmental trade-offs [31, 34].

Hence biochar in agriculture offers a multifaceted advantages like improvement in physical and chemical conditions of soil, enhances microbial life, mitigates contamination and climate-relevant emissions, valorizes waste biomass, and supports more resilient and productive farming systems. When properly matched to local soils, crops, and management practices, biochar can be a powerful, long-term tool for sustainable intensification of agriculture.

Success Stories and the Indian Context

India has generated 600 million metric tons annual agricultural residue, so it has full potential of emerging as a global venture for biochar at international scaling. By 2025, approximately 15000 Indian farmers have already adopted biochar in production that reported 10-25% yield increment along with 30–50% reduction in nitrous oxide greenhouse gas emissions, which has 273 times more potential than CO_2 . During the 2024–2025, Rajasthan and Maharashtra started a pilot project of applying 1.2 mt of biochar per acre that have showed 15% improvement crop emergence and 40% reduction in chemical fertilizer requirement. In 2024, Madhya Pradesh and Odisha nearly 1, 000 households used biochar application and improved 29.34% in their net farm profitability along with 1.5% reduction in crop production costs. In recent years, biochar has dominated the physical Carbon Dioxide Removal (CDR) market, often accounting for over 90% of all delivered carbon credits globally. In 2025, a landmark deal, Google signed industrial biochar agreement with Indian startup Varaha [45], is the largest land mark deal of purchasing 100, 000 tons of carbon removal.

Moving Forward: Overcoming the Roadblocks

To overcome these challenges, the industrial production must be shifting toward special needs of variable soil chemistry that may be called as “designer biochars”. So, the next generation biochar is engineered Biochar not just burned biomass. This specialized biochar is treated as accurate tool rather than a general

commodity to reduce economic and environmental risks significantly reduced. The biochar is transitioning strategy amendment to remove a global carbon.

Future innovations areas

Precision Design: Scientists, researchers and environmentalists are now working with AI tools and machine learning to tailoring biochars for specific purposes such as locking away heavy metals toxicity in soils or improve nutrient release as per crop needs.

Nanocomposites: Innovations like biochar-nanoparticle hybrids (using silica or chitosan) are emerging to enhance plant growth and nickel uptake in degraded industrial lands. These nano-biochar (NBC) are emerging as cost-efficient, highly effective, amendments for soil reclamation and remediation specifically for capturing pollutants and contaminants ^[2].

Industrial Integration: The next generation of biochar is not just "burned biomass"; it is Engineered Biochar. These materials are scaled as an industrial adsorbent that improve surface, soil porosity and capable of capturing CO₂ Incorporating just 2–5% biochar into concrete can improve its strength while turning buildings into permanent carbon sinks, sequestering roughly 115 kg of CO₂ per cubic meter.

Regenerative Agriculture and Farmers Participation: Integration of biochar acts as a stable "biological battery" that improve the efficiency of deep nutrient placement ^[1]. Farmers must adopt in-situ waste management, which can increase nearly 5 g/kg in one organic matter ^[2].

For Researcher and Policymakers: researcher must focus new biochar-based recommendation for improving the sustainability of agriculture in country. Government have to start village-level support programs from lab to field and subsidy provision for the farmers to enhance adoption of biochar ^[1].

Conclusion

Biochar production from abundant agricultural waste via pyrolysis is ancient practice that represents a scalable, low-cost strategy to combat climate change by locking away gigatons of CO₂ while enhancing soil fertility and reducing emissions like N₂O by 30-50%. Initiatives like Google's partnership with Varaha, engaging thousands of farmers for yield boosts and carbon credits, demonstrate practical viability in India. Embracing ancient terra preta wisdom with modern innovations such as designer biochar and nano-hybrids can drive rural jobs, integrate into carbon markets by 2026 and foster resilient agriculture against degradation and extremes.

References

1. Al-Amin HM, Rahman MM, Alam MS, Smith J, Islam M, Rees RM, *et al.* Village level fertilizer management for increasing nitrogen use efficiency, rice yield and household income. *Archives of Agronomy and Soil Science*. 2025;71(1):1-19.
2. Pandian K, Vijayakumar S, Mustaffa MRAF, Subramanian P, Chitraputhirapillai S. Biochar – a sustainable soil conditioner for improving soil health, crop production and environment under changing climate: a review. *Frontiers in Soil Science*. 2024;4:1376159.
3. Shapiro. Biochar vs. charcoal: key differences for sustainability. 2026.
4. Rosy Soil. Biochar vs charcoal: 6 key differences you should know about. 2022.
5. Haiqi Environmental Technology. Differences between charcoal and biochar. 2024.
6. Waste X. Biochar vs charcoal: 5 key differences. 2024.
7. Celignis Biomass Laboratory. Biochar as a tool for carbon sequestration. 2026.
8. Gui X, Xu X, Zhang Z, Hu L, Huang W, Zhao L, *et al.* Biochar-amended soil can further sorb atmospheric CO₂ for more carbon sequestration. *Communications Earth & Environment*. 2025;6:5.
9. Zhang T, Tang Y, Li H, Hu W, Cheng J, *et al.* A bibliometric review of biochar for soil carbon sequestration and mitigation from 2001 to 2020. *Ecotoxicology and Environmental Safety*. 2023;264:115438.
10. Biochar in global carbon cycle: towards sustainable agriculture. *Current Research in Green and Sustainable Chemistry*. 2024.
11. Use of biomass-derived biochar for carbon sequestration in soil: recent advancements. *Communications Earth & Environment*. 2025.
12. International Biochar Initiative. Biochar production technologies.
13. Biochar: an emerging soil amendment for sustaining soil health and black gold for Indian agriculture. *Journal of Experimental Agriculture International*.
14. Farmbrite. Biochar and Terra Preta: ancient soil wisdom for modern farmers.
15. Abdul-Aziz AL, Uz I. Biochar from agricultural waste enhances rhizosphere microbial activity over bulk soil. *Journal of Soil Science and Plant Nutrition*. 2026;26:2614-2627.
16. Glaser B, Lehmann J, Zech W. Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal—A review. *Biology and Fertility of Soils*. 2002;35(4):219-230.
17. Waheed A, Xu H, Qiao X, Aili A, Yiremaikeybayi Y, Haitao D, *et al.* Biochar in sustainable agriculture and climate mitigation: mechanisms, challenges, and applications in the circular bioeconomy. *Biomass and Bioenergy*. 2025;193:107531.
18. TechCrunch. Google strikes world's largest biochar carbon removal deal with Indian startup Varaha. 2025.
19. Lehmann J. Integrated biochar systems for soil fertility management. Ithaca (NY): Cornell University; 2012.
20. Shoudho KN, Khan TH, Ara UR, Khan MR, Shawon ZBZ, Hoque ME. Biochar in global carbon cycle: towards sustainable development goals. *Current Research in Green and Sustainable Chemistry*. 2024;8:100409.
21. Waheed A, Xu H, Qiao X, Aili A, Yiremaikeybayi Y, Haitao D, Muhammad M. Biochar in sustainable agriculture and climate mitigation: mechanisms, challenges, and applications in the circular bioeconomy. *Biomass and Bioenergy*. 2025;193:107531.
22. Ali A, Jabeen N, Chachar Z, Chachar S, Ahmed S, Ahmed N, *et al.* The role of biochar in enhancing soil health and interactions with rhizosphere properties and enzyme activities in organic fertilizer substitution. *Frontiers in Plant Science*. 2025;16:1595208.
23. Zhang L, Wu Z, Zhou J, Zhou L, Lu Y, Xiang Y, *et al.* Meta-analysis of the response of the productivity of different crops to parameters and processes in soil nitrogen cycle under biochar addition. *Agronomy*. 2022;12(8):1857.
24. Pandian K, Vijayakumar S, Mustaffa MRAF, Subramanian

- P, Chitraputhirapillai S. Biochar—A sustainable soil conditioner for improving soil health, crop production and environment under changing climate: a review. *Frontiers in Soil Science*. 2024;4:1376159.
25. Vaz RCA, Lopez MAR, Ferreira GMD. Unlocking the potential of MOF-biochar composites: advanced functional materials for adsorption, catalysis, and energy storage. *Materials Today Chemistry*. 2024;42:102383.
 26. Sombroek WG. Amazon soils. Wageningen: Centre for Agricultural Publications and Documentation; 1966.
 27. Brock J. Biochar and Terra Preta: ancient soil wisdom for modern farmers. *Farmbrite*. 2026 Jan 8.
 28. Ayaz M, Feizienė D, Tilvikienė V, Akhtar K, Stulpinaitė U, Iqbal R. Biochar role in the sustainability of agriculture and environment. *Sustainability*. 2021;13(3):1330.
 29. Fuchs M, Garcia-Perez M, Small P, Flora G. Campfire lessons: breaking down the combustion process to understand biochar production. *Biochar Journal*. 2014. ISSN 2297-1114.
 30. Tomczyk A, Sokołowska Z, Boguta P. Biochar physicochemical properties: pyrolysis temperature and feedstock kind effects. *Reviews in Environmental Science and Biotechnology*. 2020;19:191-215.
 31. Li S, Tasnady D. Biochar for soil carbon sequestration: current knowledge, mechanisms, and future perspectives. *C*. 2023;9(3):67.
 32. Joseph S, Cowie AL, Van Zwieten L, Bolan N, Budai A, Buss W, *et al*. How biochar works, and when it doesn't: a review of mechanisms controlling soil and plant responses to biochar. *GCB Bioenergy*. 2021;13:1731-1764.
 33. Frimpong Manso E, Barnor MT, Baba ME, Anyomi WE. Biochar-compost from cashew apple residue as a soil amendment for cashew cultivation in Ghana. *Plant-Environment Interactions*. 2026;7(2):e70138.
 34. Khare P, Deshmukh Y, Yadav V, Pandey V, Singh A, Verma K. Biochar production: a sustainable solution for crop residue burning and related environmental issues. *Environmental Progress & Sustainable Energy*. 2021;40:e13529.
 35. Khan S, Irshad S, Mehmood K, Hasnain Z, Nawaz M, Rais A, *et al*. Biochar production and characteristics, its impacts on soil health, crop production, and yield enhancement: a review. *Plants*. 2024;13(2):166.
 36. Lehmann J, Gaunt J, Rondon M. Bio-char sequestration in terrestrial ecosystems—a review. *Mitigation and Adaptation Strategies for Global Change*. 2006;11(2):403-427.
 37. Li S, Tasnady D. Biochar for soil carbon sequestration: current knowledge, mechanisms, and future perspectives. *C*. 2023;9(3):67.
 38. Kammann C, Ippolito J, Hagemann N, Borchard N, Cayuela ML, Estavillo JM, *et al*. Biochar as a tool to reduce the agricultural greenhouse-gas burden – knowns, unknowns and future research needs. *Journal of Environmental Engineering and Landscape Management*. 2018;25(2):114-139.
 39. Abdul-Aziz AL, Abukari IA, Galadima MM, Haruna A, Abubakri M, Abdulai R. Biochar effects on soil properties and yield of maize in Northern region, Ghana. *Discover Agriculture*. 2025;3:103.
 40. Debab A, Boudjabi S, Chenchouni H, Ababsa N, Brahimi A. Effects of incorporating biochar on soil quality and barley yield in microplastics-contaminated soils. *Chemosphere*. 2024;368:143760.
 41. Xu Z, Zhou R, Xu G. Global analysis on potential effects of biochar on crop yields and soil quality. *Soil Ecology Letters*. 2025;7:240267.
 42. Roberts KG, Gloy BA, Joseph S, Scott NR, Lehmann J. Limestone amendment and biochar to improve water retention. *Environmental Science & Technology*. 2010;44(4):1161-1166.
 43. Banik C, Bakshi S, Laird DA, Smith RG, Brown RC. Impact of biochar-based slow-release N-fertilizers on maize growth and nitrogen recovery efficiency. *Journal of Environmental Quality*. 2023;52(3):630-640.
 44. Hu W, Zhang Y, Rong X, Zhou X, Fei J, Peng J, *et al*. Biochar and organic fertilizer applications enhance soil functional microbial abundance and agroecosystem multifunctionality. *Biochar*. 2024;6:3.
 45. Singh J. Google strikes world's largest biochar carbon removal deal with Indian startup Varaha. *TechCrunch*. 2025 Jan 16.