


International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
NAAS Rating (2026): 5.29
IJABR 2026; 10(6): 418-423
www.biochemjournal.com
Received: 25-03-2026
Accepted: 28-04-2026

M Siva

Department of Animal Nutrition,
Veterinary College and Research
Institute, Tamilnadu Veterinary
and Animal Sciences University
(TANUVAS), Namakkal, Tamil
Nadu, India

S Senthilkumar

Professor, Department of Animal
Nutrition, Veterinary College
and Research Institute
Tamil Nadu Veterinary and
Animal Sciences University
(TANUVAS), Namakkal, Tamil
Nadu, India

K Rajamanickam

Department of Veterinary,
Physiology and Biochemistry
Veterinary College and Research
Institute, Tamil Nadu
Veterinary and Animal Sciences
University (TANUVAS) Salem,
Tamil Nadu, India

G Ramesh Kumar

Department of Veterinary
Physiology and Biochemistry,
Veterinary College and Research
Institute, Tamil Nadu
Veterinary and Animal Sciences
University (TANUVAS) Theni,
Tamil Nadu, India

B Deepika

Department of Veterinary
Physiology and Biochemistry,
Veterinary College and Research
Institute, Tamil Nadu
Veterinary and Animal Sciences
University (TANUVAS),
Orathanadu, Tamil Nadu, India

Corresponding Author:

M Siva

Department of Animal
Nutrition, Veterinary College
and Research Institute,
Tamilnadu Veterinary and
Animal Sciences University
(TANUVAS), Namakkal,
Tamil Nadu, India

Rumen microflora and their nutritional requirements: Implications for optimizing fermentation efficiency in ruminants

M Siva, S Senthilkumar, K Rajamanickam, G Ramesh Kumar and B Deepika

DOI: <https://www.doi.org/10.33545/26174693.2026.v10.i6e.8654>

Abstract

The rumen, a vast and complex fermentation chamber, hosts a remarkably diverse and dense population of microorganisms, including bacteria, protozoa, fungi, and archaea. This symbiotic microflora is essential for the ruminant animal, as it breaks down complex fibrous plant materials, which the host cannot digest, into volatile fatty acids (VFAs) - the animal's primary energy source - and microbial protein. Understanding the distinct nutritional requirements and metabolic roles of these microbial groups is paramount for optimizing rumen fermentation efficiency. This review explores the composition of the rumen ecosystem, the specific nutrient needs (energy, nitrogen, minerals, vitamins) of its key microflora, and various dietary strategies and additives employed to manipulate this microbial community. Optimizing fermentation efficiency translates into improved animal performance (growth, milk yield), enhanced feed conversion, and a reduction in environmental impact, notably the mitigation of enteric methane emissions. Continued research in this area is crucial for sustainable and productive ruminant agriculture.

Keywords: Ruminants, rumen, microflora, microbial digestion, volatile fatty acids

Introduction

The rumen can be perceived as an anaerobic and methanogenic fermentation chamber that contains microorganisms, hosting a diverse microbial community including bacteria, anaerobic fungi, archaea, protozoa, and viruses, that could utilize, and increase the productivity of, cellulolytic feeds (i.e. straw, hay, silage and grass). These microorganisms play a vital role in breaking down nutrients, producing energy-rich volatile fatty acids (VFAs) essential for the host. There are considerable benefits associated with understanding rumen function, as rumen dynamics are almost solely responsible for providing nutrients to the host animal (C. Matthews *et al.*, 2019) [10].

At birth, the digestive system is germ-free, requiring contact with the mother to acquire rumen bacteria. In 24 hours, facultative anaerobic bacteria begin colonizing the rumen wall, with bacterial diversity reaching its peak among rumen microorganisms (Wei *et al.*, 2022) [9]. These microbes collaboratively ferment cellulose, hemicellulose, starch, and proteins, generating VFAs, microbial proteins, and gases such as carbon dioxide and methane. The animal absorbs VFAs through the rumen wall to meet its energy needs, while microbial proteins are digested in the small intestine, contributing significantly to the animal's amino acid intake and overall energy supply.

Optimizing rumen fermentation efficiency is crucial in ruminant nutrition. Inefficient fermentation can result in energy wastage, decreased microbial protein synthesis, digestive issues like acidosis, and increased environmental pollution. The goal is to maximize feed energy conversion into VFAs and microbial biomass while reducing methane emissions and ammonia excretion. Achieving this balance requires a comprehensive understanding of the rumen's microbial ecosystem, including the specific functions and nutritional needs of different microbial groups. Strategies such as dietary adjustments, feed processing techniques, and targeted additives are employed to modify the rumen environment, promoting more efficient fermentation pathways. This review explores the key microbial populations within the rumen, their nutritional requirements, and the implications for

improving fermentation efficiency in ruminant production systems, ultimately aiming to enhance productivity and reduce environmental impact.

Rumen Microflora Composition and Diversity

The rumen ecosystem is incredibly dense, with bacterial

concentrations estimated at 10^{10} to 10^{11} cells per milliliter of rumen fluid, and protozoa around 10^5 to 10^6 cells/ml. Anaerobic fungi are also a vital part of the community, and archaea, though fewer in number, perform a critical, albeit sometimes detrimental, function.




Fig 1: Ruminant stomach



Fig 2: Compartments of Ruminant stomach




Fig 3: Fermentation, energy production and utilization by rumen microbes

2.1 Rumen Bacteria

Rumen bacteria are the most numerous and functionally diverse group in the rumen, accounting for roughly half of the total microbial biomass. They are responsible for most fermentation activities. Rumen bacteria can be broadly classified into different functional groups based on the substrates they prefer or utilize:

- **Fibrolytic bacteria:** These specialize in breaking down complex structural carbohydrates like cellulose and hemicellulose. Major genera include *Fibrobacter* (e.g., *F. succinogenes*), *Ruminococcus* (e.g., *R. albus*, *R. flavefaciens*), and *Butyrivibrio* (e.g., *B. fibrisolvens*). Their enzymes (cellulases, xylanases) are crucial for initial fiber degradation.
- **Amylolytic bacteria:** These ferment starch and simpler sugars. Key genera include *Streptococcus* (e.g., *S. bovis*, which can multiply rapidly on high-starch diets, potentially leading to acidosis), *Succinitrio*, *Prevotella*, and *Ruminobacter*.

- **Proteolytic bacteria:** These degrade proteins, producing peptides and amino acids, which can then be further deaminated to produce ammonia. Many genera, including *Prevotella*, *Butyrivibrio*, *Ruminococcus*, and *Streptococcus*, exhibit proteolytic activity.
- **Lactic acid-utilizing bacteria:** Important for preventing acidosis by converting lactic acid (produced by rapid starch fermentation) into other VFAs. Examples include *Megasphaera elsdenii* and *Selenomonas ruminantium*.
- **Methanogenic archaea:** Though not technically bacteria, these single-celled organisms are crucial in removing hydrogen, an essential end-product that would otherwise inhibit overall fermentation. They use hydrogen to reduce carbon dioxide, producing methane (CH_4) as a byproduct. Common genera are *Methanobrevibacter* and *Methanosphaera*.

This diversity ensures that various components of the ruminant diet can be effectively degraded, albeit at different rates and by specialized microbial guilds.




Fig 4: Rumen Bacteria

Table 1: Classification of major rumen functional bacteria.

Classification	Latin Name of Bacteria	Gram Staining	Function	Source
Cellulolytic bacteria	<i>Ruminococcus flavefaciens</i>	G+	degrade cellulose, hemicellulase, xylan	Yeoman <i>et al.</i> [18]
	<i>Ruminococcus albus</i>	G+	degrade cellulose, hemicellulase, xylan	Rodriguez Hernandez <i>et al.</i> [28]
	<i>Fibrobacter succinogenes</i>	G–	ferment cellulose and cellobiose	
Amylolytic bacteria	<i>Butyrivibrio fibrisolvens</i>	G–	utilize cellulose, starch, and other polysaccharides, secrete pectinase, utilize xylan	
	<i>Streptococcus bovis</i>	G+	degrade starch to produce lactic acid	Cerqueira <i>et al.</i> [19]
	<i>Prevotella ruminicola</i>	G–	degrade starch, xylan, pectin	Anderson <i>et al.</i> [29]
Protein-degrading bacteria	<i>Ruminobacter amylophilus</i>	G–	ferment starch, degrade protein	
	<i>Ruminobacter amylophilus</i>	G–	ferment starch, degrade protein	Anderson <i>et al.</i>
	<i>Butyrivibrio fibrisolvens</i>	G–	utilize cellulose, starch and other polysaccharides, secrete pectinase, utilize xylan	Cotta <i>et al.</i>
Fat-degrading bacteria	<i>Anaerovibrio lipolytica</i>	G–	utilize fat and lactic acid	Prins <i>et al.</i>
Lactic acid-producing bacteria	<i>Bifidobacterium lactis</i>	G+	produce acetic acid, lactic acid, inhibit spoilage bacteria	Uusitupa <i>et al.</i>
	<i>Lactobacillus acidophilus</i>	G+	produce lactic acid and acetic acid	Anjum <i>et al.</i>
	<i>Streptococcus bovis</i>	G+	produce lactic acid and acetic acid	Cerqueira <i>et al.</i>
Lactic-acid-utilising bacteria	<i>Selenomonas ruminantium</i>	G–	utilize lactic acid to produce acetic and propionic acids	Fan <i>et al.</i>
	<i>Megasphaera elsdenii</i>	G–	ferment fructose, lactic acid	Monteiro <i>et al.</i>
Methanogens	<i>Methanobrevibacter ruminantium</i>	G+	reduce CO ₂ , CH ₄	Ma <i>et al.</i>
	<i>Methanomicrobium mobile</i>	G–	reduce CO ₂ , CH ₄	Yanagita <i>et al.</i>

2.2 Rumen Protozoa

Rumen protozoa, predominantly large ciliates, constitute a significant portion of the microbial biomass (up to 50% in some cases) due to their individual size. Their numbers can fluctuate significantly depending on the diet, typically being higher on high-forage diets (Morgavi *et al.*, 2010).

Protozoa perform several important roles:

- **Fermentation stabilization:** They ingest starch and bacteria, effectively slowing down the rate of starch fermentation and reducing the population of lactic acid producing bacteria. This helps buffer the rumen pH and prevent rapid pH drops associated with acidosis.
- **Fiber digestion:** Some protozoa exhibit cellulolytic and hemicellulolytic activity, contributing directly to fiber degradation, though potentially less efficiently than bacteria.

- **Bacterial predation:** By engulfing bacteria, protozoa recycle microbial nitrogen within the rumen, which can reduce the efficiency of microbial protein synthesis reaching the small intestine. However, protozoa themselves are large, high-quality sources of protein for the host, but their slower turnover rate in the rumen means they contribute less than bacteria to the microbial protein flow.
- **Hydrogen metabolism:** Certain protozoa harbor methanogenic archaea as endo- and ectosymbionts, contributing indirectly to methane production.

Defaunation (removal of protozoa) can sometimes increase microbial protein synthesis but may destabilize rumen pH and potentially slightly reduce fiber digestion efficiency, illustrating the complex, nuanced role of protozoa in the ecosystem.



Fig 5: Rumen Protozoa

2.3 Rumen Fungi

Anaerobic fungi are another key component of the rumen microflora, particularly prominent in animals consuming high-fiber diets. Although their biomass might be relatively small compared to bacteria and protozoa, they have a unique and vital function: (Joblin & Naylor, 1994) [5]. Rumen fungi (10^3 - 10^6 zoospores/ml) are anaerobic, falling into the class Neocallimastigomycetes, consisting of 6 previously recognised genera (Anaeromyces, Caecomyces, Cyllamyces, Neocallimastix, Orpinomyces and Piromyces) with 21 known species and, using molecular techniques, 2 recently discovered genera Oontomyces and Buwchfawromyces. Their discovery required a reevaluation of the central dogma within microbiology that all fungi were aerobic (C. Matthews *et al.*, 2019) [10].

- **Physical disruption:** Fungi possess extensive filamentous hyphae that penetrate deep into tough plant material (like stems and leaf sheaths), physically breaking open the complex plant cell wall matrix.
- **Potent fibrolytic enzymes:** They produce a wide array of highly active enzymes (cellulases, xylanases, pectinases) that degrade structural carbohydrates.
- **Facilitating bacterial access:** By physically disrupting the plant tissue and initiating degradation, fungi effectively "soften up" the material, creating more surface area and points of entry for the much smaller

fibrolytic bacteria to colonize and finish the job. Fungi are thus essential for the efficient utilization of low-quality, highly lignified forages, and their activity is a prerequisite for subsequent bacterial degradation of complex fibers.




Fig 6: Rumen Anaerobic fungi

2.4 Methanogenic Archaea

Archaea in the rumen are exclusively methanogens. As mentioned, they utilize hydrogen gas (H_2), along with carbon dioxide (CO_2) or other simple carbon compounds (like formate), to produce methane (CH_4) via an anaerobic respiration pathway. This is a critical function because hydrogen accumulation in the rumen inhibits the re-oxidation of cofactors like NADH, essential for ongoing carbohydrate fermentation. Methanogens thus act as a "hydrogen sink," maintaining low hydrogen levels and ensuring efficient fermentation (Cieslak *et al.*, 2013) [8]. However, methane is a significant loss of energy for the ruminant (2-12% of gross energy intake) and a potent greenhouse gas. Consequently, methanogenesis represents both an essential microbial function and a significant inefficiency, making it a major target for mitigation strategies.

Table 2: Concentration of rumen microbes and their function

Microorganism	Number	Function in the Rumen
Bacteria	10^{10} cells/mL	Ferment/degrade a range of substrates and reproduce
Protozoa	10^5 - 10^6 cells/mL	Ferment/degrade substrates, engulf starch, bacteria, and particles, and reproduce
Archaea	10^7 - 10^9 cells/mL	Metabolize hydrogen
Fungi	10^3 - 10^5 zoospores/mL	Source of cellulolytic enzymes; Fiber degradation and digestion
Bacteriophages	10^8 - 10^9 cells/mL	Infect bacteria

3. Nutritional Requirements of Rumen Microflora

The dynamic populations of rumen microflora have specific, sometimes overlapping, but also distinct nutritional requirements for energy, nitrogen, and other essential nutrients to maintain growth, metabolism, and fermentation activity. Providing balanced and appropriate levels of these nutrients is fundamental for supporting a healthy and efficient microbial community.

3.1 Energy Requirements (Carbohydrates)

Carbohydrates are the primary energy source for almost all rumen microorganisms. However, the form and fermentation rate of these carbohydrates drastically

influence microbial growth and fermentation patterns: (Van Soest, 1994) [7].

- **Structural Carbohydrates (Fiber):** Cellulolytic bacteria and fungi are highly adapted to fermenting cellulose and hemicellulose. This process is generally slow due to the complex structure and association with lignin, making overall fiber digestion slower and yielding relatively less energy per unit of time compared to starch. Adequate amounts of fermentable fiber are necessary to maintain optimal rumen motility, rumination, and buffering, thus supporting a stable, less acidic environment suitable for these fiber-digesting microbes.

- **Non-structural Carbohydrates (Starch, Sugars):** Amylolytic bacteria and some protozoa rapidly ferment starch and soluble sugars. This rapid fermentation provides abundant energy, supporting faster microbial growth and producing high levels of propionate (a VFA and key glucogenic precursor for the host). However, excessive and rapid starch fermentation can lead to a significant drop in rumen pH (acidosis) due to the accumulation of lactic acid, which inhibits fibrolytic bacteria and protozoa, negatively impacting overall rumen health and digestion efficiency.

The balance and rate of carbohydrate availability are, therefore, crucial. Maximizing energy supply for production without compromising rumen stability requires careful management of the starch-to-fiber ratio and the inclusion of buffering capacity in the diet.

3.2 Protein Requirements

Rumen microorganisms synthesize their own protein (microbial protein) from dietary nitrogen (N) sources. This microbial protein is subsequently digested by the host animal, supplying a large proportion of its metabolizable protein needs, often of high amino acid quality (Baldwin & Allison, 1983)^[1]. Microflora primarily utilize:

- **Ammonia (NH₃):** Many bacteria (especially fibrolytic types) prefer or even require ammonia as their primary N source for protein synthesis. Dietary protein and non-protein nitrogen (NPN) sources like urea are degraded to produce ammonia in the rumen. Optimal ammonia levels (5-10 mg/100 mL of rumen fluid) are necessary to maximize microbial growth. Excessive ammonia is absorbed and excreted, representing an efficiency loss and potential environmental concern.
- **Amino Acids and Peptides:** While some microbes prefer ammonia, others (often amylolytic and proteolytic bacteria, and many protozoa) also readily utilize amino acids and peptides from dietary or microbial sources. Providing a mix of N forms can support more rapid and efficient microbial growth, particularly when energy is abundant.

Efficient microbial protein synthesis relies critically on the synchronization of fermentable energy (carbohydrates) and degradable nitrogen availability. If energy is high but N is limiting, N will reduce microbial growth. If N is high but energy is limiting, excessive ammonia will be produced and wasted. Maintaining optimal and synchronous levels of both is essential for maximizing microbial protein yield and reducing N wastage. Rumen undegradable protein (bypass protein) escapes rumen fermentation and passes directly to the lower digestive tract, useful for high-producing animals whose requirements exceed microbial protein supply, but its provision must not limit rumen microflora N needs.

3.3 Minerals

Mackie & White (1990)^[3] stated that minerals are critical cofactors for numerous microbial enzymes and essential for cell structure and energy metabolism within the rumen ecosystem:

- **Phosphorus (P):** Essential for microbial growth, cell wall structure, synthesis of nucleic acids (DNA/RNA), and energy compounds like ATP. P is typically supplemented, especially on high-fiber diets.
- **Sulfur (S):** Required for the synthesis of sulfur-containing amino acids (methionine, cysteine) and

certain B-vitamins, particularly important when non-protein nitrogen (like urea) is used as a significant portion of the N supply.

- **Magnesium (Mg), Calcium (Ca), Potassium (K):** Play vital roles in enzyme function, cell membrane stability, and osmoregulation. While requirements are generally met by common diets, adequate levels are crucial for optimal microbial activity.
- **Trace Minerals (Cobalt, Copper, Zinc, Selenium, Iron):** Essential cofactors for various microbial enzymes, including those involved in fiber digestion, hydrogen metabolism, and synthesis of essential compounds (e.g., Cobalt for vitamin B12 synthesis). Deficiencies, though less common under typical management, can significantly impair microbial function. Cobalt is particularly noteworthy due to its specific role in B12 synthesis by rumen bacteria.

Appropriate mineral supplementation, informed by diet composition and animal needs, is vital to prevent imbalances that could compromise microbial metabolism and overall fermentation efficiency.

3.4 Vitamins

Rumen microorganisms are generally capable of synthesizing all water-soluble B-vitamins and vitamin K required by the host animal, eliminating the need for dietary supplementation under normal conditions. Some synthesis is extensive (e.g., B12, riboflavin, thiamine), while others are produced in smaller quantities. However, factors like high levels of dietary concentrate, disease stress, or extreme nutrient imbalances can potentially reduce microbial vitamin synthesis, suggesting that supplementation might be beneficial in specific situations for very high-producing animals, though definitive proof is often limited.

4. Implications for Optimizing Fermentation Efficiency in Ruminants

Manipulating the rumen microflora by understanding and meeting their distinct nutritional requirements offers significant opportunities to improve fermentation efficiency and overall ruminant productivity and sustainability.

4.1 Dietary Management and Optimization

- **Balancing Forage and Concentrate:** Maintaining an optimal forage-to-concentrate ratio is perhaps the most fundamental dietary management strategy. It involves providing sufficient concentrate (and starch) for high energy intake and productivity, while ensuring adequate effective fiber for proper rumination, buffering, and support of a robust fibrolytic population. This prevents acidosis, stabilizes pH, and ensures efficient utilization of both structural and non-structural carbohydrates.
- **Enhancing Fiber Digestibility:** Improving the quality and digestibility of forages through physical processing (chopping, grinding), steam-flaking grains (increasing starch fermentability), or using alkali treatments on low-quality residues can enhance microbial access and degradation, increasing overall energy yield.
- **Improving Nitrogen-Energy Synchronization:** Optimizing the timing and amounts of rumen-degradable nitrogen (from protein and NPN) and fermentable carbohydrates in the diet is critical for maximizing microbial protein synthesis and reducing energy-costly urea recycling and excretion.

4.2 Feed Additives for Microbial Modulation

Nagaraja & Titgemeyer, (2007) ^[4] observed that various feed additives were employed to selectively manipulate rumen microflora and fermentation pathways towards desired outcomes:

- **Ionophores (e.g., monensin):** These antibiotics selectively inhibit certain gram-positive rumen bacteria, leading to several beneficial shifts: increased propionate production (improving energy efficiency), decreased methane output (energy capture), and reduced breakdown of dietary protein (increasing bypass protein and reducing ammonia). Their use significantly improves feed conversion and animal performance.
- **Buffers (e.g., sodium bicarbonate):** Directly neutralize acids produced during rapid fermentation, helping to maintain an optimal rumen pH and prevent subacute rumen acidosis (SARA). This promotes stability, supporting fibrolytic bacteria and protozoa that are sensitive to low pH.
- **Direct-Fed Microbials (Probiotics):** Additive microbes, commonly live yeasts (e.g., *Saccharomyces cerevisiae*) or beneficial bacteria (e.g., *Lactobacillus* species). Yeasts can improve fiber digestion by scavenging oxygen (creating a more anaerobic environment), stabilizing pH, and producing growth factors for cellulolytic bacteria. Bacterial probiotics may potentially improve health or competitively exclude pathogens, but their consistent efficacy in ruminants is less clear compared to monogastric, likely due to the robustness of the existing rumen community.
- **Prebiotics:** Non-digestible carbohydrates (like oligosaccharides) that can selectively stimulate the growth and activity of beneficial bacteria already present in the rumen, potentially enhancing fermentation stability and efficiency.
- **Exogenous Enzymes (e.g., cellulases, xylanases):** Can potentially complement and enhance the fibrolytic activity of rumen microflora, especially for lower-quality forages, improving fiber digestibility and energy yield, though their effectiveness is often variable and depends on many factors.
- **Plant Secondary Compounds (e.g., Tannins, Saponins, Essential Oils):** Natural compounds with antimicrobial properties. Some (like tannins) can reduce protein degradation and potentially decrease methane, though they can also negatively impact palatability and overall digestibility at higher concentrations. Saponins and certain essential oils are actively researched for their potential to inhibit protozoa and methanogens.
- **Methane Inhibitors:** Specific chemical inhibitors are under intensive research and development to directly target methanogens and significantly reduce enteric methane emissions, thereby recapturing potentially wasted energy for the animal and minimizing environmental impact. Successful application of additives and dietary strategies requires careful consideration of the specific production system, diet composition, and desired outcomes, recognizing that the rumen ecosystem is complex and can exhibit varying responses.

4.3 Conclusion

Rumen microbiology is a dynamic field that has made substantial strides in understanding the vast diversity and complex metabolic networks within this unique ecosystem. Continued research leveraging advanced metagenomic,

transcriptomic, and proteomic techniques is further elucidating the intricacies of microbial interactions, nutritional cross-feeding, and specific requirements under varied conditions.

Optimizing rumen fermentation efficiency remains a core goal for enhancing sustainable ruminant agriculture. Future opportunities lie in:

- **Detailed Microbiome Engineering:** Developing highly targeted interventions, potentially including engineered strains or specific inhibitors, to precisely manipulate microbial populations for improved nutrient utilization and reduced environmental impact.
- **Precision Feeding Strategies:** Implementing sophisticated, synchronized dietary formulations and additive regimens tailored to individual animal genetics, life stage, and environmental factors to maximize efficiency and performance.
- **Harnessing Natural Biodiversity:** Further exploring the diverse natural rumen microflora, including poorly understood groups like the vast array of protozoa and fungi, to discover novel enzymes, metabolic pathways, and approaches for enhancing fiber degradation and minimizing energy loss.

A comprehensive understanding of the rumen microflora and their distinct nutritional requirements is foundational for implementing effective strategies that optimize fermentation efficiency, translating into economically viable, productive, and environmentally responsible ruminant production systems.

References

1. Baldwin RL, Allison MJ. Rumen metabolism. *Annu Rev Microbiol.* 1983;37(1):277-302.
2. Hungate RE. *The Rumen and Its Microbes*. New York: Academic Press; 1966.
3. Mackie RI, White BA. Recent advances in rumen microbial ecology and metabolism: Potential impact on nutrient utilization. *J Dairy Sci.* 1990;73(10):2971-2995.
4. Nagaraja TG, Titgemeyer EC. Ruminal acidosis in beef cattle: The science and its application. *J Anim Sci.* 2007;85:E1-E8.
5. Joblin KN, Naylor GE. Growth of rumen fungi on cell wall components of diverse plant materials. *Lett Appl Microbiol.* 1994;19(3):196-198.
6. Morgavi DP, Forano E, Martin C, Newbold CJ. Microbial ecosystems in the guts of agricultural animals. *Gut Microbes.* 2010;1(3):154-171.
7. Van Soest PJ. *Nutritional Ecology of the Ruminant*. Ithaca (NY): Cornell University Press; 1994.
8. Cieslak A, Szumacher-Strabel M, Stochmal A, Oleszek W. Plant secondary metabolites for rumen fermentation control and methane mitigation: A review. *Anim Feed Sci Technol.* 2013;186(1-2):1-28.
9. Wei W, Zhen Y, Wang Y, Shahzad K, Wang M. Advances of rumen functional bacteria and the application of micro-encapsulation fermentation technology in ruminants: A review. *Fermentation.* 2022;8:564.
10. Matthews C, Crispie F, Lewis E, Reid M, O'Toole PW, Cotter PD. The rumen microbiome: A crucial consideration when optimising milk and meat production and nitrogen utilisation efficiency. *Gut Microbes.* 2019;10(2):115-132.