

International Journal of Statistics and Applied Mathematics

ISSN: 2456-1452
Maths 2024; 9(3): 239-241
© 2024 Stats & Maths
<https://www.mathsjournal.com>
Received: 07-05-2024
Accepted: 12-06-2024

Kishore Mahajan
Department of Applied
Statistics, Indore College of
Applied Mathematics, Indore,
Madhya Pradesh, India

Decomposing continued fraction expansions of algebraic numbers with applications to Fiji cyclone return periods

Kishore Mahajan

DOI: <https://www.doi.org/10.22271/math.2024.v9.i3d.2391>

Abstract

Continued fraction expansions of algebraic numbers encode structural information through their partial quotient sequences, yet applications beyond number theory remain rare. This research decomposes continued fractions of algebraic numbers of degree 2 through 6 and maps the resulting partial quotient patterns onto tropical cyclone return period estimation for Fiji. Cyclone records from the Fiji Meteorological Service (1969-2022, 187 named storms) were encoded as sequences whose distributional properties were matched to continued fraction templates. The best-fitting algebraic number (degree 4, discriminant 1297) produced partial quotient statistics that minimized Kolmogorov-Smirnov distance ($D = 0.073$) against the empirical return period distribution. Estimated return periods for Category 3+ cyclones were 4.7 years (continued fraction model) versus 5.1 years (Gumbel extreme value model), with the continued fraction approach yielding narrower confidence intervals (width 1.83 vs. 2.47 years). The partial quotient sequence's periodic structure captured the quasi-periodic ENSO modulation of Fiji cyclone activity better than stationary extreme value models.

Keywords: Continued fractions, algebraic numbers, partial quotients, cyclone return periods, Fiji meteorology, extreme value estimation, ENSO modulation, periodic structure, distributional matching, tropical cyclones

Introduction

A policy statement from the Fiji National Disaster Management Office in 2019 called for “improved return period estimates for major cyclones to guide infrastructure design standards.” The standard approach uses extreme value theory fitting a Gumbel or generalized extreme value (GEV) distribution to annual maximum wind speeds ^[1]. But these models assume stationarity, which conflicts with the known quasi-periodic modulation of South Pacific cyclone activity by ENSO ^[2]. Continued fractions offer an alternative encoding of periodic structure. Every quadratic irrational has an eventually periodic continued fraction expansion (by Lagrange's theorem), and algebraic numbers of higher degree produce partial quotient sequences with complex but structured patterns ^[3]. If a cyclone return period sequence shares distributional properties with the partial quotients of a particular algebraic number, the algebraic number's periodic structure can be exploited for prediction ^[4]. This research tests this idea on 54 years of Fiji cyclone data. The goals were to identify which algebraic number's continued fraction best matches the empirical return period distribution, to compare the resulting estimates against standard extreme value models, and to assess whether the continued fraction approach captures ENSO-related periodicity ^[5].

Material and methods

Material

Named tropical cyclone records (Category 1-5 on the Saffir-Simpson scale) passing within 400 km of Fiji's capital Suva were obtained from the Fiji Meteorological Service for the 1969-2022 cyclone seasons, totaling 187 storms. Annual maximum sustained wind speeds and return periods (computed as the inverse of the Weibull plotting position) were derived. ENSO

Corresponding Author:
Kishore Mahajan
Department of Applied
Statistics, Indore College of
Applied Mathematics, Indore,
Madhya Pradesh, India

State (El Niño, La Niña, neutral) for each season was obtained from NOAA's Oceanic Niño Index. The research was conducted between October 2021 and April 2023 at the Indore College of Applied Mathematics, India. Ethical clearance was obtained (ICAM-2021-037, approved 24 September 2021).

Methods

Continued fraction expansions were computed for algebraic numbers α of degrees 2 through 6, with minimal polynomials having integer coefficients bounded by 20 and discriminants up to 5,000. For each α , the first 500 partial quotients were extracted. The empirical return period sequence (sorted

annual maxima) was compared against each partial quotient distribution via the Kolmogorov-Smirnov (KS) statistic. The algebraic number minimizing KS distance was selected. Return period estimates were derived from the selected continued fraction's periodic structure using a renewal-theoretic argument. Confidence intervals used a block bootstrap (block length = ENSO cycle ≈ 4 years, $B = 2,000$ resamples). Comparison models included Gumbel, GEV, and generalized Pareto distributions fit by maximum likelihood. All computations used Sage Math 9.7 and R 4.2.1.

Results

Table 1: Best-fitting algebraic number by degree and KS distance

Degree	Min. polynomial	Discriminant	KS (D)	Period length
2	x^2-3x+1	5	0.127	1
3	x^3-2x-1	59	0.098	--
4	x^4-x^3-3x+1	1297	0.073	4
5	x^5-2x^3+x-1	2713	0.084	--
6	$x^6-x^4-2x^2+1$	4621	0.091	--

The degree-4 algebraic number with discriminant 1297 achieved the smallest KS distance (0.073), with a quasi-period of 4 in its partial quotient sequence closely matching the 3-5

year ENSO cycle that modulates Fiji cyclone activity. Quadratic irrationals (degree 2) had truly periodic expansions but were too regular to capture the data's variability.




Fig 1: Scatter plot of continued fraction depth versus estimated cyclone return period.

Table 2: Return period estimates (years) for Category 3+ cyclones by model

Model	Estimate (yr)	95% CI width (yr)	AIC
Continued fraction	4.7	1.83	--
Gumbel	5.1	2.47	287.4
GEV	4.9	2.31	285.1
GP (threshold)	4.8	2.14	283.7

The continued fraction model produced the narrowest confidence interval (1.83 years), 26% narrower than Gumbel. Point estimates were similar across models (4.7-5.1 years).

The continued fraction's narrower interval reflects its exploitation of the periodic structure that stationary models treat as noise.




Fig 2: Violin plots of cyclone return periods across four intensity categories

Discussion

The degree-4 algebraic number's quasi-period of 4 aligning with the ENSO cycle is the key finding. ENSO modulates the position of the South Pacific Convergence Zone, shifting cyclone genesis regions and affecting Fiji's exposure [2]. Standard extreme value models, which assume IID annual maxima, absorb this periodic modulation into the variance, widening confidence intervals. The continued fraction approach explicitly models the periodicity, producing tighter intervals [4]. The KS distance of 0.073 indicates a reasonably good but not perfect distributional match. The mismatch arises partly because cyclone activity has additional non-ENSO drivers (Pacific Decadal Oscillation, local SST anomalies) that the algebraic number's periodic structure cannot fully capture [2]. Incorporating multiple algebraic numbers with different periods could improve the fit. A limitation is that the method requires a sufficiently long record to estimate the period reliably. The 54-year Fiji record spans roughly 11-14 ENSO cycles, which is adequate for a period-4 pattern but might be insufficient for longer-period modulations.

Conclusion

This research matched continued fraction expansions of algebraic numbers to Fiji cyclone return period distributions. A degree-4 algebraic number with quasi-period 4 achieved the best distributional fit (KS $D = 0.073$) and produced return period estimates with 26% narrower confidence intervals than the Gumbel model. For Fiji's disaster management authorities, the continued fraction approach offers tighter uncertainty bounds for infrastructure design standards. The method's exploitation of ENSO-like periodicity gives it a structural advantage over stationary extreme value models. Future work should test the approach on other Pacific island cyclone records and incorporate multi-period continued fraction mixtures.

Acknowledgements

Funding sources

This research was funded by a departmental allocation from the Indore College of Applied Mathematics.

Institutional support

The authors thank the Department of Applied Statistics for computing resources.

Contributions not qualifying for authorship

The authors acknowledge Dr. Apenisa Ratu for Fiji Meteorological Service data.

References

1. Coles S. An introduction to statistical modeling of extreme values. London: Springer; 2001.
2. Kuleshov Y, Qi L, Fawcett R, Jones D. On tropical cyclone activity in the Southern Hemisphere: trends and the ENSO connection. *Geophys Res Lett*. 2008;35(14).
3. Hardy GH, Wright EM. An introduction to the theory of numbers. 6th ed. Oxford: Oxford University Press; 2008.
4. Khintchine AY. Continued fractions. Mineola (NY): Dover Publications; 1997.
5. Fiji Meteorological Service. Climate of Fiji and tropical cyclone records. Suva: Fiji Meteorological Service; 2022.
6. Hosking JRM, Wallis JR. Regional frequency analysis: An approach based on L-moments. Cambridge: Cambridge University Press; 1997.
7. Olds CD. Continued fractions. Washington (DC): Mathematical Association of America; 1963.
8. Wall HS. Analytic theory of continued fractions. New York (NY): Chelsea Publishing; 1948.
9. Perron O. Die Lehre von den Kettenbrüchen. Leipzig: Teubner; 1954.
10. Rockett AM, Szűsz P. Continued fractions. Singapore: World Scientific; 1992.
11. Embrechts P, Klüppelberg C, Mikosch T. Modelling extremal events for insurance and finance. Berlin: Springer; 1997.
12. Beirlant J, Goegebeur Y, Segers J, Teugels JL. Statistics of extremes: Theory and applications. Chichester: John Wiley & Sons; 2004.
13. de Melo AB, Torres RR. Tropical cyclone return periods in the South Pacific. *Int J Climatol*. 2018;38(12):4562–4578.
14. Hendon HH, Wheeler MC, Zhang C. Seasonal dependence of the MJO-ENSO relationship. *J Clim*. 2007;20(3):531–543.