

International Journal of Statistics and Applied Mathematics

ISSN: 2456-1452
NAAS Rating (2025): 4.49
Maths 2025; 10(11): 56-59
© 2025 Stats & Maths
<https://www.mathsjournal.com>
Received: 23-09-2025
Accepted: 26-10-2025

Bruno Nascimento
Department of Probability and
Statistics, Curitiba Academy of
Statistical Research, Curitiba,
Paraná, Brazil

Camila Araujo
Department of Probability and
Statistics, Curitiba Academy of
Statistical Research, Curitiba,
Paraná, Brazil

Corresponding Author:
Bruno Nascimento
Department of Probability and
Statistics, Curitiba Academy of
Statistical Research, Curitiba,
Paraná, Brazil

Computing mood test extensions for multivariate scale parameters with applications to brahmaputra flood frequency analysis

Bruno Nascimento and Camila Araujo

DOI: <https://doi.org/10.22271/math.2025.v10.i11a.2487>

Abstract

An observation that has gone largely unnoticed in the hydrological literature is that the Mood test for equal scales, originally designed for univariate samples, can be extended to the multivariate setting in ways that capture joint scale shifts across correlated flood indicators. This research develops such an extension and applies it to annual maximum discharge, peak water level, and flood duration data from 16 gauging stations along the Brahmaputra River in India and Bangladesh, covering the years 2000 to 2022. Graph-theoretic representations of the station network defined the spatial dependence structure, while the Laplace equation on the network graph provided harmonic interpolation of flood characteristics between stations. Laplace transforms of the flood duration distribution supplied moment-generating functions used to derive the asymptotic variance of the multivariate Mood statistic. Survey design principles guided the selection of a balanced panel of stations, and support vector machine classifiers benchmarked the test's ability to distinguish pre- and post-2010 flood regimes. The multivariate Mood test detected a significant scale shift ($p = 0.003$) in the joint flood distribution after 2010, a shift missed by the univariate Mood test applied to any single indicator. The support vector machine achieved 84.6 percent classification accuracy using the multivariate scale features as inputs, compared with 71.3 percent using univariate features alone. These results indicate that multivariate extensions of classical scale tests can reveal regime changes in complex flood systems that univariate analyses overlook.

Keywords: Graph theory, laplace equation, laplace transform, survey design, support vector machines, mood test, multivariate scale, flood frequency, brahmaputra river, regime detection

Introduction

Most hydrologists who have worked on the Brahmaputra would agree on one thing: the river doesn't follow the rules. Peak discharges at Pandu, India, have exceeded 70,000 cubic metres per second in some years and barely reached 40,000 in others, a variability that makes flood frequency estimation both critical and frustrating ^[1]. Standard approaches treat each flood indicator, discharge, water level, duration, separately, testing for trends or regime changes one variable at a time. But floods are multivariate events, and a shift in the joint scale of these indicators can go undetected when each margin is tested alone.

The Mood test, proposed by Mood in 1954 ^[2], uses the sum of squared deviations of the ranks from the combined sample median rank as its test statistic. It is a rank-based test for differences in scale (dispersion) between two populations, requiring no distributional assumptions beyond continuity. Its power is highest against alternatives where the two populations differ in spread but share the same location, making it well suited to flood frequency applications where non-stationarity may manifest as increased variability rather than a shift in the mean ^[3].

Extending the Mood test to multiple response variables requires a definition of multivariate ranks and a corresponding test statistic. Several approaches exist, including componentwise ranks, spatial ranks based on data depth, and graph-based ranks defined by nearest-neighbour distances ^[4]. This research adopts the graph-based approach, defining ranks on the minimum spanning tree of the inter-observation distance matrix. The resulting multivariate Mood

statistic inherits the distribution-free property of its univariate parent under the null hypothesis of equal multivariate scales. The Brahmaputra application uses data from 16 stations spanning the river's course from Dibrugarh, Assam, to Bahadurabad, Bangladesh, over 23 flood seasons (2000-2022). Graph theory provides the spatial structure, the Laplace equation on the station network interpolates between stations, and Laplace transforms supply the moment-generating machinery for deriving the test's asymptotic properties. The research was conducted at the Curitiba Academy of Statistical Research, Curitiba, Parana, Brazil.

Model description and parameterisation

The multivariate Mood statistic was constructed as follows. For two samples of trivariate flood vectors (discharge, water level, duration) of sizes m and n , the combined sample of $N = m + n$ observations was used to build a minimum spanning tree (MST) on the Euclidean distances in three-dimensional space. Each observation was assigned a graph rank equal to its depth in the MST, defined as the number of edges on the longest path from that node to a leaf ^[4]. The multivariate Mood Statistic M was defined as $M = \sum_{i \text{ in sample 1}} (R_i - (N+1)/2)^2$, where R_i is the graph rank, analogous to the univariate Mood statistic but with graph ranks replacing linear ranks. Under the null hypothesis of equal multivariate scale, M has an asymptotic normal distribution with mean and variance derived from the MST's degree sequence. The Laplace transform of the flood duration distribution, $L(s) = E[\exp(-sD)]$, was used to compute the variance correction for the heavy-tailed duration component, ensuring that the asymptotic approximation remained accurate even when the marginal distributions were skewed ^[5].

Material and methods

Material

Annual flood records for 16 gauging stations along the Brahmaputra River were obtained from the India Central Water Commission (stations in Assam) and the Bangladesh Water Development Board (stations in Bangladesh) for the flood seasons of 2000 through 2022. Three variables were recorded for each station-year: annual maximum

instantaneous discharge (cubic metres per second), corresponding peak water level (metres above datum), and flood duration (days above danger level). Station selection followed a survey design that ensured balanced representation of the upper, middle, and lower river reaches, with at least four stations per reach ^[6]. Satellite-derived inundation extent from the Dartmouth Flood Observatory was used as an auxiliary variable for stations with incomplete records. The research was conducted at the Department of Probability and Statistics, Curitiba Academy of Statistical Research, Curitiba, Parana, Brazil, between January 2022 and June 2023. Ethics approval was not required because the research used publicly available hydrological records. Analysis used R version 4.2.2 with the energy, e1071, and igraph packages.

Methods

The 23-year record was split at 2010 to define pre- and post-2010 samples, motivated by reports of intensifying flood variability in the Brahmaputra after 2010 ^[7]. The univariate Mood test was applied separately to each of the three flood indicators, using the standard squared-rank statistic with a continuity-corrected normal approximation. The multivariate Mood test was computed on the trivariate flood vectors using the MST-based graph ranks described above. The Laplace equation on the station network graph, defined with inverse-distance edge weights, was solved to interpolate flood characteristics at ungauged points between stations, providing a spatially continuous flood surface for visualisation. The Laplace transform of the flood duration distribution was estimated by fitting a gamma distribution to the duration data by maximum likelihood and evaluating $L(s) = (\beta / (\beta + s))^\alpha$ analytically. Support vector machines with a radial basis function kernel were trained to classify each station-year as pre- or post-2010 using either the three univariate scale features (squared deviations from the median for each indicator) or the single multivariate scale feature (the graph-rank-based Mood contribution). Classification accuracy was assessed by leave-one-year-out cross-validation ^[8].

Results

Table 1: Univariate and multivariate Mood test results for pre- vs post-2010 flood regimes

Test	Statistic	p-value	Conclusion
Mood (Discharge)	2.31	0.128	Not signif.
Mood (Water Level)	1.87	0.171	Not signif.
Mood (Duration)	2.64	0.104	Not signif.
Multivariate Mood	8.92	0.003	Significant

None of the three univariate Mood tests detected a significant scale shift at the 0.05 level, with p-values ranging from 0.104 (duration) to 0.171 (water level). The multivariate Mood test,

by contrast, yielded a statistic of 8.92 ($p = 0.003$), providing strong evidence for a joint scale shift in the trivariate flood distribution after 2010 (Table 1).

Table 2: Support vector machine classification accuracy for pre- vs post-2010 regime

Feature Set	Accuracy (%)	Sensitivity (%)	Specificity (%)
Univariate scale features	71.3	68.4	74.1
Multivariate scale feature	84.6	82.1	87.0
Combined features	86.2	84.3	88.1

The SVM trained on the multivariate scale feature achieved 84.6 percent accuracy, a 13.3 percentage point gain over the univariate feature set (71.3 percent). Combining both feature sets gave a marginal further improvement to 86.2 percent

(Table 2), suggesting that most of the discriminative information was already captured by the multivariate Mood contribution.




Fig 1: Annual flood frequency index at Brahmaputra gauging stations, 2000-2022




Fig 2: Proportional contribution of flood components to total Brahmaputra discharge variability

Discussion

The multivariate Mood test's ability to detect a joint scale shift that all three univariate tests missed illustrates a well-known but often ignored principle: marginal tests can miss multivariate signals. The joint shift appeared because post-2010 floods were not just larger or longer on average but more variable in their joint behaviour, years with high discharge but short duration alternated with years of moderate discharge but exceptionally long duration, a pattern that inflates the multivariate scale without necessarily inflating any single marginal scale ^[9].

The graph-theoretic construction of multivariate ranks via the minimum spanning tree has the advantage of being affine-invariant: rotating or rescaling the flood variables doesn't change the MST structure or the resulting ranks. This property is important because the three flood indicators have different units and vastly different magnitudes ^[4]. The Laplace equation interpolation on the station network produced smooth flood surfaces that agreed visually with satellite inundation maps, though formal validation against the Dartmouth Flood Observatory data showed a mean absolute error of 8.3 percent in inundation extent.

The SVM results confirm that the multivariate scale shift has practical predictive value: an 84.6 percent accuracy in distinguishing pre- from post-2010 flood regimes means that the joint scale feature carries regime information that could inform early warning systems. The radial basis kernel's implicit mapping to a high-dimensional feature space may have captured nonlinear interactions between the flood indicators that the linear Mood statistic does not explicitly model ^[8].

A limitation is the relatively short record: 11 years before and 12 years after the putative 2010 breakpoint provide limited power for detecting subtle shifts. The choice of 2010 as the split point was informed by prior literature ^[7] rather than estimated from the data, which avoids selection bias but means the test is conditional on this choice being correct.

Sensitivity analysis

To assess the sensitivity of the multivariate Mood test to the choice of split point, the analysis was repeated with breakpoints at every year from 2005 to 2015. The test statistic exceeded the 0.05 critical value for split points at 2009, 2010, and 2011, with the maximum statistic at 2010 ($M = 8.92$).

Split points before 2008 or after 2012 yielded non-significant results, supporting the interpretation that the regime change was concentrated around 2010. Robustness to the distance metric was checked by replacing Euclidean distance with Mahalanobis distance in the MST construction; the resulting p-value shifted from 0.003 to 0.005, still clearly significant.

Conclusion

This research extended the Mood test for equal scales to the multivariate setting using minimum spanning tree graph ranks and applied it to trivariate flood data from the Brahmaputra River. The multivariate test detected a significant joint scale shift after 2010 that no univariate test identified, demonstrating the value of multivariate scale diagnostics in flood frequency analysis. Support vector machine classification confirmed the practical discriminative power of the multivariate scale feature. Graph-theoretic spatial interpolation via the Laplace equation provided smooth flood surfaces between stations. These results argue for routine use of multivariate scale tests alongside univariate analyses in flood regime assessment and suggest that the Brahmaputra's flood variability structure underwent a genuine change around 2010. Future research should extend the multivariate Mood framework to accommodate spatial correlation among stations and test it on other large river systems with suspected non-stationary flood behaviour.

Acknowledgements

Funding sources

Internal departmental resources supported this research. No external funding was received.

Institutional support

The authors thank the Curitiba Academy of Statistical Research, Curitiba, Brazil, for providing computing facilities.

Contributions not qualifying for authorship

The authors acknowledge Mr. Rafael Ferreira for data retrieval assistance and Dr. Mariana Silva for suggestions on graph-based ranking.

References

1. Immerzeel WW, van Beek LPH, Bierkens MFP. Climate change will affect the Asian water towers. *Science*. 2010;328(5984):1382–1385.
1. Mood AM. On the asymptotic efficiency of certain nonparametric two-sample tests. *Ann Math Stat*. 1954;25(3):514–522.
2. Kundzewicz ZW, Robson AJ. Change detection in hydrological records: a review of the methodology. *Hydrol Sci J*. 2004;49(1):7–19.
3. Friedman JH, Rafsky LC. Multivariate generalizations of the Wald-Wolfowitz and Smirnov two-sample tests. *Ann Stat*. 1979;7(4):697–717.
4. Billingsley P. Probability and measure. 3rd ed. New York: John Wiley and Sons; 2012. p. 1–624.
5. Lohr SL. Sampling: design and analysis. 2nd ed. Boca Raton: Chapman and Hall/CRC; 2010. p. 1–596.
6. Mirza MMQ. Climate change, flooding in South Asia and implications. *Reg Environ Change*. 2011;11(S1):95–107.
7. Cortes C, Vapnik V. Support-vector networks. *Mach Learn*. 1995;20(3):273–297.
8. Anderson TW. An introduction to multivariate statistical analysis. 3rd ed. New York: John Wiley and Sons; 2003. p. 1–721.

9. Prim RC. Shortest connection networks and some generalizations. *Bell Syst Tech J*. 1957;36(6):1389–1401.
10. Merz B, Thielen AH. Flood risk curves and uncertainty bounds. *Nat Hazards*. 2009;51(3):437–458.
11. Chebana F, Ouara TBMJ. Multivariate quantiles in hydrological frequency analysis. *Environmetrics*. 2011;22(1):63–78.
12. Grover PL, Burn DH, Cunderlik JM. A comparison of index flood estimation procedures for ungauged catchments. *Can J Civ Eng*. 2002;29(5):734–741.
13. West DB. Introduction to graph theory. 2nd ed. Upper Saddle River: Prentice Hall; 2001. p. 1–588.
14. Salvadori G, De Michele C. Multivariate multiparameter extreme value models and return periods. *Water Resour Res*. 2010;46(10): W10501.
15. Koutsoyiannis D. Statistics of extremes and estimation of extreme rainfall. *Hydrol Sci J*. 2004;49(4):575–590.
16. Scholz FW, Stephens MA. K-sample Anderson-Darling tests. *J Am Stat Assoc*. 1987;82(399):918–924.
17. Coles S. An introduction to statistical modeling of extreme values. London: Springer; 2001. p. 1–208.