



International Journal of Veterinary Sciences and Animal Husbandry



ISSN: 2456-2912

NAAS Rating (2026): 4.61

VET 2026; 11(6): 187-190

© 2026 VET

www.veterinarypaper.com

Received: 11-04-2026

Accepted: 15-05-2026

Dr. B Deepika

Assistant Professor, Department of Veterinary Physiology and Biochemistry, Veterinary College and Research Institute, Orathanadu, Tamil Nadu, India

Dr. G Rameshkumar

Assistant Professor, Department of Veterinary Physiology and Biochemistry, Veterinary College and Research Institute, Theni, Tamil Nadu, India

Dr. K Rajamanickam

Assistant Professor, Department of Veterinary Physiology and Biochemistry, Veterinary College and Research Institute, Salem, Tamil Nadu, India

Dr. M Siva

Assistant Professor, Department of animal Nutrition, Veterinary College and Research Institute, Namakkal, Tamil Nadu, India

Dr.S Senthilkumar

Professor, Department of animal Nutrition, Veterinary College and Research Institute, Namakkal, Tamil Nadu, India

Corresponding Author:

Dr. B Deepika

Assistant Professor, Department of Veterinary Physiology and Biochemistry, Veterinary College and Research Institute, Orathanadu, Tamil Nadu, India

Recent approaches in rumen fermentation manipulation for sustainable ruminant production

Deepika B, Rameshkumar Ganesan, Rajamanickam Kandasamy, Siva M and Senthilkumar S

DOI: <https://www.doi.org/10.22271/veterinary.2026.v11.i6c.3521>

Abstract

The microbial activity occurring within the rumen enables ruminants to convert fibrous plant materials into readily usable nutrients essential for their growth and productivity. The ruminal microbial ecosystem contains bacteria, protozoa, fungi, and methanogenic archaea that collectively degrade dietary carbohydrates and proteins into volatile fatty acids and microbial proteins. Despite its importance, ruminal fermentation is often associated with considerable energy losses through methane production and excessive nitrogen excretion, thereby reducing feed efficiency and contributing to environmental pollution. Recent advances in nutritional and microbial strategies have focused on optimizing rumen fermentation to improve productivity and sustainability in livestock production systems. Several strategies such as dietary modification, supplementation with probiotics, ionophores, organic acids, phytobiotics, essential oils, exogenous enzymes, and plant-derived secondary compounds have shown considerable potential in enhancing ruminal fermentation and lowering methane production. Emerging genetic and biotechnological techniques directed toward rumen microorganisms are gaining attention for their ability to improve feed utilization efficiency and promote environmentally sustainable livestock production. This review highlights the recent advancements in rumen fermentation modulation and emphasizes their significance in enhancing animal performance while reducing ecological impact.

Keywords: rumen fermentation, methane mitigation, probiotics, phytobiotics, feed additives, sustainable livestock production

1. Introduction

Ruminant animals possess a specialized digestive system capable of converting low-quality fibrous feed materials into valuable animal products such as milk, meat, and wool. The rumen acts as a large anaerobic fermentation chamber inhabited by a diverse population of microorganisms including bacteria, protozoa, fungi, and methanogens. These microbes ferment dietary carbohydrates and proteins into volatile fatty acids (VFAs), microbial proteins, gases, and other metabolites that support animal growth and production.

Among the fermentation products, acetate, propionate, and butyrate serve as major energy sources for ruminants. However, the fermentation process is not completely efficient because a portion of dietary energy is lost as methane. Methane production also contributes significantly to greenhouse gas emissions from livestock. In addition, inefficient protein utilization in the rumen leads to excessive ammonia production and nitrogen wastage.

Improvement of rumen fermentation has therefore become an important strategy in sustainable livestock production. Manipulation of rumen microbial activity can enhance feed utilization, improve growth and milk production, reduce methane emission, and minimize environmental pollution. Various nutritional, microbial, and biotechnological interventions have been investigated to achieve these objectives.

2. Rumen Microbial Ecosystem

The rumen contains a highly complex microbial ecosystem composed of bacteria, protozoa, fungi, archaea, and bacteriophages (Gebrehiwot, 2014) [9]. Bacteria are the predominant microorganisms and are mainly responsible for carbohydrate and protein degradation.

Fibrolytic bacteria digest cellulose and hemicellulose, whereas amylolytic bacteria ferment starch-rich feed materials.

Protozoa play an important role in starch utilization and maintenance of ruminal fermentation stability; however, their close association with methanogenic microorganisms contributes indirectly to methane synthesis (Patra, 2012). Anaerobic fungi facilitate the digestion of fibrous feed materials by invading plant tissues and disrupting complex lignocellulosic structures. Methanogenic archaea utilize the hydrogen released during microbial fermentation for methane formation.

The balance among these microbial populations determines fermentation efficiency and nutrient availability in the host animal (Gworgwor *et al.*, 2006) ^[10]. Modification of ruminal microbial populations through nutritional and microbial interventions can markedly affect fermentation patterns, nutrient utilization, and overall animal performance.

3. Significance of Optimizing Rumen Fermentation

Ruminant animals serve as major contributors to dietary protein supply, especially in regions where livestock predominantly depend on low-quality roughages and agro-industrial lignocellulosic by-products. However, these feed resources are generally poor in nutritional quality and often limit optimum animal productivity.

4. Methods of Rumen Manipulation

- Genetic Approaches
- Non-Genetic Approaches (Dietary manipulation)

4.1 Genetic Approaches

Genetic Approaches

According to Jansen (1975) ^[12], genetic modification of rumen microorganisms provides an opportunity to improve their inherent functions or introduce desirable metabolic characteristics. Genetically engineered microbes can be developed to degrade lignin and other complex fibrous components more efficiently, neutralize toxic substances, synthesize essential amino acids, reduce methane formation, and tolerate acidic ruminal conditions.

McSweeney *et al.* (1999) reported that modified rumen microorganisms may enhance fermentation efficiency by improving fiber digestion and reducing undesirable fermentation end products. Similarly, Chang (1996) ^[2] demonstrated that incorporation of specific genes into gastrointestinal microbes could enhance their metabolic activities and functional capabilities. Furthermore, studies conducted by Forsberg *et al.* (1993) ^[7] indicated that the introduction of superior microbial strains or novel microbial species into the rumen ecosystem may improve feed utilization efficiency and overall animal productivity.

4.2 Non-Genetic Approaches

4.2.1 Feed Additives

Dietary feed additives are commonly included in animal rations to improve nutrient digestibility, enhance productive performance, reduce metabolic complications, and minimize the environmental burden associated with livestock feeding systems (MacRae and Lobley, 1982) ^[14]. These supplements contribute to better digestive activity, improved feed efficiency, and reduced environmental contamination.

4.2.2 Probiotics

Stewart *et al.* (1988) stated that feed digestion in ruminants

primarily depends on microbial fermentation and enzymatic activities carried out by the diverse microbial population present within the rumen. In recent years, increasing emphasis has been placed on understanding the importance of ruminal and intestinal microbiota in maintaining animal health and enhancing productive performance. Disturbances in microbial balance caused by the proliferation of undesirable microorganisms can adversely affect digestion, health status, and productivity. To overcome these challenges, researchers have investigated the use of probiotics comprising beneficial bacteria, fungi, and yeast cultures as natural feed additives capable of regulating rumen fermentation and improving animal performance.

Furthermore, Fuller (1989) ^[8] reported that probiotic supplementation could improve growth performance, enhance disease resistance, increase production efficiency, and provide an economical feeding strategy. The use of microbial probiotics has also been associated with significant improvements in body weight gain and milk yield, with production increases of approximately 7–8% observed in supplemented animals.

4.2.3 Ionophores

Ionophores are widely used as nutritional modifiers for regulating ruminal fermentation, thereby enhancing feed utilization efficiency and overall animal productivity. These compounds are carboxylic polyether antibiotics produced by different species of *Streptomyces* and primarily act by altering the metabolism of Gram-positive bacteria within the rumen. Their action promotes greater propionate production while reducing acetate synthesis, leading to improved energy efficiency. Ionophores also decrease ruminal proteolysis and ammonia production, thereby improving nitrogen utilization. Commonly used ionophores include monensin, lasalocid, tetronasin, salinomycin, narasin, lysocellin, laidlomycin, nigericin, and valinomycin. By facilitating selective ion transport across microbial cell membranes, these compounds modify ruminal microbial populations and enhance fermentation efficiency.

4.2.4 Buffering Agents

Buffering agents are incorporated into dairy rations to counteract excessive acidity within the digestive system and maintain normal ruminal conditions. These compounds support the animal's natural buffering mechanisms, particularly salivary buffering, thereby preventing acid accumulation associated with high-concentrate feeding and disorders such as lactic acidosis. Commonly used buffering substances include sodium bicarbonate, calcium carbonate, magnesium oxide, sodium sesquicarbonate, and bentonite. Inclusion of these agents in the diet helps maintain optimum ruminal pH, supports microbial activity, sustains milk production, and promotes overall herd health.

4.2.5 Organic Acids

Organic acids such as aspartate, fumarate, and malate have been investigated for their ability to enhance lactate utilization by *Selenomonas ruminantium*, an important ruminal microorganism. Among these compounds, malate has been considered the most effective due to its role as an alternative electron acceptor, redirecting hydrogen toward succinate and subsequently propionate production during fermentation. Supplementation of DL-malate has been reported to stabilize ruminal pH, decrease methane production, and improve volatile fatty acid profiles, thereby reducing the risk of lactic

acidosis.

4.2.6 Essential Oils

Plant-derived essential oils are aromatic volatile substances possessing strong antimicrobial properties. These bioactive compounds alter microbial cell membrane integrity and interfere with vital cellular activities such as energy production, protein movement, and ion transport. Their inhibitory action is mainly directed against Gram-positive bacteria, similar to the effects produced by monensin, thereby influencing ruminal fermentation characteristics. Because of their natural origin and targeted antimicrobial action, essential oils are gaining importance as eco-friendly alternatives for rumen fermentation modulation.

4.2.7 Exogenous Fibrolytic

Supplementation of exogenous enzymes, particularly cellulases, hemicellulases, xylanases, and related fibrolytic enzymes, has shown considerable potential in enhancing fiber degradation within the rumen. These enzyme additives improve the digestibility of plant cell wall components, support better animal growth, and modify ruminal fermentation patterns in a manner that may lower methane production. Moreover, enzyme supplementation often decreases the acetate-to-propionate ratio, thereby improving fermentation efficiency.

4.2.8 Fat Supplementation

Inclusion of fat in ruminant diets is considered an effective nutritional strategy for increasing dietary energy density while simultaneously reducing methane emission. Medium-chain fatty acids (C8–C14), particularly those present in coconut oil and palm oil, have shown significant inhibitory effects on methanogenesis. Since fats undergo minimal fermentation within the rumen and do not contribute hydrogen required for methane synthesis, they play an important role in methane mitigation and improvement of feed efficiency.

4.2.9 Phytobiotics

Phytobiotics, which are plant-based feed additives, are increasingly incorporated into ruminant diets to enhance fiber digestion, promote the growth of beneficial microbial populations, and improve ruminal fermentation patterns. These natural compounds have been reported to increase propionate production, reduce methane emission, and enhance overall rumen efficiency. Owing to their natural origin and multiple beneficial effects, phytobiotics are gaining considerable importance in modern animal nutrition.

4.2.10 Plant Secondary Metabolites (PSM)

Plant secondary metabolites such as tannins, saponins, flavonoids, and essential oils are known to influence ruminal fermentation by suppressing methanogenic archaea and altering protozoal activity. Some of these compounds also enhance propionate production by redirecting hydrogen away from methane synthesis pathways. Due to their dual benefits in improving fermentation efficiency and reducing environmental pollution, these bioactive compounds are considered promising natural modifiers of rumen fermentation.

4.2.11 Biological Additives and Enzymes

Numerous direct-fed microbial products are currently available commercially and are widely utilized for improving animal health and productive performance. These

preparations commonly include beneficial microorganisms such as *Lactobacillus*, *Bifidobacterium*, and *Propionibacterium*, which can be administered in the form of probiotics, prebiotics, or synbiotics. Such microbial supplements support rumen development, enhance digestive efficiency, encourage competitive exclusion of pathogenic microorganisms, and reduce dependence on antibiotic growth promoters. Similarly, fibrolytic enzymes including endoglucanases, xylanases, and esterases, particularly those derived from ruminal microorganisms, are being increasingly explored for commercial application. These biological additives provide environmentally sustainable approaches for improving feed utilization and enhancing livestock productivity.

5. Defaunation

Defaunation, which refers to the removal of ruminal protozoa, can markedly influence rumen fermentation dynamics and methane production. Elimination of protozoal populations has been shown to increase bacterial growth, enhance microbial protein synthesis, and improve nitrogen flow to the duodenum, particularly in animals receiving protein-limited diets. Although defaunation may slightly reduce the digestion of structural carbohydrates present in plant cell walls, its beneficial effects including improved protein utilization, enhanced animal performance, and reduced methane emission are considered highly advantageous.

5.1 Methods of Defaunation

1. Neonatal isolation

Agarwal *et al.* (1991) ^[1] described neonatal isolation as an effective defaunation technique in which newborn ruminants are separated from their dams for two to three days after birth. This practice prevents the transmission of ciliate protozoa while permitting colonization by indigenous bacterial populations, thereby establishing a protozoa-free rumen environment.

2. Chemical Defaunation

Santra and Karim (2002) reported that ruminal protozoa can be selectively eliminated through the use of chemical agents such as copper sulphate, monoxol, and sodium lauryl sulphate. These compounds are capable of producing protozoa-free animals without causing major disturbances to the overall ruminal microbial balance.

3. Diet-Based Defaunation

Fonty *et al.* (1984) ^[6] demonstrated that defaunation can also be achieved through dietary manipulation of the rumen environment. Ruminal ciliate protozoa are highly sensitive to acidic conditions and are unable to survive when ruminal pH falls below 5.8, particularly below 5.0. Feeding animals with high-energy diets such as barley or maize following a 24-hour fasting period induces temporary ruminal acidosis, which may ultimately result in complete elimination of protozoal populations.

Conflict of Interest

Not available

Financial Support

Not available

References

1. Agarwal N, Kewalramani N, Kamra DN, Agrawal DK, Nath K. Hydrolytic enzymes of buffalo rumen:

- comparison of cell-free rumen fluid, bacterial and protozoal fractions. Buffalo Journal. 1991;2(2):203-207.
2. Chang H. Genetic engineering to enhance microbial interference and related therapeutic applications. Nature Biotechnology. 1996;14(4):423-431.
 3. Dominguez-Bello MG, Escobar A. Rumen manipulation for the improved utilization of tropical forages. Animal Feed Science and Technology. 1997;69(1):91-102.
 4. Donald EJ, Ward GM. Estimates of animal methane emissions. Environmental Monitoring and Assessment. 1996;42(1-2):133-141.
 5. Egan AR, Frederick F, Dixon RM. Improving efficiency of use supplements by manipulation of management procedures. In: Dixon RM, editor. Ruminant Feeding Systems Utilizing Fibrous Agricultural Residues. International Development Program of Australian Universities and Colleges (IDP); 1986. p. 69-81.
 6. Fonty G, Jouany JP, Senaud J, Gouet PH, Grain J. The evolution of microflora, macrofauna and digestion in the rumen of lambs from birth to four months. Canadian Journal of Animal Science. 1984;64(1):165-169.
 7. Forsberg CW, Cheng KJ, Krell PJ, Phillips JP. Establishment of rumen microbial gene pools and their manipulation to benefit digestion by domestic animals. In: Proceedings of the Seventh World Conference on Animal Production; 1993; Edmonton, Alberta, Canada. p. 281-316.
 8. Fuller R. A review: probiotics in man and animals. Journal of Applied Bacteriology. 1989;66(4):365-378.
 9. Gebrehiwot T. Rumen manipulation for enhanced feed utilization and improved productivity performance of ruminants: a review. Momona Ethiopian Journal of Science. 2014;6(2):3-17.
 10. Gworgwor Z, Mbahi TF, Yakubu B. Environmental implications of methane production by ruminants: a review. Journal of Sustainable Development in Agriculture and Environment. 2006;2(1):794-867.
 11. Hess HD, Beuret RA, Lötscher M, Hindrichsen IK, Machmüller A. Ruminant fermentation, methanogenesis and nitrogen utilization of sheep receiving tropical grass hay-concentrate diets offered with *Sapindus saponaria* fruits and *Cratylia argentea* foliage. Animal Science. 2004;79(2):177-189.
 12. Jansen DH. Ecology of plants in the tropics. In: Arnold E, editor. Studies in Biology. UK; 1975. p. 66.
 13. Kamra DN. Rumen microbial ecosystem. Current Science. 2005;89(1):134-135.
 14. MacRae JC, Lobley GE. Some factors which influence thermal energy loss during the metabolism of ruminants. Livestock Production Science. 1982;9(3):477-479.

How to Cite This Article

Deepika B, Ganesan R, Kandasamy R, Siva M, Senthilkumar S. Recent approaches in rumen fermentation manipulation for sustainable ruminant production. International Journal of Veterinary Sciences and Animal Husbandry. 2026;11(6):187-190.

Creative Commons (CC) License

This is an open-access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.