



International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
 ISSN Online: 2617-4707
 NAAS Rating (2026): 5.29
 IJABR 2026; 10(6): 1131-1136
www.biochemjournal.com
 Received: 01-04-2026
 Accepted: 04-05-2026

R Suresh

Assistant Professor,
 Department of Veterinary
 Medicine, CVSc, Korutla,
 PVNRTGVU, Hyderabad,
 Telangana, India

M Mahesh Babu

M.V.Sc., Department of
 Animal Nutrition, CVSc,
 Rajendranagar, PVNRTGVU,
 Hyderabad, Telangana, India

B Sravanthi

Ph.D. Scholar, Department of
 Animal Nutrition, CVSc,
 Rajendranagar, PVNRTGVU,
 Hyderabad, Telangana, India

N Nalini Kumari

Professor, Department of
 Animal Nutrition, CVSc,
 Rajendranagar, PVNRTGVU,
 Hyderabad, Telangana, India

G Vinay

M.V.Sc. Scholar, Department
 of Animal Nutrition, CVSc,
 Rajendranagar, PVNRTGVU,
 Hyderabad, Telangana, India

J Adhithya

M.V.Sc. Scholar, Department
 of Animal Nutrition, CVSc,
 Rajendranagar, PVNRTGVU,
 Hyderabad, Telangana, India

A Laxman

M.V.Sc. Scholar, Department
 of Animal Nutrition, CVSc,
 Rajendranagar, PVNRTGVU,
 Hyderabad, Telangana, India

Corresponding Author:**B Sravanthi**

Ph.D. Scholar, Department of
 Animal Nutrition, CVSc,
 Rajendranagar, PVNRTGVU,
 Hyderabad, Telangana, India

Comprehensive ontogeny of the rumen: Microbial succession, molecular regulation, and nutritional programming under global and tropical smallholder paradigms

R Suresh, M Mahesh Babu, B Sravanthi, N Nalini Kumari, G Vinay, J Adhithya and A Laxman

DOI: <https://www.doi.org/10.33545/26174693.2026.v10.i6n.8841>

Abstract

The developmental transition of the neonatal ruminant from a functional pseudo-monogastric state to a mature, multi-compartmental obligate herbivore represents a highly coordinated biological milestone. This transition requires synchronized adaptations across gastric histomorphology, physiological architecture, and microbial ecology. Early nutritional interventions dictate the velocity of this transition, modulating both the physical development of the ruminal wall and the metabolic functionality of its epimural and luminal microbiota. This review synthesizes current knowledge on the structural development of the forestomach, the ecological succession of the four microbial kingdoms, and the underlying molecular, endocrine, and epigenetic signaling networks such as the somatotrophic axis and volatile fatty acid (VFA)-mediated cellular cascades that govern mucosal growth. Furthermore, it examines the practical applications of starter processing and physically effective fiber (peNDF), while addressing key mathematical discrepancies in gastrointestinal tract fill calculations. Finally, this paper provides a targeted analysis of rumen development under tropical smallholder conditions, focusing on India, where climatic stress, high-lignin crop residues, and distinct breed genetics alter standard developmental timelines. This synthesis provides an integrated framework for designing nutritional interventions that optimize long-term production efficiency and health in both temperate and tropical production systems.

Keywords: Rumen development, forestomach maturation, rumen microbiome, volatile fatty acids

Introduction

The economic and physiological sustainability of global ruminant production relies heavily on the management of the post-natal pre-weaning window. At birth, the young ruminant possesses an anatomically underdeveloped forestomach complex, functioning effectively as a pseudo-monogastric organism dependent on liquid milk diets digested exclusively within the abomasum (Diao *et al.*, 2019) [2]. The subsequent transition to solid feed requires a profound physiological and anatomical reorganization. This process is driven by complex interactions between dietary physical forms, microbial fermentation end-products, and host tissue transcriptomic reprogramming (Ghaffari *et al.*, 2025) [5].

Optimizing this developmental window accelerates early growth, lowers rearing costs, and programs the lifetime metabolic efficiency of the adult animal. Conversely, mismanagement during this critical period can lead to structural damage to the ruminal epithelium, systemic metabolic disorders, and delayed physiological maturity.

While fundamental models of rumen ontogeny have been established in temperate, intensive production environments, applying these frameworks directly to tropical regions such as the Indian subcontinent presents unique challenges. Smallholder systems in these areas face high ambient temperatures, structural forage scarcities, and a heavy reliance on low-digestibility crop residues.

This review provides a comprehensive analysis of the anatomical, microbiological, and molecular mechanisms governing rumen development. It evaluates global nutritional strategies and introduces targeted management interventions designed to address the

environmental and nutritional challenges of tropical production systems.

Morphological and Anatomical Ontogeny of the Rumen Gastric Compartment Mass Dynamics

The post-natal development of the ruminant stomach involves major shifts in the relative mass and volume of its four distinct chambers. In the newborn calf or lamb, the abomasum is the largest compartment, accounting for approximately 49% of the total gastric tissue weight, whereas the reticulorumen represents only 32% to 38% (Yu and Wu, 2026) ^[12]. This structural balance is highly temporary.

As the animal consumes solid feed, the forestomachs undergo rapid growth. By 6 to 8 weeks of age, the reticulorumen expands to represent 61.23% to 63% of the total stomach mass, while the abomasum declines to roughly 25.37%. Upon reaching physiological maturity between 12 and 16 weeks of age, the adult profile stabilizes: the reticulorumen dominates at over 70% of the total stomach mass, the omasum constitutes approximately 18%, and the abomasum reduces to a secondary metabolic role at 15% (Yu and Wu, 2026, Li *et al.*, 2024) ^[12, 8].

Histomorphology of the Rumen Wall and Epithelial Stratification

The expansion of the ruminal wall requires synchronized development across its primary histological layers: the outer *tunica muscularis*, the highly vascularized *tunica submucosa*, and the internal *tunica mucosa*. The *tunica mucosa* differentiates into a specialized stratified squamous epithelium divided into four distinct functional strata:

1. **Stratum Basale:** Positioned adjacent to the basement membrane, this layer is packed with active mitochondria. It drives continuous cellular division and supplies the energy needed to sustain the electrochemical gradients that power transmembrane nutrient and electrolyte transport.
2. **Stratum Spinosum:** Located above the basal layer, these cells are heavily involved in regional short-chain fatty acid (SCFA) metabolism. They convert absorbed butyrate into beta-hydroxybutyrate (BHBA), generating local cellular energy.
3. **Stratum Granulosum:** Characterized by tight junction complexes (including claudins and occludins), this layer forms a selective barrier that prevents the systemic translocation of luminal bacteria or endotoxins into deeper tissues.
4. **Stratum Corneum:** The outermost layer, composed of highly keratinized, non-viable cells. This cornified shield protects the underlying tissue from mechanical abrasion by fibrous feed particles and regularly sloughs off into the lumen.

While the *tunica muscularis* expands primarily in response to the physical volume and scratch factor of coarse forage, the development of the mucosal papillae is driven by chemical stimuli. The absorption and metabolic processing of volatile fatty acids (VFAs) act as the primary trigger for papillae elongation and vascularization. Among these, butyrate exerts the most potent stimulatory effect on epithelial mitosis, followed sequentially by propionate and acetate (Ghaffari *et al.*, 2025, Diddeniya *et al.*, 2024) ^[5, 3].

Microbial Colonization and Succession Dynamics

The Prenatal Colonization Controversy

The classic view of gastrointestinal microbiology maintained that the mammalian fetus remains entirely sterile *in utero*, with microbial colonization initiating exclusively during and after birth via exposure to the maternal vaginal canal, skin, and immediate environment. However, the use of high-throughput, next-generation sequencing and multi-omics technologies has challenged this paradigm. Several studies have detected low-density, metabolically active microbial signatures in fetal meconium, amniotic fluid, and placental membranes prior to birth (Yu and Wu, 2026, Mao *et al.*, 2021) ^[12, 9].

This has generated an ongoing scientific debate. Some researchers suggest these findings indicate a baseline prenatal colonization that prepares the fetal gut for post-natal life, while others argue that these low-biomass signals result from external contamination by DNA reagents or sequencing equipment. Resolution of this debate requires rigorous, low-biomass isolation protocols and synchronized negative controls.

Ecological Succession across the Four Microbial Kingdoms

Post-natal colonization of the rumen lumen and epithelium follows a highly structured ecological succession, even before the animal begins consuming solid feed:

1. Bacteria (50% to 80% of total microbial biomass)

Pioneer communities establish within 20 minutes of birth, dominated by facultative anaerobes such as *Streptococcus* and *Enterococcus* spp. (Perez *et al.*, 2024) ^[10]. These organisms consume residual oxygen within the rumen cavity, lowering the redox potential and establishing a strictly anaerobic environment.

Once oxygen is depleted, obligate anaerobes begin to dominate. The microbial community undergoes a major shift at the phylum level, as *Proteobacteria* are progressively replaced by *Bacteroidetes* and *Firmicutes*. At the genus level, *Bacteroides* functions as a key transitional group before *Prevotella* establishes itself as the dominant genus in the mature rumen (Perez *et al.*, 2024) ^[10]. Crucially, specialized cellulolytic and amylolytic strains colonize the ruminal fluid long before the host starts consuming solid feeds, ensuring the necessary enzymatic machinery is present ahead of dietary transitions.

2. Archaea and Methanogens (0.3% to 3% of total population)

Strictly anaerobic methanogens, predominantly *Methanobrevibacter* spp., appear within the first 20 minutes post-birth and achieve adult-like population densities within 10 to 14 days (Diao *et al.*, 2019) ^[12]. Methanogens play a critical biochemical role by utilizing hydrogen (H₂) gas to reduce carbon dioxide (CO₂), formate, or methyl compounds into methane (CH₄).

By continuously consuming free H₂, methanogens maintain an exceptionally low partial pressure of hydrogen within the rumen. This low hydrogen level is thermodynamically required for ruminal bacteria to spontaneously oxidize NADH back into NAD⁺, preventing fermentation stasis. However, this methane production represents a direct loss of 2% to 12% of the host's gross dietary energy intake.

3. Fungi (8% to 12% of total biomass)

Anaerobic fungi colonize the rumen within the first week of life. They serve as primary colonizers of complex plant cell walls, using their rhizoid networks to physically penetrate lignified tissue. Fungi possess a specialized array of highly active cellulases and hemicellulases that break open tough fibers, making them accessible to cellulolytic bacteria.

Fungi lack classic mitochondria and instead utilize specialized organelles called hydrogenosomes for metabolic electron transport. This process generates large quantities of H₂, formate, and acetate. As a result, methanogenic archaea frequently exhibit physical ectosymbiosis, attaching directly to the outer surfaces of fungal structures to capture these substrates immediately.

4. Protozoa (20% to 50% of total biomass)

Eukaryotic protozoa establish later, typically around 2 weeks of age, requiring direct contact with older animals or dams for transmission due to the lack of an environmentally stable spore phase. Flagellates dominate the early neonatal period but are systematically replaced by ciliate protozoa as the rumen matures.

Protozoa engulf and ferment starch granules and plant fibers, stabilizing the ruminal environment by slowing down rapid bacterial starch degradation. They also regulate bacterial populations via predation, which contributes significantly to intra-ruminal nitrogen recycling. Complete removal of protozoa (defaunation) typically reduces overall feed digestion and animal growth metrics by approximately 10%.

Cellular, Hormonal, and Epigenetic Regulatory Networks

The morphological adaptation of the rumen epithelium from a non-absorptive membrane into a highly active transport barrier is governed by a network of endocrine signals and intracellular signaling cascades.

The Somatotrophic GH/IGF-1 Axis

Systemic regulation of mucosal tissue expansion is driven by the growth hormone/insulin-like growth factor axis. Pituitary-derived Growth Hormone (GH) binds directly to growth hormone receptors (GHR) distributed throughout the ruminal tissue (Yu & Wu, 2026) ^[12]. This receptor binding triggers the localized, autocrine/paracrine synthesis of Insulin-Like Growth Factor-1 (IGF-1) and upregulates its corresponding receptor, IGF-1R.

The activation of IGF-1R initiates rapid mitosis and cellular differentiation within the *stratum basale*. This hormonal cascade is highly regulated by local Insulin-Like Growth Factor Binding Proteins (IGFBPs). During active rumen development, tissue expression of IGFBP-5 increases to amplify IGF-1 signaling, while the expression of IGFBP-2, -3, and -6 decreases, removing their baseline inhibition over cell proliferation and suppressing apoptotic pathways (Yu & Wu, 2026) ^[12]. This shift upregulates key transmembrane VFA and electrolyte transporters, such as Sodium-Hydrogen Exchanger-2 (NHE2) and Monocarboxylate Transporter-1, increasing the tissue's total absorptive capacity.

Key Intracellular Signaling Cascades

Four main biochemical pathways coordinate mucosal growth and tissue homeostatic control:

1. The VFA-Mediated RAP1A/GPR41/PPAR Pathway

The physical absorption of acetate, propionate, and butyrate initiates two distinct intracellular signaling loops. First, the presence of these VFAs downregulates the expression of RAP1A, which removes its baseline inhibition on the RAS protein. This activates the downstream RAS/cAMP/MAPK cascade, triggering the transcription of genes required for cellular division.

Second, VFAs act as direct extracellular ligands for the G-protein coupled receptor GPR41. Activating GPR41 turns on the PI3K/AKT/mTOR signaling pathway, upregulating cellular translation and protein deposition. Simultaneously, absorbed VFAs activate Peroxisome Proliferator-Activated Receptors, specifically upregulating PPARG and PPARG post-weaning (Yu & Wu, 2026) ^[12]. This upregulation enhances mitochondrial beta-oxidation and lipid metabolism, allowing the tissue to meet the high energy demands of rapid structural growth.

2. The Wnt/ β -catenin Pathway

During the introduction of starch-rich diets, amylolytic bacterial fermentation produces metabolites such as indole-3-carboxaldehyde. This metabolite acts as an upstream signal that activates the β -catenin signaling pathway. The activation of this pathway induces structural morphogenesis and the physical elongation of the ruminal papillae, expanding the available surface area for nutrient absorption.

3. The Ca⁺² Muscular Hypertrophy Pathway

The ingestion of coarse, fibrous forage creates mechanical friction against the rumen wall, stimulating the microbial production of prostaglandin D2 (PGD2). PGD2 binds to receptors on the smooth muscle cells of the *tunica muscularis*, triggering an intracellular calcium Ca⁺² influx. This calcium influx activates calmodulin-dependent protein kinases, specifically upregulating CAMK2A and CASQ1, which drive muscle fiber hypertrophy and enhance ruminal motility.

4. The Hippo Pathway

To prevent uncontrolled, malignant tissue overgrowth during periods of intense nutritional stimulation, the Hippo signaling pathway functions as a homeostatic size regulator. Operating through the LATS2-YAP1-BIRC5 cellular axis, this pathway monitors mechanical cell-to-cell contact and density. It supports cell proliferation during early growth phases but activates apoptotic checkpoints if over-proliferation threatens the structural integrity of the tissue barrier (Ghaffari *et al.*, 2025) ^[5].

Epigenetic Imprinting and Chromatin Remodeling

Early nutritional inputs can induce long-term changes in host gene expression through epigenetic programming. Fermentation-derived butyrate functions naturally as a histone deacetylase inhibitor (HDACi). By blocking histone deacetylation, butyrate maintains an open, hyperacetylated chromatin structure (particularly at the H3K27ac locus). This open configuration allows transcription factors sustained access to promoter regions, permanently upregulating genes involved in VFA transport and ketogenesis. This epigenetic imprinting helps optimize the animal's long-term metabolic efficiency post-weaning.

Dietary Interventions and Nutritional Programming

Liquid Versus Solid Feed Dynamics

The neonatal ruminant relies on the esophageal (rumoreticular) groove reflex to survive on liquid diets. Environmental and physiological cues, such as the anticipation of nursing, stimulate vagal nerve pathways that cause the muscular folds of the reticular groove to invert and form a temporary channel. This channel directs milk or milk replacer past the immature reticulorumen directly into the abomasum for enzymatic digestion (Diao *et al.*, 2019) [2].

If this reflex fails often due to poor bucket-feeding practices, thermal stress, or disease liquid feed can leak into the undeveloped rumen sac. This condition, known as "rumen drinking," leads to rapid bacterial fermentation of milk carbohydrates, causing severe ruminal and metabolic acidosis, mucosal inflammation, and systemic unthriftiness. While liquid feed bypasses the forestomach, early access to solid starter feed is required to stimulate rumen development. The introduction of dry feed provides mechanical friction that encourages muscular contractions and expands total ruminal volume. More importantly, its fermentation generates the VFAs required to drive mucosal vascularization and papillae growth.

Starter Grain Processing and the Pathophysiology of Parakeratosis

The physical form and processing method of calf starters heavily influence ruminal health. Texturized or pelleted feeds containing whole, rolled, or steam-flaked grains are superior to finely ground meals. For proper rumen development, calf starters should be formulated so that at least 75% of all dietary particles exceed 1190 μm in diameter (Ghaffari *et al.*, 2025, Kang *et al.*, 2023) [5, 7].

Finely ground starters are rapidly fermented by amylolytic bacteria, causing a drop in ruminal pH below 6.0. This acidic environment inhibits beneficial cellulolytic bacteria and promotes the growth of lactic acid producers like *Streptococcus bovis*. The resulting lactic acid accumulation can cause acute or subacute ruminal acidosis (SARA).

At the tissue level, prolonged acid exposure disrupts regular cell differentiation within the stratified squamous epithelium. This leads to epithelial parakeratosis, a condition where the cells of the *stratum corneum* fail to lose their nuclei and form dense, non-functional cell clusters. These layers lose their flexibility, develop deep cracks, and exhibit papillae clumping. This structural breakdown compromises the tight junctions of the *stratum granulosum*, allowing bacteria and endotoxins to enter the portal circulation and cause liver abscesses and systemic inflammation.

Forage Inclusion and Physically Effective Fiber (peNDF)

Including low levels of high-quality forage in early diets can help mitigate the risks of high-starch starters. This strategy operates on the concept of physically effective NDF (peNDF > 4mm). Providing between 7.5% Tifton hay and 10% corn silage on a dry matter (DM) basis within a Total Mixed Ration (TMR) delivers an ideal peNDF level of 0.8% to 3.7% (Toledo *et al.*, 2024) [11].

This physical fiber provides several benefits:

- It creates a mechanical "scratch factor" that clears sloughed keratinized cells from between the mucosal papillae, preventing parakeratosis.

- It stimulates rumination and increases saliva flow, which contains natural bicarbonate and phosphate buffers that stabilize ruminal pH above 6.0.
- It reduces abnormal, non-nutritive oral behaviors, such as hair licking or pen chewing.

However, forage levels must be carefully managed. Because forage has a lower energy density and a higher gut fill effect than grain concentrates, exceeding 15% forage inclusion can limit total dry matter and energy intake, which may delay overall body tissue growth.

The Gut Fill Mathematical Correction Fallacy

In nutrition trials evaluating calf growth, researchers often correct for individual gastrointestinal tract fill using standard equations, such as the *Jahn and Chandler equation*, to determine true empty body weight (EBW) gains. However, this equation was optimized using older cattle (8 to 20 weeks of age) fed high-straw diets.

Recent empirical slaughter data indicate that applying this classic 6% correction factor overestimates true tissue growth in younger calves (Toledo *et al.*, 2024, Gasiorek *et al.*, 2020) [11, 4]. In pre-weaned and newly weaned calves, the actual ratio of gastrointestinal tract fills to total live body weight is closer to 12%. Using this updated 12% baseline ensures more accurate measurements of real structural tissue growth versus transient digesta accumulation.

Environmental Stressors and Management Modalities: Global Perspectives and Indian Tropical Paradigms

Climatic Stressors and Epithelial Integrity

Extreme ambient temperatures can disrupt the physiological development of the rumen. In temperate regions, cold stress shifts energy usage toward metabolic thermogenesis, which can reduce the efficiency of nutrient utilization for structural tissue growth. Conversely, heat stress creates significant challenges for ruminant production globally, particularly during tropical summers.

When animals suffer from severe heat stress, blood flow is diverted from the splanchnic bed to the periphery to assist with cooling. This hyperthermia-induced ischemia deprives the ruminal epithelium of oxygen and nutrients, leading to ATP depletion within the *stratum basale*. As a result, the tight junction proteins within the *stratum granulosum* break down, increasing mucosal permeability. This structural failure allows luminal endotoxins (lipopolysaccharides, LPS) to enter the bloodstream, triggering systemic inflammatory responses and metabolic issues.

The Indian Ruminant Production Paradigm

Applying intensive Western calf rearing models directly to the Indian livestock sector is difficult due to several distinct environmental and socio-economic factors:

1. Smallholder Micro-Economics and Management Challenges

Over 70% of India's milk is produced by smallholder and landless farmers managing small herds of 2 to 5 animals. In these systems, commercial texturized calf starters and high-quality milk replacers are often economically impractical or unavailable.

Calves are frequently raised using restricted maternal suckling protocols, which prioritizes milk for human consumption and commercial sale. This restriction often

limits early nutrient intake, delaying both general structural growth and the development of the forestomach. Furthermore, widespread individual tethering and a lack of group housing eliminate the benefits of social facilitation, which has been shown to encourage early solid feed intake.

2. Breed Genetics (*Bos indicus* vs. *Bos taurus* crossbreds)

The Indian dairy herd consists of indigenous Zebu (*Bos indicus*) cattle, crossbred cattle (*Bos taurus* X *Bos indicus*), and water buffaloes (*Bubalus bubalis*). Indigenous Zebu breeds and water buffaloes exhibit unique physiological traits, including lower baseline maintenance energy requirements and higher resistance to heat stress compared to purebred *Bos taurus* animals.

However, crossbred calves (such as Holstein-Friesian or Jersey crosses with Sahiwal or Kankrej) often lack these protective mechanisms. They are highly susceptible to the high Temperature-Humidity Index (THI) typical of Indian summers, which frequently results in depressed dry matter intake, altered ruminal fermentation, and delayed weaning maturity.

3. Low-Digestibility, High-Lignin Crop Residues

Due to competition for arable land between human food crops and animal forage, young ruminants in India are commonly transitioned onto diets based on crop residues, such as paddy straw (*Oryza sativa*) or wheat straw (*Triticum aestivum*). These straws are characterized by high concentrations of heavily lignified NDF and silica fractions, along with low crude protein levels (< 4%).

Transitioning young animals directly onto these high-lignin residues presents several physiological challenges:

- The lack of readily fermentable non-fiber carbohydrates (NFC) limits the production of propionate and butyrate, which are necessary to stimulate papillae growth.
- The high physical fiber content fills the immature rumen cavity, reducing total dry matter intake through gut fill limitations.
- Low dietary nitrogen restricts the growth of the newly establishing ruminal microbiota, delaying the maturation of normal fermentation kinetics.

Targeted Nutritional Interventions for Indian Conditions

To improve rumen development within these constraint-driven systems, several targeted, low-cost nutritional strategies can be applied:

- **Complete Feed Blocks (CFB):** Compacting finely chopped straw (e.g., wheat or paddy straw) with precise levels of molasses, urea, and locally available oil cakes (such as mustard or cotton seed cake) into high-density blocks provides several benefits (Haloi *et al.*, 2021) [6]. The complete feed system provides a quantitative mixture of all dietary ingredients, blended thoroughly to prevent separation and selection, fed as a sole source of nutrients (Beigh *et al.*, 2016) [1]. This practice stabilizes rumen fermentation, minimizes fermentation loss, and ensures more uniform load on the rumen with less fluctuation in the release of ammonia, which supports more efficient utilization of ruminal non-protein nitrogen (Beigh *et al.*, 2016; Haloi *et al.*, 2021) [1, 6].
- **Strategic Green Forage Supplementation:** Introducing small, controlled amounts of high-quality local leguminous green forages such as berseem

(*Trifolium alexandrinum*), cowpea (*Vigna unguiculata*), or lucerne (*Medicago sativa*) at 10% of total dry matter intake can help balance crop-residue diets. These forages provide highly bioavailable protein, vitamins, and structural fiber that support optimal ruminal pH and encourage papillae growth without causing gut fill limitations.

- **Solid-State Fermentation (SSF) and Crop Residues:** Pre-treating poor-quality straws with white-rot fungi (such as *Pleurotus ostreatus*) or commercial fibrolytic enzyme complexes can help break down complex lignin-cellulose bonds before feeding. This pre-digestion increases the straw's total soluble carbohydrate availability, improving VFA production and accelerating rumen development.
- **Direct-Fed Microbials (DFM) and Prebiotics:** Supplementing diets with live yeast cultures (*Saccharomyces cerevisiae*) or specific postbiotics can help improve ruminal development under high-temperature conditions (Beigh *et al.*, 2016) [1]. Live yeast consumes residual oxygen within the rumen, helping to maintain a low redox potential and supporting the growth of beneficial fiber-digesting bacteria (Beigh *et al.*, 2016) [1]. This stabilization helps maintain consistent fermentation rates and protects epithelial health during periods of environmental heat stress.

Conclusions and Future Perspectives

The development of the ruminant forestomach is a dynamic biological process regulated by complex physiological, microbiological, and endocrine mechanisms. Transitions in dietary forms from liquid milk to solid feed induce a complete rearrangement of gastric tissue mass, modify the stratification of the squamous epithelium, and guide the ecological succession of the ruminal microbiota. These structural modifications are supported at the molecular level by the somatotrophic axis and VFA-driven intracellular signaling cascades, which balance cell proliferation with tissue homeostasis.

While intensive management strategies using precise starter grain processing and physically effective fiber are well established in temperate regions, their direct application to tropical smallholder systems requires careful adjustment. Addressing the challenges of high climatic heat stress, low-quality crop residues, and limited resource availability requires targeted nutritional solutions.

Implementing practical interventions such as complete feed blocks, pre-treatment of straws, and strategic supplementation with local green legumes can help optimize rumen development in these environments. Future research should focus on identifying the specific epigenetic mechanisms driven by local tropical feedstuffs, allowing for the design of targeted nutritional programs that improve lifetime performance, health, and climate resilience in diverse production systems.

References

1. Beigh YA, Ganai AM, Ahmad HA. Prospects of complete feed system in ruminant feeding: A review. Vet World. 2017;10(4):424-437.
2. Diao Q, Zhang R, Fu T. Review of strategies to promote rumen development in calves. Animals. 2019;9(8):490-508.

3. Diddeniya G, Ghaffari MH, Hernandez-Sanabria E, Guan LL, Malmuthuge N. Invited review: Impact of maternal health and nutrition on the microbiome and immune development of neonatal calves. *J Dairy Sci.* 2024;107(10):7504-7519.
4. Gasiorek M, Stefanska B, Pruszyńska-Oszmalek E, Taciak M, Komisarek J, Nowak W. Effect of oat hay provision method on growth performance, rumen fermentation and blood metabolites of dairy calves during preweaning and postweaning periods. *Animal.* 2020;14(10):2054-2062.
5. Ghaffari MH, Hammon HM, Koch C. Early rumen development in calves: Biological processes and nutritional strategies—A mini-review. *JDS Commun.* 2025;6(3):427-431.
6. Haloi S, Bhuyan R, Borah L, Saikia BN. Complete feed block as a mode of enhancing ruminant production: A review. *Indian J Anim Health.* 2021;60(1):10-15.
7. Kang X, Li C, Liu S, Baldwin RL, Liu GE, Li CJ. Genome-wide acetylation modification of H3K27ac in bovine rumen cell following butyrate exposure. *Biomolecules.* 2023;13(7):1137-1150.
8. Li B, Wu K, Duan G, Yin W, Lei M, Yan Y, *et al.* Folic acid and taurine alleviate the impairment of redox status, immunity, rumen microbial composition and fermentation of lambs under heat stress. *Animals.* 2024;14(7):998-1014.
9. Mao H, Zhang Y, Yun Y, Ji W, Jin Z, Wang C, *et al.* Weaning age affects the development of the ruminal bacterial and archaeal community in *Hu* lambs during early life. *Front Microbiol.* 2021;12:636865.
10. Perez HG, Stevenson CK, Lourenco JM, Callaway TR. Understanding rumen microbiology: An overview. *Encyclopedia.* 2024;4(1):148-157.
11. Toledo AFD, Silva APD, Barbosa FVL, Barboza RDF, Oliveira ICR, Marino ED, *et al.* Forage sources in total mixed rations on rumen fermentation, gut fill, and development of the gastrointestinal tract of dairy calves. *Sci Rep.* 2024;14(1):30762.
12. Yu G, Wu D. Research advances in rumen development and regulatory mechanisms in young ruminants. *PeerJ.* 2026;14:e20626.