



International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693
ISSN Online: 2617-4707
NAAS Rating (2026): 5.29
IJABR 2026; SP-10(9): 1087-1099
www.biochemjournal.com
Received: 01-07-2026
Accepted: 02-08-2026

T Balaji Goud
Department of Genetics and
Plant Breeding, S.V.
Agricultural College, Tirupati,
Andhra Pradesh, India

Dr. N Sabitha
Associate Professor, S.V.
Agricultural College, Tirupati,
Andhra Pradesh, India

Dr. I Sudhir Kumar
Senior Scientist (Plant
Breeding), Agricultural
Research Station,
Peddapuram, Andhra Pradesh,
India

**Dr. Chikkappa Gangadhar
Karjagi**
Principal Scientist, Regional
Maize Research and Seed
Production Centre, ICAR-
Indian Institute of Maize
Research, Begusarai, Bihar,
India

Dr. M Raghavendra
Department of Soil Science and
Agricultural Chemistry, S.V.
Agricultural College, Tirupati,
Andhra Pradesh, India

Dr. G Mohan Naidu
Department of Statistics and
Computer Applications, S.V.
Agricultural College, Tirupati,
Andhra Pradesh, India

Corresponding Author:
Dr. N Sabitha
Associate Professor, S.V.
Agricultural College, Tirupati,
Andhra Pradesh, India

Pleiotropic master regulators in crop plants: Mechanisms and strategies for multi-stress tolerance and biofortification

**T Balaji Goud, N Sabitha, I Sudhir Kumar, Chikkappa Gangadhar
Karjagi, M Raghavendra and G Mohan Naidu**

DOI: <https://www.doi.org/10.33545/26174693.2026.v10.i9Sn.9965>

Abstract

Modern agricultural systems face the dual pressure of ensuring global nutritional security while mitigating the devastating impacts of climate-driven abiotic and biotic stresses. Traditional breeding approaches, which often address isolated phenotypic traits, frequently overlook the complex, intertwined nature of plant genetic architecture. This review synthesizes the critical role of pleiotropic master regulators genes that exert simultaneous control over multiple physiological and metabolic pathways as foundational tools for crop improvement. By examining high-value loci such as *IPAI* in rice, *tb1* in maize and *NAM-B1* in wheat, we demonstrate that precise manipulation of these hubs allows for the decoupling of negative physiological trade-offs, enabling the concurrent enhancement of grain yield, nutrient density and stress resilience. Furthermore, we evaluate the transition from conventional backcrossing to advanced biotechnological interventions, including CRISPR-mediated promoter engineering and integrated multi-omics, which facilitate the surgical tuning of genetic networks. As the scientific community pivots toward *de novo* domestication and precision breeding, the management of these complex pleiotropic networks emerges as the most viable strategy to develop climate-resilient, biofortified staple crops capable of meeting the compounding agronomic challenges of the 21st century and permanently alleviating the global hidden hunger epidemic.

Keywords: Pleiotropy, biofortification, abiotic stress, CRISPR-Cas9, nutritional security, linkage drag, multi-omics integration, epistasis

1. Introduction

The global agricultural sector currently faces a convergence of severe environmental, nutritional and population-driven pressures. The rapid expansion of the human population necessitates significant increases in food yields, a mandate that is heavily undermined by erratic climatic shifts. These climatic variations increase the likelihood and intensity of simultaneous abiotic and biotic stresses in farming environments (Mickelbart *et al.*, 2015; Husaini, 2022). In natural field conditions, crops are frequently subjected to multiple overlapping stressors rather than isolated adverse events (Hasanuzzaman *et al.*, 2012). Because the biological mechanisms governing stress tolerance are highly complex, managing multiple simultaneous threats poses a profound challenge to plant systems (Bailey-Serres *et al.*, 2019). To confront this systemic vulnerability, major global organizations including the Food and Agriculture Organization (FAO), the International Fund for Agricultural Development (IFAD), the United Nations World Food Programme (WFP) and the Consultative Group on International Agricultural Research (CGIAR) convened a High-Level Conference on World Food Security in 2008. During this event, 181 nations ratified a declaration underscoring the vital need to enhance the structural resilience of modern food systems against climate-induced disruptions (Bailey-Serres *et al.*, 2019) [3, 32, 58].

Changing weather patterns not only drive the emergence of new agricultural diseases but also exacerbate the loss of genetic diversity in global crops, thereby diminishing their natural evolutionary defenses (Husaini, 2022). Beyond raw yield reductions, there is an escalating qualitative crisis categorized as "hidden hunger" (Yadava *et al.*, 2018) [39, 94]. Past agricultural initiatives, most notably the Green Revolution, succeeded in elevating caloric production but often neglected micronutrient density, resulting in pervasive nutritional deficiencies

worldwide (Yadava *et al.*, 2018) ^[94]. Thus, contemporary agricultural frameworks must transition away from a strictly yield-centric model toward one that ensures comprehensive nutritional security, especially in degrading ecosystems.

Limitations of Conventional Crop Improvement Programs: Despite the pressing need to develop cereal varieties that offer both high nutritional value and robust environmental adaptability, traditional plant breeding faces significant hurdles in concurrently optimizing these distinct traits (Hawkesford *et al.*, 2013). The frequent generation of lines that are nutritionally dense but vulnerable to drought or conversely, drought-resilient but nutritionally deficient highlights a fundamental limitation in orthodox breeding methodologies (Bouis & Saltzman, 2017). Three primary physiological and genetic hurdles underpin this ongoing shortfall ^[9, 33]:

Compartmentalized Breeding Objectives

Historically, agricultural research programs have operated in silos. Initiatives aimed at bolstering stress tolerance and those focused on biofortification have largely remained separate, yielding crop varieties that excel in isolated metrics but fail to perform holistically in complex environments (Horton *et al.*, 2018; Saltzman *et al.*, 2013). This fragmented approach struggles to provide the comprehensive genetic architectures necessary for confronting multi-stress field conditions (Tester & Langridge, 2010) ^[35, 70, 82].

Resource Allocation and Metabolic Penalties

Intrinsic physiological trade-offs further impede dual-trait optimization. When subjected to acute stress, plants initiate survival mechanisms that divert metabolic energy away from reproductive development, consequently diminishing grain fill rates and the deposition of essential micronutrients (Cakmak, 2008; Chapin *et al.*, 1990). As a result, genotypes capable of surviving severe environmental shocks often suffer from reduced yields and poorer nutritional quality due to disrupted source-sink dynamics (Foyer *et al.*, 2016) ^[12, 15, 26].

Linkage Drag and Genetic Confounding

The introduction of stress-resilience genes from wild relatives into elite commercial lines frequently drags along tightly linked, unfavorable alleles. This phenomenon often creates an illusion of negative pleiotropy, where enhanced resilience appears intrinsically tied to yield or quality penalties (Fridman *et al.*, 2004; Tanksley & McCouch, 1997). In reality, these detrimental correlations are largely the product of intense linkage disequilibrium (LD) rather than a singular gene exerting multiple adverse effects (Mackay *et al.*, 2009) ^[27, 51, 81].

Standard bi-parental backcrossing lacks the precision and speed required to sever these tight genetic linkages. To overcome these structural barriers, researchers increasingly rely on highly recombined mapping panels, such as Multi-parent Advanced Generation Inter-Cross (MAGIC) and Nested Association Mapping (NAM) populations, to break deleterious associations and isolate genuine pleiotropic regulatory hubs (Cavanagh *et al.*, 2008; Yu *et al.*, 2008) ^[14, 99].

1.2 Addressing Quantitative Complexity via Pleiotropy

The multifaceted challenges of agricultural adaptation are exacerbated by the quantitative nature of critical agronomic traits. Characteristics like overall yield and climatic resilience are governed by numerous quantitative trait loci (QTLs), each contributing minor effects that fluctuate based on environmental interactions (Liu *et al.*, 2012). Overcoming this complexity requires advanced genetic interventions to design crop ideotypes capable of withstanding heat, salinity, drought, flooding and pathogenic pressures without sacrificing productivity (Eckardt *et al.*, 2023). Systematically developing these biofortified, resilient cultivars represents a critical pathway toward sustainable farming (Yadava *et al.*, 2018) ^[23, 49, 94].


Rather than painstakingly pyramiding individual genes a practice that lengthens breeding cycles and elevates the risk of linkage drag modern strategies focus on exploiting high-value pleiotropic loci. These single genetic elements hold the capacity to concurrently regulate multiple advantageous characteristics. Targeting a unified control point allows breeders to manipulate various physiological pathways simultaneously, bypassing the biological trade-offs that have traditionally stymied agricultural progress (Dwivedi *et al.*, 2024; Husaini, 2022) ^[22].

The primary strategic benefit of pleiotropy is the ability to enhance several traits at once without incurring physiological penalties. Specific regulatory loci can be activated to simultaneously deploy defensive structures, trigger immune responses and stimulate nutritional accumulation pathways (Husaini, 2022). Notable candidate genes like *OsDREB1A* and *Osmotin* have demonstrated the ability to foster multi-stress endurance while simultaneously boosting yields and nutritional density (Husaini, 2022).

For instance, pinpointing a single pleiotropic QTL can trigger mechanisms for moisture conservation while simultaneously enhancing the metabolic pathways responsible for loading micronutrients into the developing seed (Husaini, 2022; Ren *et al.*, 2022). This structural efficiency greatly streamlines introgression efforts. By negating the need for complex gene pyramiding, the utilization of single-locus hubs drastically simplifies the target genetic architecture, enabling the rapid development of superior crop lines (Dwivedi *et al.*, 2024; Husaini, 2022). Mapping these specific networks offers a reliable, non-transgenic blueprint for securing food safety, combating global malnutrition and supporting long-term agricultural stability (Bailey-Serres *et al.*, 2019; Shahzad *et al.*, 2024) ^[3, 22, 39, 73].

1.3 The Genetic Architecture of Pleiotropy

Exploiting pleiotropy requires a clear delineation of its functional mechanisms compared to other inheritance patterns.



1.3.1 Differentiating Pleiotropy from Polygenic Inheritance: Breeding programs must clearly distinguish between biological pleiotropy and polygenic traits, as they operate via inverse structural pathways (Dwivedi *et al.*, 2024).

The Polygenic Model ("Many-to-One")

Polygenic traits rely on the cumulative action of multiple, disparate genes to express a single continuous phenotype, such as baseline yield or plant height (LibreTexts Genetics, 2021). Because numerous loci must be manipulated simultaneously to achieve significant phenotypic alterations, breeding for these traits is notoriously slow and resource-intensive (Dwivedi *et al.*, 2024; LibreTexts Genetics, 2021).

The Pleiotropic Model ("One-to-Many")


Conversely, pleiotropy functions as a centralized regulatory switch (Jordan *et al.*, 2019). A single master locus dictates multiple, distinct physiological outcomes. This allows geneticists to initiate a broad cascade of advantageous phenotypic adjustments by targeting just one locus, making these genes highly sought after for complex trait integration (Dwivedi *et al.*, 2024; Jordan *et al.*, 2019; LibreTexts Genetics, 2021) [22].

Structural Sub-Classifications

At the molecular tier, pleiotropy generally arises when a single molecular function cascades into disparate downstream outcomes (He & Zhang, 2006). This takes three primary forms (Jordan *et al.*, 2019) [41]:


Vertical Pleiotropy

A cascading effect wherein a gene alters a primary phenotype, which subsequently serves as the biological foundation for a secondary trait (e.g., deeper root architecture leading to improved mineral uptake).




Horizontal Pleiotropy

True biological pleiotropy, where a single master gene independently and simultaneously initiates multiple distinct pathways, providing highly robust multi-trait enhancement.



Linkage Disequilibrium (Spurious Pleiotropy)

An artifact where two distinct genes governing different traits are located exceptionally close to one another on a chromosome. Overcoming this false pleiotropy demands heavy recombination to physically separate the linked traits (Solovieff *et al.*, 2013) [75].



Functional Validation Strategies

Before deploying a target locus into elite germplasm, its pleiotropic authenticity must be rigorously confirmed across various genetic backgrounds to ensure it does not harbor concealed yield penalties.

Mutant Crosses and Knock-out Verification

The deployment of targeted knock-out genotypes within stable genetic lines remains a gold standard for assessing pleiotropy (Dudley *et al.*, 2005) [21]. By inducing a loss-of-function mutation at the target locus, scientists can verify if multiple traits collapse concurrently, thereby validating the presence of a centralized regulatory node.


Recombination-Based Validation

Using high-resolution mapping populations enables breeders to definitively separate true horizontal pleiotropy from tight linkage disequilibrium. This methodology prevents the introduction of unfavorable physiological trade-offs and accelerates the creation of resilient crops using conventional and precision breeding techniques.

Advanced Mapping Populations: The Recombination Toolkits: Distinguishing authentic pleiotropy from spurious LD is a foundational challenge in quantitative genetics (Mackay *et al.*, 2009; Solovieff *et al.*, 2013). Older mapping techniques, such as standard Recombinant Inbred Lines (RILs), suffer from vast linkage blocks that artificially group distinct traits together, producing false positive signals (Cavanagh *et al.*, 2008; Jordan *et al.*, 2019). Modern analysis requires sophisticated multi-parent mating designs (Yu *et al.*, 2008) [14, 41, 51, 75, 99].


Nested Association Mapping (NAM)

NAM architectures merge the statistical robustness of traditional linkage mapping with the high-resolution advantages of genome-wide association studies (Yu *et al.*, 2008) [99]. By hybridizing a well-characterized central parent with diverse global founder lines, NAM minimizes population stratification while accumulating historical recombination events. This design empowers researchers to isolate precise QTLs governing both resilience and nutritional quality (McMullen *et al.*, 2009; Tian *et al.*, 2011) [55, 84].



Multi-parent Advanced Generation Inter-Cross (MAGIC)

MAGIC schemas utilize a highly complex, rotating hybridization strategy involving multiple founders (Bandillo *et al.*, 2013; Cavanagh *et al.*, 2008). This forces the rapid degradation of long-range linkage disequilibrium, shrinking genomic linkage blocks down to a near-gene scale (Dell'Acqua *et al.*, 2015). When analyzed across varied environmental pressures, MAGIC populations expose true horizontal pleiotropy, yielding verified master hubs primed for commercial introgression (Huang *et al.*, 2015) [4, 14, 18, 37].



The Functional Interplay of Pleiotropy in Cross-Tolerance: High-value pleiotropic loci function as sophisticated metabolic switchboards, orchestrating communication across disparate biological networks.

Hormonal Crosstalk and Transcriptional Networks

The mechanism underpinning horizontal pleiotropy frequently involves upstream transcription factors binding to conserved cis-regulatory sequences present across diverse downstream pathways (He & Zhang, 2006). These loci adeptly manage environmental and pathogenic responses by balancing signaling hormones like Abscissic Acid (ABA) against defense hormones such as Jasmonic Acid (JA) and Salicylic Acid (SA) (Fujita *et al.*, 2006; Husaini, 2022) [29, 39].

ROS Scavenging Mechanisms

Nearly all acute stresses trigger an influx of highly destructive Reactive Oxygen Species (ROS) within plant cells (Hasanuzzaman *et al.*, 2012; Mittler, 2017). Master regulatory genes mitigate this by initiating systemic antioxidant cascades, synthesizing critical enzymes like Superoxide Dismutase (SOD) and Catalase (CAT) to neutralize oxidative damage (Gill & Tuteja, 2010; Mittler *et al.*, 2004). This single intervention safeguards the plant against extreme weather and cellular degradation simultaneously (Residiwati *et al.*, 2021) [31, 32, 60, 61].

Nutrient-Mediated Immunity

An advanced facet of pleiotropy involves the coupling of systemic immunity with enhanced root mineral acquisition (Datnoff *et al.*, 2007). Pleiotropic networks that improve the plant's capacity to forage and translocate essential micronutrients concurrently furnish the cellular building blocks required for immune defense enzymes. This dual-action pathway inherently links long-term biofortification with enhanced physical survival in the field (Cabot *et al.*, 2019; Cakmak, 2008; Dwivedi *et al.*, 2024; Husaini, 2022) [11, 12, 17, 39].

Systems-Level Discovery Mechanisms

Transitioning beyond single-trait observational studies requires multi-dimensional biological methodologies to capture the breadth of pleiotropic expression (Dwivedi *et al.*, 2024; Mackay *et al.*, 2009; Solovieff *et al.*, 2013) [22, 51, 75].

Multi-Trait GWAS (MT-GWAS)

MT-GWAS applies sophisticated multivariate linear mixed models to concurrently evaluate multiple phenotypic readouts (Korte *et al.*, 2012; Zhou & Stephens, 2014). Analyzing variance-covariance matrices allows geneticists to locate specific chromosomal regions that co-vary with disparate traits simultaneously, drastically improving statistical validity (Ren *et al.*, 2022; Stephens, 2013) [44, 78, 100].

Transcriptomic Network Analysis

Exposing plants to concurrent stresses and conducting temporal RNA-sequencing enables the application of Weighted Gene Co-expression Network Analysis (WGCNA) (Langfelder & Horvath, 2008; Mittler, 2017). This technique effectively groups co-expressed networks to pinpoint specific transcription factors that simultaneously direct defense and nutrient loading operations (He & Zhang, 2006; Mao *et al.*, 2009) [47, 53, 60].

Expression QTL (eQTL) Integration

eQTL strategies identify distant regulatory hotspots (*trans*-eQTLs) that exert wide-reaching authority over expansive metabolic networks, linking localized chromosomal signatures with broad phenotypic changes (Albert & Kruglyak, 2015) [1].

Reverse Genetics

Validating candidate loci heavily relies on mutagenic libraries created via techniques like Targeting Induced Local Lesions in Genomes (TILLING) (Kurowska *et al.*, 2011; McCallum *et al.*, 2000). Identifying individuals with singular point mutations that display a concurrent loss of multiple target traits confirms the existence of a high-value pleiotropic regulator (Dudley *et al.*, 2005) [21, 46, 54].

Precision Introgression and Recombination Breeding

Once validated, these pivotal genetic assets must be meticulously transferred into established commercial varieties (Hospital, 2009) [36].

Marker-Assisted Backcrossing (MABC)

Employing high-density molecular markers, breeders perform foreground selection to trace the pleiotropic locus while simultaneously executing background selection to rapidly reinstate the commercial parent's native genome, effectively circumventing linkage drag (Frisch & Melchinger, 2005; Hospital, 2009; Tanksley & McCouch, 1997) [28, 36, 81].

Genomic Selection (GS)



This advanced methodology computes Genomic Estimated Breeding Values (GEBVs) to account for the total genetic background (Bernardo & Yu, 2007; Meuwissen *et al.*, 2001). By weaving target markers into predictive statistical models, breeders can lock in major multi-trait hubs without disrupting the plant's broader polygenic yield capacity (Dwivedi *et al.*, 2024; Spindel *et al.*, 2015) [6, 22, 57, 77, 99].

High-Resolution Haplotype Tuning

Generating intense localized recombination across upstream regulatory sequences allows breeders to selectively capture elite promoter haplotypes (Huang *et al.*, 2015; Salvi & Tuberosa, 2005; Tian *et al.*, 2011). This permits non-transgenic optimization of gene expression timing and spatial distribution (Husaini, 2022) [37, 39, 71, 84].

Biofortification: A Structural Approach to Nutrition

Tackling widespread micronutrient deficiencies demands a shift from temporary supplementation to permanent agronomic frameworks (Bouis & Welch, 2010). Biofortifying primary staple crops provides an inherently sustainable, highly scalable method for delivering critical nutrition directly into the diets of susceptible demographics (Garg *et al.*, 2018) [9, 30].



However, altering these nutritional pathways inevitably engages pleiotropic networks, requiring a delicate biochemical balance:

Elevating Beneficial Compounds

Initiatives aim to significantly increase the concentrations of key minerals (Fe, Zn), essential vitamins and rate-limiting amino acids like tryptophan and lysine (Mertz *et al.*, 1964). This process frequently triggers unintended metabolic shifts. For example, augmenting provitamin-A via the *crtRB1* gene inherently alters downstream ABA synthesis, modifying the plant's innate response to drought (Yan *et al.*, 2010). Likewise, driving minerals into wheat grains via *NAM-B1* forcibly accelerates the senescence of vegetative tissues (Uauy *et al.*, 2006) [56, 86, 95].

Suppressing Anti-Nutritional Elements


It is equally vital to diminish compounds that inhibit human nutrient absorption, such as phytic acid. Yet, silencing these pathways (e.g., through *lpa* variants) routinely depresses overall seed viability and agronomic vigor, manifesting as severe negative pleiotropy (Raboy, 2001) [67].

Strategies for Managing Nutritional Pleiotropy Agronomic Methodologies


Utilizing external practices such as direct soil inoculation, foliar nutrient sprays and seed priming to temporarily bypass genetic hurdles and boost nutrient density (Cakmak, 2008) [12].

Advanced Recombination Breeding

The primary long-term solution. This involves exploring natural allelic diversity and executing precise cross-hybridizations to aggressively fracture linkage blocks (Bouis & Welch, 2010) [9]. By securing rare recombination events, breeders can successfully untangle high nutritional profiles from deleterious physiological side effects, creating stable, biofortified elite lines.



2. Comprehensive Catalog of Pleiotropic Master Regulators



Rice (*Oryza sativa*): Due to its relatively compact diploid genome and self-pollinating nature, rice is the premier model organism for elucidating genetic regulatory hubs.

***IPA1 / OsSPL14*:** The *OsSPL14* (*IPA1/WFP*) transcription factor optimizes rice architecture and yield by reducing vegetative tillering and significantly boosting reproductive panicle branching. Mechanistically, *OsSPL14* is regulated post-transcriptionally by the microRNA OsmiR156. During vegetative growth, high OsmiR156 levels cleave *OsSPL14* transcripts to permit normal tillering; during early panicle development, OsmiR156 is repressed, allowing *OsSPL14* accumulation to promote branching. Specific mutations, such as a C-to-A point mutation in the *OsmiR156*-binding site of the Aikawa 1 cultivar, prevent this microRNA-directed cleavage. This leads to abnormal *OsSPL14* accumulation that severely represses axillary bud development while promoting massive panicle branching, increasing total grain yield by up to 52% (Miura et al., 2010; Wang et al., 2018) ^[62, 89].

Ghd7 & Ghd8 / DTH8

The *Ghd7* locus encodes a CCT domain protein that regulates rice heading date, plant height and grain yield. Under long-day conditions, light enhances *Ghd7* expression, causing the protein to act as a robust floral repressor in the nucleus. It specifically suppresses *Ehd1*, which in turn downregulates the *Hd3a* mobile flowering signal, delaying the transition to heading. This extended vegetative growth phase promotes extensive cell proliferation, driving major increases in stem thickness, panicle size and overall grain number. Conversely, natural nonfunctional alleles remove this repression, enabling early heading and allowing rice cultivation to expand into high-latitude, shorter-season environments (Xue et al., 2008; Yan et al., 2011)^[93].

 Q_{SDST}

The *DST* (drought and salt tolerance) transcription factor is a primary negative regulator of osmotic and salinity endurance

in rice. Under favorable conditions, this C2H2-type zinc finger protein binds to a specific promoter motif (DBS) to transactivate genes like *Peroxidase 24 precursor*, which actively neutralizes hydrogen peroxide. However, environmental stress rapidly downregulates *DST* transcription, halting peroxidase production. The resulting accumulation of hydrogen peroxide acts as a secondary messenger that triggers wide-scale stomatal closure. Consequently, *dst* loss-of-function mutations dramatically improve drought and salt survival by permanently reducing stomatal density and keeping apertures closed, limiting transpirational water loss and toxic sodium migration without compromising yield (Huang et al., 2009) ^[38].

***OsDREB1A*:** The *OsDREB* (dehydration-responsive element-binding) family encodes vital transcription activators that orchestrate multi-stress endurance in plants. Environmental triggers activate distinct family members: low temperatures stimulate *OsDREB1A* and *OsDREB1B*, while water deficits and salinity activate *OsDREB2A*. These regulatory proteins migrate to the nucleus and bind specifically to DRE/CRT (dehydration-responsive element/C-repeat) promoter motifs in target genes. This binding drives robust transactivation of a broad stress-inducible gene network, initiating physiological and biochemical adjustments that stabilize cellular structures and optimize osmotic balance, culminating in significantly elevated tolerance to drought, cold and high-salt conditions (Dubouzet et al., 2003) ^[20].

***GS3* & *GW2*:** The *GS3* and *GW2* loci act as classic negative regulators of panicle architecture and grain geometry. The wild-type *GS3* encodes a transmembrane protein that functions as a gamma subunit in G-protein signaling to actively halt longitudinal grain lengthening. Concurrently, the wild-type *GW2* operates as a RING-type E3 ubiquitin ligase that degrades cell-cycle promotion proteins to constrain lateral cell expansion. Domesticated loss-of-function mutations in these loci disrupt their respective repressive pathways, leading to unchecked longitudinal and latitudinal cell proliferation. This combined cellular expansion significantly thickens and enlarges the floral hulls, dramatically increasing grain length, width and individual panicle weight to maximize overall crop productivity (Fan et al., 2006; Song et al., 2007) ^[25, 76].

OsPRR37: The *OsPRR37* (*LHD7*) gene regulates reproductive maturity and crop adaptability in rice. While typically a repressor, in specific genetic backgrounds (like *japonica* Zhonghua 11) under long days, *OsPRR37* paradoxically acts to accelerate flowering. It achieves this by directly binding and suppressing the expression of the prominent floral repressor *Ghd7*. This targeted silencing relieves chronic repression on the flowering pathway, triggering a massive downstream upregulation of *Ehd1*, *Hd3a* and *RFT1* mobile signals. By accelerating the vegetative-to-reproductive phase transition, active *OsPRR37* induces significantly earlier heading dates, resulting in minor reductions in plant height and overall grain yield to ensure optimal environmental synchronization (Koo et al., 2013) [43].

2.2 Maize (*Zea mays*): The historical open-pollination framework of maize creates a rapid decay in linkage blocks,

providing an ideal system for high-resolution genetic decoupling.

tb1

The *teosinte branched1* (*tb1*) gene is a major evolutionary locus selected during maize domestication to control apical dominance, converting the highly branched wild teosinte into a modern single-stalk architecture. It encodes a nuclear transcription factor that represses organ growth. In teosinte, *tb1* expression is restricted to secondary axillary meristems, allowing primary branches to elongate. During domestication, human selection targeted a cis-regulatory variant that expanded *tb1* expression directly into primary axillary meristems and doubled transcript accumulation. This heightened expression strongly arrests primary branch elongation, forcing them to develop into short shoots capped by female inflorescences (ears) instead of elongated branches (Doebley et al., 1997)^[19].

tga1

The *teosinte glume architecture 1* (*tga1*) gene encodes an SBP-domain transcriptional regulator that liberated maize kernels from the hardened protective casing of wild teosinte. The functional divergence is driven by a single missense substitution in exon 1 that replaces a conserved lysine with an asparagine. This modification drastically reduces functional TGA1 protein abundance in domesticated maize compared to teosinte. While high teosinte TGA1 levels drive heavy lignification to form a stony fruitcase, the reduced protein in maize releases this suppression, producing small, loosely bound glumes that leave the kernels naked and exposed on the central cob (Wang et al., 2005)^[88].

Vgt1

Vegetative to generative transition 1 (*Vgt1*) is a pleiotropic quantitative trait locus controlling the timing of floral induction, leaf node formation, and plant height. It functions as a long-range, cis-acting regulatory element located upstream of *ZmRap2.7*, an AP2-like repressor of the vegetative-to-reproductive transition. The active wild-type *Vgt1* allele maintains elevated *ZmRap2.7* expression, delaying flowering to prolong vegetative growth. Conversely, domesticated early-flowering variants contain a transposon (MITE) insertion that severely dampens *ZmRap2.7* transcription. This releases the repressor, accelerating the floral transition to shorten overall stature and optimize synchronization for shorter growing seasons (Salvi et al., 2007)^[72].

ZmVPP1

The *ZmVPP1* locus is a primary determinant of drought tolerance and root system development in maize. It encodes a vacuolar H⁺ pyrophosphatase pump. In drought-tolerant genotypes, a naturally selected 366-base pair promoter insertion provides binding sites for MYB transcription factors. During dehydration, these factors drive strong stress-inducible upregulation of the *ZmVPP1* allele. The resulting abundance of H⁺ pumps actively translocates protons into the vacuole, enhancing cellular osmoregulation and turgor maintenance. This elevated electrochemical gradient significantly upgrades photosynthetic efficiency and stimulates vigorous root development, maximizing subterranean water foraging and seedling survival under prolonged drought (Wang et al., 2016)^[90].

crtRB1

The *crtRB1* gene is the principal quantitative trait locus dictating beta-carotene concentration in maize kernels, offering a vital tool for provitamin A biofortification. It encodes beta-carotene hydroxylase 1, an enzyme that typically degrades beta-carotene into downstream hydroxylated products with lower provitamin A activity. Favorable natural variants of this locus contain specific structural polymorphisms (insertions and deletions) that heavily depress *crtRB1* transcript expression exclusively within the endosperm. This stark downregulation minimizes beta-hydroxylase activity during grain filling, effectively blocking beta-carotene degradation and resulting in a dramatic accumulation of provitamin A within the mature kernel (Yan et al., 2010)^[95].

ZmMADS69

The *ZmMADS69* MADS-box transcription factor governs the vegetative-to-reproductive transition in maize, historically enabling its geographic expansion into high-latitude temperate zones by breaking strict tropical daylength requirements. It operates via a double-negative transcriptional regulatory circuit. The functional *ZmMADS69* protein directly downregulates the expression of the AP2-like floral repressor *ZmRap2.7* in leaf tissues. Relieving this chronic genetic block powerfully induces the expression of the principal mobile florigen signal, *ZCN8*. The abundant synthesis of *ZCN8* accelerates reproductive floral initiation, significantly advancing heading dates to prevent frost damage in short-season climates (Liang et al., 2019)^[48].

Fea2

The *FASCIATED EAR2* (*FEA2*) locus regulates inflorescence meristem proliferation and kernel row number (KRN) in maize. It encodes a leucine-rich repeat (LRR) transmembrane receptor that binds a mobile CLAVATA3-like signaling peptide. This ligand-receptor interaction normally initiates a negative feedback loop on the WUSCHEL transcription factor, strictly preventing stem cell over-proliferation. However, weak or null mutant alleles, such as *fea2-1328*, dampen this regulatory feedback. This allows the inflorescence meristem to expand its physical diameter, providing more space for additional spikelet meristem rows and ultimately yielding a higher number of kernel rows per ear without compromising structural integrity (Bommert et al., 2013)^[7].

UB2 and UB3: The *unbranched2* (*ub2*) and *unbranched3* (*ub3*) genes encode functionally redundant SBP-box transcription factors that critically regulate maize inflorescence architecture and yield potentials. Localized throughout the meristematic tissue (but absent from the central self-renewal domain), these proteins modulate the transition of indeterminate stem cells into specialized lateral organ primordia. By balancing the rate of meristem renewal against lateral organ initiation, UB2 and UB3 dictate kernel row number and total ear productivity. Their crucial role is underscored by double-mutant plants (*ub2;ub3*), which exhibit severe defects in tassel branching and ear development (Chuck et al., 2014)^[16].

d8: The *Dwarf8* (*d8*) locus is a pleiotropic gene regulating vegetative plant height and floral induction timing in maize, structurally orthologous to the loci that drove the "Green

Revolution." It encodes a nuclear-localized DELLA protein that acts as a negative regulator of the gibberellin response pathway, curbing cell elongation. Natural quantitative variations in *d8* are driven by combined non-synonymous polymorphisms, including a 6-base pair coding deletion near a binding-essential SH2-like domain and promoter MITE insertions. Rather than acting as null mutations, these variations generate subtle modifications in transcriptional interactivity, fine-tuning stalk height and shortening the lifecycle to maximize spatial adaptation (Thornberry et al., 2001) [83].

Wheat (*Triticum aestivum*)

The expansive hexaploid genome of wheat inherently features extensive redundancy, rendering master pleiotropic hubs vital for generating noticeable phenotypic shifts across multiple sub-genomes simultaneously.

NAM-B1

The *Gpc-B1* locus encodes NAM-B1, an NAC transcription factor that controls leaf senescence and nutrient remobilization in wheat. In wild-type plants, *NAM-B1* expression upregulates during grain maturation, accelerating senescence and driving the efficient translocation of nitrogen, zinc and iron from flag leaves into developing grains. Modern cultivated varieties often carry a non-functional frame-shift mutation that delays senescence but significantly lowers nutritional value. Restoring functional transcripts is essential for developing biofortified crops, as knocking down *NAM-B1* traps nutrients in vegetative tissues, dropping grain protein, zinc and iron concentrations by over 30% (Uauy et al., 2006) [86].

Lr34 / Yr18 / Pm38

The hexaploid wheat gene *Lr34* confers durable, partial "slow-rusting" resistance to multiple fungal pathogens, remaining effective for over 50 years without pathogens developing increased virulence. It encodes an ABC transporter resembling a pleiotropic drug resistance (PDR) protein. Resistant alleles feature specific sequence polymorphisms including a 3-bp deletion and a single-nucleotide polymorphism that alter the transporter's first transmembrane domain. *Lr34* is natively upregulated in adult flag leaves independent of pathogen presence, where it triggers localized senescence-like processes and may translocate defensive metabolites into the apoplast to restrict intercellular fungal expansion during the critical grain-filling stage (Krattinger et al., 2009) [45].

Rht-B1 and Rht-D1

The "Green Revolution" genes *Rht-B1* and *Rht-D1* confer a semi-dwarf wheat phenotype that dramatically increases grain yield and lodging resistance by optimizing the harvest index. The wild-type alleles encode nuclear transcription factors that act as growth repressors, which are normally degraded by the hormone gibberellin. However, semi-dominant dwarfing mutant alleles feature point mutations creating premature stop codons. Translational reinitiation produces truncated repressor proteins lacking the gibberellin-signaling domain. These constitutively active mutant repressors escape gibberellin-mediated degradation, continuously suppressing stem elongation to maintain a sturdy dwarf stature despite high endogenous hormone levels (Peng et al., 1999) [66].

Ppd-D1

The *Photoperiod-D1* (*Ppd-D1*) locus regulates flowering time and photoperiod sensitivity in wheat. In wild-type plants, a pseudo-response regulator (PRR) gene oscillates to restrict flowering under short days. However, the semi-dominant *Ppd-D1a* mutant contains a 2-kilobase promoter deletion that causes constant misexpression of the PRR gene. This overabundance continuously drives the expression of the floral integrator *Flowering Locus T* (*FT*), rendering the plant photoperiod-insensitive. By bypassing the natural long-day requirement, this mutation significantly accelerates heading, allowing crops to escape end-of-season heat and drought stress while increasing the harvest index and 1000-grain weight under moisture-limited conditions (Beales et al., 2007) [5].

Vrn1

The *VRN1* gene defines the primary genetic toggle separating winter and spring wheat classes by orchestrating the vegetative-to-reproductive phase transition. In winter cultivars, flowering is chronically blocked because the *VRN2* repressor suppresses *VRN1* transcription. Prolonged winter cold (vernalization) gradually downregulates *VRN2*, allowing the *VRN1* MADS-box transcription factor to accumulate and activate downstream meristem identity factors for spike differentiation. Conversely, dominant spring alleles contain promoter deletions that prevent *VRN2* binding. This entirely bypasses the cold requirement, driving constitutive, winter-independent flowering to stabilize yield potentials across variable regional climates (Yan et al., 2003) [96].

TaGW2

The *TaGW2* gene encodes an E3 ubiquitin ligase that acts as a critical negative regulator of grain development in bread wheat. Unlike in rice, allelic functional variations in wheat are driven by single-nucleotide polymorphisms (SNPs) within the promoter region that regulate transcription. The Hap-6A-G allele maintains high expression, strongly suppressing cellular expansion to produce slimmer, lower-weight grains. In contrast, the favorable Hap-6A-A allele exhibits downregulated promoter activity and reduced transcript accumulation. This suppression of the E3 ligase pathway relieves negative growth constraints, significantly expanding grain width, increasing one-thousand grain weight and slightly advancing crop maturity (Su et al., 2011).

2.4 Sorghum (*Sorghum bicolor*)

Sorghum utilizes specific pleiotropic hubs to maintain robust vegetative integrity and reproductive success under acute heat and arid conditions.

Mal

The *Mal* gene encodes the nuclear transcription factor SbPRR37, the primary repressor of sorghum flowering under long-day (LD) environments. Natural loss-of-function mutations disable this repressor to prompt early flowering, an essential genetic adaptation that enabled tropical short-day sorghum to expand into temperate zones. Conversely, functional alleles delay flowering under extended light, prolonging vegetative growth to maximize biomass for forage and bioenergy production.

Mechanistically, SbPRR37 integrates light signals with the circadian clock. Under long days, light-activated PRR37 peaks in both the morning and evening, upregulating the

floral inhibitor CONSTANS (CO) while suppressing key floral activators (Ehd1, FT and ZCN8) to block reproduction. Under short days, the evening expression phase shifts into darkness, preventing *SbPRR37* upregulation. This absence of PRR37 de-represses the floral activators, successfully triggering the switch to reproductive flowering (Murphy et al., 2011)^[64].

Dw3

The *brachytic2* (*br2*) locus in maize and its sorghum ortholog *dwarf3* (*dw3*) regulate stalk morphology to maximize structural lodging resistance. Both loci encode ATP-binding cassette (ABC) membrane proteins that modulate the downward polar transport of auxin. Loss-of-function variants severely disrupt this auxin flux, restricting cellular elongation in the lower stem internodes. This yields a highly desired, compact dwarf phenotype with an improved harvest index and enhanced mechanical strength, completely sparing reproductive and leaf development (Multani et al., 2003)^[63].

Bmr12

Brown midrib (*Bmr*) loci are major pleiotropic determinants of cell wall composition and forage digestibility in sorghum. The wild-type locus encodes caffeic acid O-methyltransferase (COMT), an enzyme essential for synthesizing syringyl (S) lignin. Induced mutations, such as *bmr12*, introduce premature stop codons that severely downregulate COMT transcripts. This enzymatic block reduces S-lignin and esterified p-coumaric acid levels, manifesting as a classic brown coloration of leaf midribs, decreasing cell wall rigidity to improve ruminant feeding efficiency and inducing delayed flowering for optimized seasonal biomass accumulation (Bout and Vermerris, 2003)^[10].

Tannin1

The *Tannin1* (*Tan1*) gene dictates grain chemical defense and seed quality in sorghum. The functional allele encodes a WD40 repeat domain protein that coordinates with transcription factors to activate the late flavonoid and proanthocyanidin biosynthetic pathways in the pigmented testa. This accumulation of condensed tannins produces dark-seeded lines with prominent resistance to birds and grain molds. Conversely, natural loss-of-function mutations (*tan1-a* and *tan1-b*) strip the WD40 protein of its functional domain, halting tannin deposition to yield bright, nontannin grains with dramatically increased protein digestibility and livestock feed efficiency (Wu et al., 2012)^[92].

2.5 Barley (*Hordeum vulgare*)

Breeding initiatives in barley rely on targeted pleiotropy to manipulate highly specific architectural features of the inflorescence.

Vrs1

The *vrs1* (*six-rowed spike 1*) gene is the primary genetic toggle that enabled early agriculturalists to triple barley yields by converting sterile, two-rowed wild progenitors into fully fertile six-rowed cultivars. The dominant wild-type *Vrs1* allele encodes a homeodomain-leucine zipper (HD-ZIP) I-class transcription factor expressed exclusively in the lateral spikelet primordia, where it actively represses normal organogenesis and renders the lateral rows sterile. The cultivated six-rowed phenotype is universally driven by

independent recessive *vrs1* loss-of-function mutations including single-nucleotide substitutions, premature stop codons and large genomic deletions that deactivate this repressor. This genetic de-repression restores full fertility and spikelet development in the lateral rows, a yield-enhancing trait that evolved repeatedly across different geographical zones (Komatsuda et al., 2007)^[42].

Ppd-H1

The *Photoperiod-H1* (*Ppd-H1*) locus is a master regulator that dictates when barley shifts from vegetative growth to reproductive spike formation. The dominant wild-type allele encodes a circadian-associated pseudo-response regulator (PRR) protein that accelerates flowering under long-day conditions by rhythmically modulating the expression of CONSTANS-like genes (*HvCO1* and *HvCO2*). This drives the upregulation of florigen mobile signals (*HvFT*) to initiate reproduction. Conversely, recessive *ppd-H1* loss-of-function variants delay flowering under long days due to altered *HvCO* rhythms and crippled *HvFT* synthesis. This elongation of the vegetative phase holds high agronomic value, allowing spring-sown cultivars to maximize biomass and secure stable grain yields (Turner et al., 2005)^[85].

Nud

The *Nud* (*nudum*) locus is the molecular toggle separating covered (hulled) and naked (hulless) barley. Agronomically, covered barley is ideal for malting and livestock feed, while naked variants thresh free, yielding high-fiber grains optimized for direct human consumption. The wild-type *Nud* allele encodes an ethylene response factor (ERF) transcription factor that upregulates localized lipid biosynthesis. These secreted lipids form an adhesive layer on the outer pericarp epidermis, permanently fusing the maturing seed to the surrounding floral hulls.

Conversely, the naked barley phenotype is driven by recessive *nud* loss-of-function mutations, most commonly a large 17-kilobase deletion that obliterates the ERF coding sequence. Without the functional NUD transcription factor, the lipid biosynthetic cascade remains inactive and the adhesive epidermal lipid layer fails to form. Lacking this crucial biochemical glue, organ fusion is prevented, permitting the hull to cleanly separate from the grain during threshing (Taketa et al., 2008)^[80].

HvCbf14

The C-repeat binding factor (Cbf) gene family acts as a master switch for cold acclimation in plants, providing a critical genetic pathway for developing freeze-resistant crops. When environmental temperatures drop, an upstream regulator (ICE) rapidly initiates the transcription of Cbf genes. The resulting CBF proteins, which are AP2/ERF transcriptional activators, migrate to the nucleus and bind to specific DNA motifs, such as C-repeat or dehydration-responsive elements in the promoters of downstream target loci.

This precise binding triggers a transcriptional cascade that upregulates cold-regulated (Cor) genes, which synthesize cryoprotective proteins and osmolytes to stabilize cellular membranes and prevent intracellular ice crystallization. Operating under a temperature-dependent and circadian-regulated diurnal rhythm, this complex signal transduction network translates chilling sensations into a robust

physiological defense, ultimately elevating freezing tolerance across the plant's tissues (Campoli *et al.*, 2009)^[13].

Future Directions and Perspectives

To guarantee future agricultural resilience, scientific focus must pivot from the rudimentary introgression of single-effect traits toward the highly precise, systems-level calibration of interconnected genetic networks.

Precision Tuning via Regulatory Haplotypes

Moving forward, crop engineering will rely less on binary gene knockouts and more on the sophisticated modulation of expression timing. By identifying and recombining specific cis-regulatory promoter variants derived from multi-parent panels, geneticists can achieve highly localized, conditional gene expression. This ensures that energy-intensive biofortification or defense cascades are initiated only during optimal developmental windows, thereby mitigating the severe metabolic penalties historically linked to multi-trait improvements.

Deployment of Breeder-Friendly Assays

Translating high-level genomic discoveries into applied agronomic progress requires the rapid development of accessible, high-throughput marker systems (e.g., KASP or localized SNP arrays) (Rasheed *et al.*, 2017)^[68]. These tools allow global breeding networks to aggressively track and introgress complex multi-trait loci without the prohibitive costs of continual field phenotyping.

Resolving Environmental Plasticity

As environmental variables fluctuate, the stability of epistatic interactions can falter. Forthcoming research must meticulously evaluate how pleiotropic networks react across highly varied geographic testing environments to guarantee that targeted yield enhancements do not collapse under unforeseen climatic interactions (El-Soda *et al.*, 2014)^[24].

Integration of Multi-Omics and Predictive AI

Unlocking the complete mechanistic depth of pleiotropy requires harmonizing vast datasets spanning transcriptomics, proteomics and metabolomics (Yang *et al.*, 2021)^[98]. Coupling this multi-tiered biological data with Machine Learning architectures will empower researchers to accurately forecast epistatic behavior and biochemical outcomes prior to executing costly, long-term physical field crosses.

Accelerated Breeding Frameworks

The immediate deployment of these highly refined pleiotropic hubs is being radically accelerated through controlled "Speed Breeding" methodologies. By artificially manipulating light and temperature cycles within specialized glasshouses, breeders can rapidly compress generation times, achieving up to six full life cycles of staple cereals per calendar year, dramatically accelerating the race against climate instability (Watson *et al.*, 2018)^[91].

References

1. Albert FW, Kruglyak L. The role of regulatory variation in complex traits as revealed by eQTL studies. *Nature Reviews Genetics*. 2015;16(4):197-212.
2. Antony G, Zhou J, Huang S, Li T, Liu B, White F, Yang B. Rice SWEET14 glucose transporter genes as targets for TAL effectors of *Xanthomonas oryzae* pv. *oryzae*. *Plant Cell*. 2010;22(11):3864-3876.
3. Bailey-Serres J, Parker JE, Ainsworth EA, Oldroyd GED, Schroeder JI. Genetic strategies for improving crop yields. *Nature*. 2019;575(7781):109-118.
4. Bandillo N, Raghavan C, Muyco PA, Sevilla MAL, Lobina IT, Dilla-Ermita CJ, Tung CW, McCouch S, Thomson M, Mauleon R, Singh RK, Craufurd P, Leung H, Lafitte R. Multi-parent advanced generation inter-cross (MAGIC) populations in rice: progress and potential for genetics research and breeding. *Rice*. 2013;6(1):11.
5. Beales J, Turner A, Griffiths S, Snape JW, Laurie DA. A pseudo-response regulator is misexpressed in the photoperiod insensitive Ppd-D1a mutant of wheat. *Theoretical and Applied Genetics*. 2007;115(5):721-733.
6. Bernardo R, Yu J. Prospects for genome-wide selection for quantitative traits in maize. *Crop Science*. 2007;47(3):1082-1090.
7. Bommert P, Nagasawa N, Jackson D. Quantitative variation in maize kernel row number is controlled by the FASCIATED EAR2 locus. *Nature Genetics*. 2013;45(3):334-338.
8. Bouis HE, Saltzman A. Improving nutrition through biofortification: a review of evidence from HarvestPlus, 2003 through 2016. *Global Food Security*. 2017;12:49-58.
9. Bouis HE, Welch RM. Biofortification a sustainable agricultural strategy for reducing micronutrient malnutrition in the global south. *Crop Science*. 2010;50(Suppl 1):S20-S32.
10. Bout S, Vermerris W. A candidate-gene approach to clone the sorghum Brown midrib gene encoding caffeic acid O-methyltransferase. *Molecular Genetics and Genomics*. 2003;269(2):205-214.
11. Cabot C, Martos S, Llugany M, Gallego B, Tolrà R, Poschenrieder C. Roles of zinc in plant tolerance to stress, nutrient-mediated immunity and biofortification. *Frontiers in Plant Science*. 2019;10:1147.
12. Cakmak I. Enrichment of cereal grains with zinc: Agronomic or genetic biofortification? *Plant and Soil*. 2008;302(1-2):1-17.
13. Campoli C, Matus-Cádiz MA, Pozniak CJ, Cattivelli L, Fowler DB. Comparative expression of Cbf genes in the Triticeae under different acclimation temperatures. *Molecular Genetics and Genomics*. 2009;282(2):141-152.
14. Cavanagh C, Morell M, Mackay I, Powell W. From concept to practice: unlocking semi-exotic germplasm with multi-parent advanced generation inter-cross (MAGIC) populations. *Crop Science*. 2008;48(1):S25-S40.
15. Chapin FS, Schulze ED, Mooney HA. The ecology and economics of storage in plants. *Annual Review of Ecology and Systematics*. 1990;21(1):423-447.
16. Chuck G, Brown PJ, Meeley R, Jackson D. The SBP-box transcription factor unbranched3 affects yield traits in maize. *Nature Genetics*. 2014;46(11):1214-1217.
17. Datnoff LE, Elmer WH, Huber DM. Mineral nutrition and plant disease. St Paul: American Phytopathological Society; 2007. p. 1-278.
18. Dell'Acqua M, Gatti DM, Pea G, Cattonaro F, Coppens F, Magris G, Hlaing AL, Aung HH, Nelissen H, Baute J, Frascaroli E, Churchill GA, Inzé D, Morgante M, Pè ME.

- Genetic properties of the MAGIC maize population: a new platform for high-resolution quantitative trait locus mapping. *Genome Biology*. 2015;16(1):167.
19. Doebley J, Stec A, Hubbard L. The evolution of apical dominance in maize. *Nature*. 1997;386(6624):485-488.
 20. Dubouzet JG, Sakuma Y, Ito Y, Kasuga M, Dubouzet EG, Miura S, Seki M, Shinozaki K, Yamaguchi-Shinozaki K. OsDREB genes in rice, *Oryza sativa* L., encode transcription activators that function in drought-, high-salt- and cold-responsive gene expression. *Plant Journal*. 2003;33(4):751-763.
 21. Dudley AM, Janse DM, Tanay A, Shamir R, Church GM. A quantitative screen for gene interaction in *Saccharomyces cerevisiae*. *Molecular Systems Biology*. 2005;1(1):2005.0001.
 22. Dwivedi SL, Heslop-Harrison P, Amas J, Ortiz R, Edwards D. Epistasis and pleiotropy-induced variation for plant breeding. *Plant Biotechnology Journal*. 2024;22(11):2788-2807.
 23. Eckardt NA, Ainsworth EA, Bahuguna RN, Broadley MR, Busch W, Carpita NC, Castrillo G, Chory J, DeHaan LR, Duarte CM, Henry A. Climate change challenges, plant science solutions. *Plant Cell*. 2023;35(1):24-66.
 24. El-Soda M, Malosetti M, Zwaan BJ, Koornneef M, Aarts MGM. Genotype × environment interaction QTL mapping in plants: lessons from *Arabidopsis*. *Trends in Plant Science*. 2014;19(6):390-398.
 25. Fan C, Xing Y, Mao H, Lu T, Han B, Xu C, Li X, Zhang Q. GS3, a major QTL for grain length and weight and minor QTL for grain width and thickness in rice, encodes a putative transmembrane protein. *Theoretical and Applied Genetics*. 2006;112(6):1164-1171.
 26. Foyer CH, Rasool B, Hunter A, Hancock RD. Is ascorbate peroxidase 1 a specific monitor of intracellular reactive oxygen species? *Plant Physiology*. 2016;171(4):2315-2317.
 27. Fridman E, Carrari F, Liu YS, Fernie AR, Zamir D. Zooming in on a quantitative trait for tomato yield using interspecific introgressions. *Science*. 2004;305(5691):1786-1789.
 28. Frisch M, Melchinger AE. Selection strategies for marker-assisted backcrossing. *Crop Science*. 2005;45(4):1231-1244.
 29. Fujita M, Fujita Y, Noutoshi Y, Takahashi F, Narusaka Y, Yamaguchi-Shinozaki K, Shinozaki K. Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Current Opinion in Plant Biology*. 2006;9(4):436-442.
 30. Garg M, Sharma N, Sharma S, Kapoor P, Kumar A, Chunduri V, Arora P. Biofortified crops generated by breeding, agronomy and transgenic approaches are yielding success in the field. *Frontiers in Chemistry*. 2018;6:462.
 31. Gill SS, Tuteja N. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiology and Biochemistry*. 2010;48(12):909-930.
 32. Hasanuzzaman M, Hossain MA, Silva JATD, Fujita M. Plant response and tolerance to abiotic oxidative stress: antioxidant defense is a key factor. In: *Crop stress and its management: Perspectives and strategies*. 2012. p. 261-315.
 33. Hawkesford MJ, Araus JL, Park R, Calderini D, Miralles D, Shen T, Zhang J, Parry MAJ. Prospects of macronutrient use efficiency improvements. In: *Plant macronutrient use efficiency*. 2013. p. 1-24.
 34. He X, Zhang J. Toward a molecular understanding of pleiotropy. *Genetics*. 2006;173(4):1885-1891.
 35. Horton P, Banwart SA, Brock D, Cleaver KJ, Campi Cuatínio A, Dorward PT, Griffiths H, Halford NG, Hawkesworth DL, Kemp S, Macaulay M, Scholes MC, Smith J, Stringer LC. Technologies to deliver food and nutritional security through agriculture. *Nature Plants*. 2018;4(5):250-255.
 36. Hospital F. Marker-assisted backcrossing. In: *Marker-assisted selection*. 2009. p. 137-159.
 37. Huang BE, George AW, Forrest KL, Kilian A, Hayden MJ, Morell MK, Cavanagh CR. MAGIC populations in crops: progress, challenges and opportunities. *Theoretical and Applied Genetics*. 2015;128(6):999-1017.
 38. Huang XY, Chao DY, Gao JP, Zhu MZ, Shi M, Lin HX. A previously unknown zinc finger protein, DST, regulates elevated stomatal density and drought tolerance in rice. *Genes & Development*. 2009;23(15):1805-1817.
 39. Husaini AM. High-value pleiotropic genes for developing multiple stress-tolerant biofortified crops for 21st-century challenges. *Heredity*. 2022;128(6):460-472.
 40. Ikeda A, Ueguchi-Tanaka M, Sonoda Y, Kitano H, Koshioka M, Futsuhara Y, Matsuoka M, Yamaguchi J. Slender rice, a constitutive gibberellic acid response mutant, is caused by a null mutation of the SLR1 gene, an ortholog of the DELLA protein. *Plant Cell*. 2001;13(5):999-1010.
 41. Jordan DM, Verbanck M, Do R. HOPS: a quantitative score reveals pervasive horizontal pleiotropy in human genetic variation is driven by extreme polygenicity of human traits and diseases. *Genome Biology*. 2019;20(1):22.
 42. Komatsuda T, Pourkheirandish M, He C, Azhaguvel P, Kanamori H, Perovic D, Stein N, Graner A, Wicker T, Tagiri A, Lundqvist U, Fujimura T, Matsuoka M, Matsumoto T, Yano M. Six-rowed barley originated from a mutation in a homeodomain-leucine zipper I-class homeobox gene. *Proceedings of the National Academy of Sciences*. 2007;104(4):1424-1429.
 43. Koo BH, Yoo SC, Park JW, Kwon CT, Lee BD, An G, Zhang Z, Li J, Li Z, Paek NC. OsPRR37 controls reproductive transit and flowering time in rice. *Plant Physiology*. 2013;163(1):478-491.
 44. Korte A, Vilhjálmsson BJ, Segura V, Platt A, Long Q, Nordborg M. A mixed-model approach for genome-wide association studies of correlated traits in structured populations. *Nature Genetics*. 2012;44(9):1066-1071.
 45. Krattinger SG, Lagudah ES, Spielmeier W, Singh RP, Huerta-Espino J, McFadden H, Bossolini E, Selter LL, Keller B. A putative ABC transporter confers durable resistance to multiple fungal pathogens in wheat. *Science*. 2009;323(5919):1360-1363.
 46. Kurowska M, Daszkowska-Golec A, Gruszka D, Marzec M, Szurman M, Szarejko I, Maluszynski M. TILLING - a strategy in plant functional genomics. *Scientific Research and Essays*. 2011;6(8):1617-1625.

47. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics*. 2008;9(1):559.
48. Liang Y, Liu Q, Wang X, Huang C, Xu G, Hey S, Lin H, Li C, Xu D, Wu L, Wang C, Wu W, Xia J, Han X, Lu S, Zhang J, Wu Y. ZmMADS69 functions as a flowering activator through the ZmRap2.7-ZCN8 regulatory module in maize. *Plant Cell*. 2019;31(10):2364-2373.
49. Liu R, Jia H, Cao X, Huang J, Li F, Tao Y, Qiu F, Zheng Y, Zhang Z. Fine mapping and candidate gene prediction of a pleiotropic quantitative trait locus for yield-related trait in *Zea mays*. *PLoS One*. 2012;7(11):e49836.
50. Ma JF, Tamai K, Yamaji N, Mitani N, Konishi S, Katsuhara M, Ishiguro M, Murata Y, Yano M. A silicon transporter in rice. *Nature*. 2006;440(7084):688-691.
51. Mackay TF, Stone EA, Ayroles JF. The genetics of quantitative traits: challenges and prospects. *Nature Reviews Genetics*. 2009;10(8):565-577.
52. Mao H, Sun S, Yao J, Wang C, Yu S, Xu C, Li X, Zhang Q. Linking a cell cycle regulator (GS3) to quantitative variation of grain size in rice. *Proceedings of the National Academy of Sciences*. 2010;107(45):19579-19584.
53. Mao L, Van Hemert JL, Dash S, Dickerson JA. Rice network biology: gene co-expression networks and beyond. *Rice*. 2009;2(4):161-177.
54. McCallum CM, Comai L, Greene EA, Henikoff S. Targeting Induced Local Lesions In Genomes (TILLING) for plant functional genomics. *Plant Physiology*. 2000;123(2):439-442.
55. McMullen MD, Kresovich S, Villeda HS, Bradbury P, Li H, Sun Q, Flint-Garcia S, Thornsberry J, Acharya C, Bottoms C, Brown P, Browne C, Eller M, Guill K, Harjes C, Kroon D, Lepak N, Mitchell SE, Peterson B, Pressoir G, Romero S, Oropeza RO, Salvo S, Settles H, Vergara WA, Villegas WA, Ward W, Holland JB, Gleason C, Doebley JF, Buckler ES. Genetic properties of the maize nested association mapping population. *Science*. 2009;325(5941):737-740.
56. Mertz ET, Bates LS, Nelson OE. Mutant gene that changes protein composition and increases lysine content of maize endosperm. *Science*. 1964;145(3629):279-280.
57. Meuwissen TH, Hayes BJ, Goddard ME. Prediction of total genetic value using genome-wide dense marker maps. *Genetics*. 2001;157(4):1819-1829.
58. Mickelbart MV, Hasegawa PM, Bailey-Serres J. Genetic mechanisms of abiotic stress tolerance that translate to crop yield stability. *Nature Reviews Genetics*. 2015;16(4):237-251.
59. Mitani-Ueno N, Yamaji N, Ma JF. Functional characterization of OsLsi1 as an arsenite channel in rice roots. *Plant and Cell Physiology*. 2011;52(3):512-519.
60. Mittler R. ROS are good. *Trends in Plant Science*. 2017;22(1):11-19.
61. Mittler R, Vanderauwera S, Gollery M, Van Breusegem F. Reactive oxygen species gene network of plants. *Trends in Plant Science*. 2004;9(10):490-498.
62. Miura K, Ikeda M, Matsubara A, Song XJ, Ito M, Asano K, Matsuoka M, Kitano H, Ashikari M. OsSPL14 promotes panicle branching and higher grain productivity in rice. *Nature Genetics*. 2010;42(6):545-549.
63. Multani DS, Briggs SP, Chamberlin MA, Blakeslee JJ, Murphy AS, Johal GS. Loss of an MDR transporter in compact stalks of maize br2 and sorghum dw3 mutants. *Science*. 2003;302(5642):81-84.
64. Murphy RL, Klein RR, Morishige DT, Brady JA, Rooney WL, Miller FR, Dugas DV, Klein PE, Mullet JE. Coincident light and clock regulation of pseudoreponse regulator protein 37 (PRR37) controls photoperiodic floral induction in sorghum. *Proceedings of the National Academy of Sciences*. 2011;108(40):16469-16474.
65. Oliva R, Ji C, Atienza-Grande G, Huguet-Tapia JC, Perez-Quintero A, Li T, Eom JS, Li C, Nguyen H, Liu B, Auguy F, Sciallano C, Luu VT, Dossa GS, Cunnac S, Schmidt SM, Slamet-Loedin IH, Vera Cruz C, Szurek B, Frommer WB, White FF, Yang B. Broad-spectrum bacterial blight resistance in rice mediated by natural promoter variants of SWEET genes. *Nature Biotechnology*. 2019;37(11):1338-1344.
66. Peng J, Richards DE, Hartley NM, Murphy GP, Devos KM, Flintham JE, Beales J, Fish LJ, Worland AJ, Pelica F, Sudhakar D, Christou P, Snape JW, Gale MD, Harberd NP. 'Green revolution' genes encode mutant gibberellin response modulators. *Nature*. 1999;400(6741):256-261.
67. Raboy V. Seeds for a better future: 'low phytate' mutant crops. *Trends in Plant Science*. 2001;6(10):458-462.
68. Rasheed A, Hao Y, Xia X, Balesdent MH, Tuberosa R, Edwards KJ, Isaac P, Appels R, Schnable PS, Sorrells ME, He Z. Crop breeding chips and genotyping platforms: progress, challenges and perspectives. *Molecular Plant*. 2017;10(8):1047-1064.
69. Ren P, Zhao D, Zeng Z, Yan X, Zhao Y, Lan C, Wang C. Pleiotropic effect analysis and marker development for grain zinc and iron concentrations in spring wheat. *Research Square Platform LLC*. 2022:1-20.
70. Saltzman A, Birol E, Bouis HE, Boy E, De Moura FF, Islam Y, Pfeiffer WH. Biofortification: progress toward a more nourishing future. *Global Food Security*. 2013;2(1):9-17.
71. Salvi S, Tuberosa R. To clone or not to clone plant QTLs: realities and prospects. *Current Opinion in Plant Biology*. 2005;8(2):113-124.
72. Salvi S, Sponza G, Morgante M, Tomes D, Niu X, Fengler KA, Meeley R, Ananiev EV, Svtashev S, Bruggemann E, Li B, Hainey CF, Radovic S, Zaina G, Rafalski JA, Tingey SV, Miao GH, Phillips RL, Tuberosa R. Conserved noncoding genomic sequences associated with a flowering-time quantitative trait locus in maize. *Proceedings of the National Academy of Sciences*. 2007;104(27):11376-11381.
73. Shahzad L, Waheed A, Sharif F, Ghafoor GZ, Rafique A. Understanding role of climatic parameters and adaptation strategies in agriculture productivity of South Asian countries. *Sustainable Environment*. 2024;10(1):2345453.
74. Singh NR. Pleiotropy vs. Polygenic Inheritance. *LibreTexts Genetics: Introduction to Genetics*. 2021:1.
75. Solovieff N, Cotsapas C, Lee PH, Purcell SM, Smoller JW. Pleiotropy in complex traits: challenges and strategies. *Nature Reviews Genetics*. 2013;14(7):483-495.
76. Song XJ, Huang W, Shi M, Zhu MZ, Lin HX. A QTL for rice grain width and weight encodes a previously

- unknown RING-type E3 ubiquitin ligase. *Nature Genetics*. 2007;39(5):623-630.
77. Spindel J, Begum H, Akdemir D, Virk P, Collard B, Redoña E, Atlin G, Jannink JL, McCouch SR. Genomic selection and association mapping in rice breeding populations: utilization of linkage disequilibrium and its property. *PLoS Genetics*. 2015;11(2):e1004982.
 78. Stephens M. A unified framework for association analysis with multiple related phenotypes. *PLoS One*. 2013;8(7):e65245.
 79. Su Z, Hao C, Wang L, Dong Y, Zhang X. Identification and development of a functional marker of TaGW2 associated with grain weight in bread wheat. *Theoretical and Applied Genetics*. 2011;122(1):211-223.
 80. Taketa S, Amano S, Tsujino Y, Sato T, Saisho D, Kakeda K, Nomura M, Suzuki T, Matsumoto T, Sato K, Kanamori H, Kawasaki S, Takeda K. Barley grain with adhering hulls is controlled by an ERF family transcription factor gene regulating a lipid biosynthesis pathway. *Proceedings of the National Academy of Sciences*. 2008;105(10):4062-4067.
 81. Tanksley SD, McCouch SR. Seed banks and molecular maps: unlocking genetic potential from the wild. *Science*. 1997;277(5329):1063-1066.
 82. Tester M, Langridge P. Breeding technologies to increase crop production in a changing world. *Science*. 2010;327(5967):818-822.
 83. Thornsberry JM, Goodman MM, Doebley J, Kresovich S, Nielsen D, Buckler ES. Dwarf8 polymorphisms associate with variation in flowering time. *Nature Genetics*. 2001;28(3):286-289.
 84. Tian F, Bradbury PJ, Brown PJ, Hung H, Sun Q, Flint-Garcia S, Rocheford TR, McMullen MD, Holland JB, Buckler ES. Genome-wide association study of leaf architecture in the maize nested association mapping population. *Nature Genetics*. 2011;43(2):159-162.
 85. Turner A, Beales J, Faure S, Dunford RP, Laurie DA. The pseudo-response regulator Ppd-H1 provides adaptation to photoperiod in barley. *Science*. 2005;310(5750):1031-1034.
 86. Uauy C, Distelfeld A, Fahima T, Blechl A, Dubcovsky J. A NAC gene regulating senescence improves grain protein, zinc and iron content in wheat. *Science*. 2006;313(5791):1698-1701.
 87. Uga Y, Sugimoto K, Ogawa S, Rane J, Ishitani M, Hara N, Hasegawa Y, Yoshiharu Y, Kano-Nakata M, Maruyama M, Chen J, Bheemanahalli R, Menguer PK, Ishiyama K, Yano M, Matsuoka M. Control of root system architecture by DEEPER ROOTING 1 increases rice yield under drought conditions. *Nature Genetics*. 2013;45(9):1097-1102.
 88. Wang H, Nussbaum-Wagler T, Li B, Zhao Q, Vigouroux Y, Faller M, Bomblies K, Lukens L, Doebley J. The origin of the naked grains of maize. *Nature*. 2005;436(7051):714-719.
 89. Wang J, Zhou L, Shi H, Chern M, Yu H, Yi H, He M, Yin J, Zhu X, Li Y, Li W, Liu J, Wang J, Chen X, Qing H, Wang Y, Liu G, Wang W, Li P, Wu X, Zhu L, Croft L, Ronald PC, Li S, Chen X. A single transcription factor promotes both yield and immunity in rice. *Science*. 2018;361(6406):1026-1028.
 90. Wang X, Wang H, Liu S, Ferjani A, Li J, Yan J, Yang X, Qin F. Genetic variation in ZmVPP1 contributes to drought tolerance in maize seedlings. *Nature Genetics*. 2016;48(10):1233-1241.
 91. Watson A, Ghosh S, Williams MJ, Cuddy WS, Simmonds J, Rey MD, Asyraf Md Hatta M, Hinchliffe A, Steed A, Reynolds D, Adamski NM, Breakspear A, Korolev A, Rayner T, Dixon LE, Riaz A, Martin W, Ryan M, Edwards D, Batley J, Raman H, Carter J, Stephens C, Tomkins PJ, Boulton MI, Macaulay M, Morris J, Hayes BJ, Michelmore R, Lister C, Shen Q, Jones JDG, Robinson S, Trick M, Wallington EJ, Werner P, Goltsman E, Wulff BBH, Hickey LT. Speed breeding is a powerful tool to accelerate crop research and breeding. *Nature Plants*. 2018;4(1):23-29.
 92. Wu Y, Li X, Xiang W, Zhu C, Lin Z, Wu Y, Li J, Pandravada S, Ridder DD, Bai G, Wang ML, Trick HN, Bean SR, Tuinstra MR, Tesso TT, Yu J. Allelic variation of Tannin1 controls tannin content in sorghum. *Proceedings of the National Academy of Sciences*. 2012;109(26):10281-10286.
 93. Xue W, Xing Y, Weng X, Zhao Y, Tang W, Wang L, Zhou H, Yu S, Xu C, Li X, Zhang Q. Natural variation in Ghd7 is an important regulator of heading date and yield potential in rice. *Nature Genetics*. 2008;40(6):761-767.
 94. Yadava DK, Hossain F, Mohapatra T. Nutritional security through crop biofortification in India: Status & future prospects. *Indian Journal of Medical Research*. 2018;148(5):621-631.
 95. Yan J, Kandianis CB, Harjes CE, Bai L, Kim EH, Yang X, Skinner DJ, Fu Z, Mitchell S, Li Q, Fernandez MGS, Zaharieva M, Babu R, Fu Y, Palacios N, Li J, DellaPenna D, Brutnell T, Buckler ES, Warburton ML, Rocheford T. Rare genetic variation at *Zea mays* crtRB1 increases β -carotene in maize grain. *Nature Genetics*. 2010;42(4):322-327.
 96. Yan L, Loukoianov A, Tranquilli G, Helguera M, Fahima T, Dubcovsky J. Positional cloning of the wheat vernalization gene VRN1. *Proceedings of the National Academy of Sciences*. 2003;100(10):6263-6268.
 97. Yan WH, Wang P, Chen HX, Zhou HJ, Li QP, Wang CR, Ding ZH, Zhang YS, Yu SB, Xing YZ, Zhang QF. A major QTL, Ghd8, plays pleiotropic roles in regulating grain productivity, plant height and heading date in rice. *Molecular Plant*. 2011;4(2):319-330.
 98. Yang Y, Saand MA, Huang L, Abdelaal WB, Zhang J, Wu Y, Li J, Sirohi MH, Wang F. Applications of multi-omics technologies for crop improvement. *Frontiers in Plant Science*. 2021;12:563953.
 99. Yu J, Holland JB, McMullen MD, Buckler ES. Genetic design and statistical power of nested association mapping in maize. *Genetics*. 2008;178(1):539-551.
 100. Zhou X, Stephens M. Efficient multivariate linear mixed model algorithms for genome-wide association studies. *Nature Methods*. 2014;11(4):407-409.